

DISEASES OF THE HEART

BY THE SAME AUTHOR

CLINICAL DISORDERS OF THE HEART BEAT

CLINICAL ELECTROCARDIOGRAPHY

THE MECHANISM AND GRAPHIC REGISTRATION
OF THE HEART BEAT

LECTURES ON THE HEART

BLOOD VESSELS OF THE HUMAN SKIN AND
THEIR RESPONSES

DEPARTMENT OF CLINICAL RESEARCH,
UNIVERSITY COLLEGE HOSPITAL, LONDON

DISEASES OF THE HEART

DESCRIBED FOR
PRACTITIONERS AND STUDENTS

By SIR THOMAS LEWIS
CBE, FRS, MD., DSc, LL.D., FRCP.

PHYSICIAN IN CHARGE OF DEPARTMENT OF CLINICAL RESEARCH UNIVERSITY COLLEGE
HOSPITAL LONDON, PHYSICIAN OF THE STAFF OF THE MEDICAL RESEARCH COUNCIL
HONORARY CONSULTING PHYSICIAN TO THE MINISTRY OF PENSIONS, CONSULTING
PHYSICIAN CITY OF LONDON HOSPITAL FELLOW OF UNIVERSITY COLLEGE LONDON

MACMILLAN AND CO., LIMITED
ST MARTIN'S STREET, LONDON

1933

COPYRIGHT

First Published on January 1933
Reprinted September 1933

PRINTED IN GREAT BRITAIN
BY R & R CLARK LIMITED EDINBURGH

PREFACE

If I had set out to describe the many and often very complex methods of examination now in use, had attempted to discuss at any length the pathology of heart disease, had included all known disorders of the heart, giving to each of these the space commensurate with our knowledge of it, and without consideration of its commonness or rarity, this book would have grown to form an unwieldy treatise. The impulse to write a work of reference has not stirred me, but I have had the desire to place at the disposal of students and medical practitioners the outline of my clinical teaching on diseases of the heart, as this has developed in my talks to my own hospital students.

In beginning that teaching twenty and more years ago, I determined that the basis of what I taught should be that which I myself had seen and proved to be true. A second ideal that I have striven hard to attain is *simplicity in teaching*, this was not so easy during the years in which my work lay largely amongst pulse and galvanometric curves. But I have become more and more convinced of the need for simplicity, and it was apparent that little of the graphic work could profitably be taught to men entering practice, whose foremost interest must always be in observations they themselves can make upon their patients.

Such a vast amount of knowledge has accumulated about human disease, that we can now cut ruthlessly away from text books all that we perceive to rest upon an insecure basis, and we can dispense with many trivial details, having something more substantial to put in their places. Some space has been regretfully given to the explanation of such pruning. Otherwise I have endeavoured to keep to what practice needs and to discuss such things thoroughly. I have tried to strip my subject of intricacies and redundances, of unnecessary technical terms, named signs, and the old trite phrases, for these begin to stifle Medicine. It has been important to try to achieve a proper perspective of values so as not to place undue

weight on this or that because its novelty attracts or because it has a strong personal interest I shall not be accused I think of over emphasising electrocardiography The few graphic records published suffice only to show the chief directions in which these records prove serviceable Radiography I have treated similarly by using illustrative orthodiagrams chosen from a large collection of my own taking In the book chief emphasis has fallen upon physical signs that are within the immediate reach of all I have laid deliberate stress on common forms of disease but here and there use the rare when it illuminates

What is said I have tried to arrange whenever possible in a natural and logical sequence of ideas and not simply as a series of disconnected facts There is abundant opportunity in the study of patients for the exercise of thought and no system of teaching that relies chiefly upon memory or neglects to inculcate simple processes of reasoning can have real educational merit A chief reform needed in medical education to day is that students should be encouraged to take a more leisurely and intelligent interest in phenomena not of the laboratory with which they will soon lose all connection but of everyday practice An underlying philosophy when it can be found is invaluable in practice not only because it quickens and maintains interest but because it forms a stable guide to action when experience fails as it often will in face of the unusual to give precise or particular direction I have tried and perhaps not always with success to follow and obey the recognised law that there is but one cause for one effect and to group symptoms where they belong so that the same group is not accredited to two distinct lesions but to one common state

And now to consider the arrangement of the subject matter It is traditional to classify diseases whenever possible upon an anatomical basis My first reaction to an anatomical classification of heart disease is that heart diseases cannot be classified Medicine is top heavy with classifications which often illogical parody those of the systematic sciences My second reaction is that anatomy fails as a chief basis of thought where heart disease is concerned When Surgery classifies and likewise thinks anatomically it has at least the excuse that it is so impelled because anatomy rigidly governs treatment of the diseases falling within its sphere Medicine has the privilege of greater freedom can view the body more as one whole see disease more broadly and picture a series of states of disordered function recognised only during life No less of heart disease cannot

be cast in anatomical moulds, and the persevering efforts so to fashion them have provided, and continue still to provide in our text books images of disease that are often so grotesque in their representation that they largely fail in their excellent purpose of forming guides to the management of disease. This is said in deliberately vigorous terms, that it may draw and hold attention. It is with the symptoms of disease that the patient, and that the doctor mainly, contends, and the symptoms of heart disease may be said to derive almost exclusively from faults in function. Therefore, in managing our patients, our thoughts must be chiefly set in terms of function and not of structure. To whom I fail to teach this first simple, but essential, lesson I have nought to teach. Who consents to it provisionally or who grasps it will be encouraged to pursue the idea by this book's arrangement. For it is with this object in mind that I have departed from past precedents and have placed in the foreground those things an understanding of which is of supreme importance in the management of heart cases, namely, cardiac failure and angina pectoris. The book differs much in its arrangement and in its outlook from other works dealing with affections of the heart but that is perhaps the soundest reason for its publication.

It is a pleasure to state my indebtedness to those who have helped me in making this book. I am especially grateful to my colleagues, Professor T. R. Elliott and Dr R. T. Grant, who have so carefully read and criticised the complete manuscript. Dr Grant has also aided much in checking the proof sheets. Dr B. E. Schlesinger has kindly read for me the chapter on Rheumatic Carditis. To Dr

graphs of the text of Chapters II and VI. In writing of disorderly heart action I have adopted sometimes a form of description similar to but briefer than that of *Clinical Disorders of the Heart Beat*, I do so with the sanction of Messrs Shaw & Sons who publish that book for me. My grateful thanks are due to my secretary, Miss J. M. Searle upon whom the arduous task of making accurate copies in type has mainly fallen and who has given me other and considerable help.

July 1932

CONTENTS

CHAPTER I

CARDIAC FAILURE BREATHLESSNESS

	PAGE
THE CENTRAL PROBLEM	1
CARDIAC FAILURE (GENERAL STATEMENT)	1
BREATHLESSNESS ON EFFORT	3
Simple Test	4
Strenuous Test	5

CHAPTER II

CARDIAC FAILURE WITH VENOUS CONGESTION

SYMPTOMS	7
Breathlessness	7
Fatigue and Exhaustion	7
Fulness of Head	7
Pain	7
Cough	7
Vomiting	8
SIGNS	8
Measures of Venous Pressure	8
Direct Method and Principle	8
Venous Swelling and Collapse	9
Venous Pulse as Index	12
Other Signs of Congestion	14
The Liver	14
The Kidneys	15
The Lungs	15
Cyanosis and Oedema	15
Malnutrition	15
Delirium	15
Diagnosis of Congestion Correlation of Signs	15
Discrepancies	16

CHAPTER III

CARDIAC OEDEMA

FACTORS IN CAUSATION OF DROPSY	18
RECOGNITION	19

	PAGE
DISTRIBUTION	19
DIFFERENTIAL DIAGNOSIS	20
EXUDATES INTO SEROUS CAVITIES	21

CHAPTER IV

FAILURE WITH CONGESTION (CAUSE, PROGNOSIS,
AND TREATMENT)

CAUSE OF CARDIAC FAILURE	22
PROGNOSIS	24
TREATMENT	25
Severe Congestion	26
Rest	26
Posture	26
Sleep	27
Diet	27
Venesection	28
Oxygen	29
Cardiac Tonics and Stimulants	29
Pain	29
Cough	30
Oedema Drainage	30
Cases presenting Auricular Fibrillation	31
After care	32
Return to Exercise, Supervision	33
Refractory Cases	34

CHAPTER V

SKIN COLOUR AND CIRCULATORY RATE

PALLOR	35
CYANOSIS AND ITS INTERPRETATION	36
Cause of Cyanosis	36
Cyanosis arising Centrally	36
Cyanosis arising Peripherally	37
Cyanosis of Failure with Congestion	38
Malar Flush	39
CUTANEOUS BLOOD FLOW	39
Capillary Pulsation	40

CHAPTER VI

CARDIAC ISCHAEMIA CORONARY THROMBOSIS

CORONARY ARTERIES AND CARDIAC ISCHAEMIA	41
The Coronary Arteries	41
Pathological Anatomy	41
Anginal Pain and Muscular Ischaemia	42

CONTENTS

xi
PAGE

CORONARY THROMBOSIS	43
Pathology	43
Symptoms and Signs	44
Death	44
Pain and Collapse	44
Manifestations of Cardiac Failure	45
Delayed Signs (fever Leucocytosis Ruptured Heart, Embolism and Friction)	46
Diagnosis	47
Course and Prognosis	48
Treatment	49

CHAPTER VII

ANGINA PECTORIS

HEBERDEN'S ANGINA PECTORIS ANGINA OF EFFORT	50
Clinical Associations	50
Symptoms	50
Diagnosis	52
A SPECIAL ANGINAL SYNDROME	53
Diagnosis	54
COURSE AND PROGNOSIS	54
TREATMENT	55
Rest and Exercise	56
Amyl Nitrite	57
Nitroglycerine	58
Theobromine	58
Ammonium Bromide	58
Luminal	58
Surgical Interference	58
SPECIAL CASES	59
Neurasthenia	59
Tobacco and Angina	59
" "	60
" "	60
" "	60

CHAPTER VIII

MINOR PULSE IRREGULARITIES

SINUS ARRHYTHMIA	62
INTERMITTENCE COUPLING AND GROUP BEATING	63
Intermittent Pulse	63
Coupling and Group Beating	65
Palpitation from Extrasystoles	66
Associations of Extrasystoles	66
Prognostic Significance of Extrasystoles	66
Treatment of Extrasystolic Palpitation	68

CHAPTER IX TACHYCARDIAS

	PAGE
SIMPLE TACHYCARDIA	69
Symptoms	70
Signs Associated	70
PAROXYSMAL TACHYCARDIA	71
Clinical Associations	71
Symptoms	72
Cardiac Reactions	72
AURICULAR FLUTTER	73
DIAGNOSIS AND DIFFERENTIATION OF TACHYCARDIAS	73
Flutter	75
PROGNOSIS	76
Simple Tachycardia	76
Paroxysmal Tachycardia	76
Auricular Flutter	78
TREATMENT	78
Simple Tachycardia	78
Paroxysmal Tachycardia	78
Auricular Flutter	79

CHAPTER X

AURICULAR FIBRILLATION (IRREGULAR TACHYCARDIA)

INTRODUCTORY	81
Clinical Associations	81
Paroxysmal Fibrillation	82
SYMPTOMATOLOGY	82
CARDIAC REACTIONS	82
RECOGNITION OF AURICULAR FIBRILLATION	83
PROGNOSIS	85
TREATMENT	86
Digitalis Therapy	86
Quinidine Therapy	90

CHAPTER XI

BRADYCARDIA SYNCOPE AND SUDDEN DEATH

BRADYCARDIA	92
Simple Bradycardia	92
Heart block	92
Differentiation of Bradycardias	94
Prognosis and Treatment	95
Simple Bradycardia	95
Heart block	95
SYNCOPE AND RELATED PHENOMENA	96
Postural Faintness	97
Vasovagal Attacks	98

CONTENTS

XIII
PAGE

Cardiac Syncope	99
Ventricular Arrest	99
Rapid Action of the Ventricles	101
Differential Diagnosis of Fainting	101
Prognosis and Treatment	102
Postural Giddiness	102
Vasovagal Attacks	102
Ventricular Arrest	103
Sudden Death Ventricular Fibrillation	103

CHAPTER VII

CARDIAC ENLARGEMENT

CAUSES OF PATHOLOGICAL DILATATION AND HYPERTROPHY	105
Cause of Dilatation	105
Cause of Hypertrophy	106
SYMPTOMS	106
MEASURING THE HEART'S SIZE IN LIFE	107
The Orthodiagraph	107
The Maximal Impulse	107
Impulse in Children and Growing Lads	109
Diffusion of the Impulse	109
Movement and Prominence of Ribs or Sternum	109
Epigastric Pulsation	110
Percussion	110
ACUTE DILATATION	112
RIGHT AND LEFT HYPERTROPHY	112
GREAT DILATATION OF LEFT AURICLE	112
DISTINCTION BETWEEN HYPERTROPHY AND DILATATION IN CHRONIC	
CASES	113
COMPENSATION	114
DEGREES OF CHRONIC ENLARGEMENT	116
PROGNOSIS	117
Dilatation	117
Enlargement	117
TREATMENT	118

CHAPTER VIII

DISEASE OF THE AORTIC VALVE

PATHOLOGICAL ANATOMY	120
AORTIC REGURGITATION	120
Symptoms	121
Signs	121
Water hammer Pulse	121
Diastolic Murmur	123
Differential Diagnosis including Pulmonary Regurgitation	124
The Pulse	124
The Murmur	124
Ruptured Aortic Cusp	125

	PAGE
Consecutive Phenomena	125
Rapid Pulse	125
Vasodilatation	125
Enlargement	126
Anginal Pain	126
AORTIC STENOSIS	127
Symptoms, Signs, and Diagnosis	127
Anacrotic Pulse	127
Systolic Thrill	128
Systolic Murmur	128
Presence of Aortic Regurgitation	128

CHAPTER XIV

DISEASE OF THE ATRICULO VENTRICULAR VALVES

PATHOLOGICAL ANATOMY	129
Mitral Valve	129
Tricuspid Valve	130
MITRAL STENOSIS	130
Its Recognition	130
With Normal Rhythm	131
Early Diagnosis	132
In Auricular Fibrillation	134
First S Murmur	136
MITRAL REGURGITATION	136
Systolic Apical Murmurs	136
Cardiorespiratory Murmur	136
Isocardial Murmurs	137
Inconstant Murmurs	137
Constant Murmurs	137
Mitral Regurgitation and its Sign	138
RECOGNITION OF MITRAL DISEASE	139
TRICUSPID AFFECTIONS	141
Tricuspid Stenosis	141
Tricuspid Regurgitation	141

CHAPTER XV

SIGNIFICANCE OF VALVULAR DISEASE

PROGNOSIS	143
Mitral Regurgitation	146
MANAGEMENT	148

CHAPTER XVI

HEART STRAIN, WORK, AND FAILURE

HEART STRAIN	151
The Burden	151
Acute Strain Dilatation	151
PROLONGED OVERTWORK AND HEART FAILURE	154

CHAPTER XVII

EFFORT SYNDROME BREATHLESSNESS OF EFFORT

	PAGE
EFFORT SYNDROME (ATHLETE'S HEART SOLDIER'S HEART)	158
Symptoms	159
Signs	160
Significance of the Syndrome	161
Prognosis	162
Management	162
DIFFERENTIATION OF BREATHLESSNESS	163

CHAPTER XVIII

PULMONARY CONGESTION, OEDOMA AND INFARCTION

CHRONIC PASSIVE CONGESTION AND EARLY OEDOMA OF LUNGS	165
Symptoms	165
Signs	166
ACUTE CONGESTION AND OEDOMA OF LUNGS	166
Acute Congestion with Early Oedema (Cardiac Asthma)	166
The Attack	166
Diagnosis	167
Acute Pulmonary Oedema	168
Prognosis	169
Treatment	169
PULMONARY INFARCTION	170
Symptoms Signs and Course	170
Treatment	171
HAEMOPTYSIS	171

CHAPTER XIX

PERICARDITIS

ACUTE PERICARDITIS AND PERICARDIAL EFFUSION	172
Pathology	172
Symptoms	173
Signs	173
Cardiac Impulse	173
Distension of Veins	173
Cardiac Dulness	174
Friction Sounds	174
Basal Signs	175
X ray Signs	175
Diagnosis of Effusion	175
Course and Treatment	175
Pain	176
Paracentesis	176
ADHERENT PERICARDIUM AND MEDIASTINOPERICARDITIS	176
Pathology	176
Symptoms and signs of an Obliterated Sac	177

Symptoms and Signs of Mediastinopericarditis and of Constrictive Pericarditis	177
Systolic Retractions	177
Immobilized Impulse	178
Dilatation of Veins	179
Inspiratory Swelling of Veins	179
Inspiratory Decline or Failure of Pulse	179
Absolute Cardiac Dulness fixed	180
Pleuropericardial Crepitus	180
X-ray Signs	181
Associated Phenomena	181
Diagnosis	181
Significance and Prognosis	181
Special Treatment	182

CHAPTER XX

BACTERIAL ENDOCARDITIS

SUBACUTE BACTERIAL ENDOCARDITIS	183
Pathology	183
Chief Clinical Manifestations	184
Onset	184
Local Signs	185
Symptoms and Signs of Infection	185
Embolic Accidents	188
Diagnosis	189
Course and Prognosis	190
Treatment	191
ACUTE BACTERIAL ENDOCARDITIS	191
Pathology	191
Chief Clinical Manifestations	192
Onset	192
Local Signs	192
Symptoms and Signs of Infection	192
Embolic Accidents	192
Diagnosis	193
Course Prognosis and Treatment	193

CHAPTER XXI

RHEUMATIC CARDITIS

INTRODUCTORY	194
RHEUMATISM OF CHILDHOOD	195
Pathological Anatomy	195
Symptoms and Signs	196
Onset	196
Pulse and Fever	196
Joints	197
Carditis and its Manifestations	197

CONTENTS

xvii

Nodules	198
Chorea	198
Course and Prognosis	198
Treatment	201

CHAPTER XXII

CHRONIC RHEUMATIC HEART DISEASE

CHIEF MANIFESTATIONS	202
Valve Disease	202
Pericardial Adhesions	203
Enlargement	203
Breathlessness and Failure	204
Anginal Pain	205
Precordial Ache	205
Palpitation	205
Auricular Fibrillation	205
Paroxysmal Tachycardia and Auricular Flutter	206
Emboic and Thrombotic Accidents	206
Acute Pulmonary Oedema	207
OUTLINE OF TYPE TREATMENT AND PROGNOSIS	207
Infection	210
Degenerative Changes	210

CHAPTER XXIII

SYPHILIS OF HEART AND AORTA

PATHOLOGICAL ANATOMY	211
DILATATION AND ANEURYSM OF AORTA	213
Symptoms	213
Local Signs	213
Pulsations	213
Pulsating Tumour	214
Tracheal Tug	215
Dullness	215
Heart Sounds	215
X ray Shadow	216
Symptoms and Signs of Pressure	217
Descending Aorta	220
Distant Arterial Signs	220
CARDIOAORTIC SYPHILIS	221
Chief Manifestations	222
Dilatation of the Aorta	222
Aortic Regurgitation	222
Angina Pectoris	222
Cardiac Failure	222
Types	223
Diagnosis	223
Course and Prognosis	224
Treatment	225

CHAPTER XXIV

ESSENTIAL HYPERTENSION

PATHOLOGICAL ANATOMY	PAGE
SYMPTOMS	227
RECOGNITION OF HIGH BLOOD PRESSURE	228
Accentuation of the 2nd Sound	229
Pulse Tension	230
Sphygmomanometry	230
RECOGNITION OF ARTERIOLAR SPASM	231
ACCOMPANYING MANIFESTATIONS	231
Vascular Manifestations	231
Cardiac Manifestations	232
Renal Manifestations	232
Inflammation	233
A FINAL STATE	233
DIFFERENTIATION OF ESSENTIAL HYPERTENSION FROM CHRONIC NEPHRITIS	234
COURSE AND PROGNOSIS	234
TREATMENT	236
Introductory	236
Mental Quiet	237
Work and Exercise	237
Sleep	237
Diet	237
Lowering Blood Pressure	238
A Regime	239
Headache	239
Angina Pectoris	239
Cardiac Failure	239

CHAPTER XXV

ARTERIOSCLEROSIS SIMILAR HEART MYOCARDIUM

ARTERIOSCLEROSIS	PAGE
Pathological Anatomy	240
Recognition of Arterial Disease	241
Consequences of Arterial Disease	242
HEART DISEASE IN THE AGED	244
Pathological Anatomy	244
Clinical Features	244
THE MYOCARDIUM	245
Some Physical Signs of Myocardial Involvement	245
Pulse Strength	245
Heart Sound Intensity	246
Gallop Rhythm	246
T' Wave inverted	247
Bundle Branch Block	247
Pulsus Alternans	248

CONTENTS

	XIX
	PAGE
Myocarditis Fatty and Fibroid Heart	250
Myocarditis	250
Fatty Heart	250
Fibroid Heart	251
Myocardial Weakness	251

CHAPTER XXVI

BRONCHITIS AND EMPHYSEMA

INTRODUCTORY	253
PATHOLOGICAL ANATOMY OF HEART	253
THE CAUSE OF THE CARDIAC FAILURE	254
CARDIAC SYMPTOMS AND SIGNS	254
Regression of the Heart	254
Cardiac Failure	255
PROGNOSIS AND TREATMENT DURING FAILURE	255

CHAPTER XXVII

THYROTOXIC STATE

INTRODUCTORY	257
CARDIOVASCULAR MANIFESTATIONS	257
Pathological Anatomy	257
Vasodilatation	257
Pulse	257
Precordial Signs	257
Auricular Fibrillation	257
Cardiac Failure	257
Anginal Pain	257
DIAGNOSIS	257
TREATMENT AND PROGNOSIS	257

CHAPTER XXVIII

CONGENITAL MALFORMATION

PERSISTENT DUCTUS ARTERIOSUS	258
BICUSPID AORTIC VALVE	258
COARCTATION OF THE AORTA (ADULT TYPE)	258
OPEN FORAMEN OVAL	258
DEFECTIVE INTERVENTRICULAR SEPTUM	258
UNEEXPANDED INFUNDIBULUM (PULMONARY STENOSIS)	258
DIAGNOSIS	258
PROGNOSIS AND TREATMENT	258

CHAPTER XXIX

CHILD BEARING ANAESTHETICS AND OPERATIONS

CHILD BEARING	
Risk of Child Bearing	

DISEASES OF THE HEART

Failure and Child Bearing
 Management of Pregnancy
 ANAESTHETICS AND OPERATIONS

CHAPTER XIX

DIAGNOSIS PROGNOSIS, AND TREATMENT

DIAGNOSTIC TERMS AND SUMMARIES
 PROGNOSIS

A Normal Heart

Chronic Heart Disease and Prognostic Groupings .

Conversing with the Patient and his Friends

TREATMENT

Healthy Habits

Avoidance of Infection

Tobacco

Exercise and Manual Work

Strenuous Acts

Diet

INDEX

275

277

277

278

281

283

283

284

284

284

285

285

289

CHAPTER I

CARDIAC FAILURE BREATHLESSNESS

THE CENTRAL PROBLEM

It is impossible thoughtfully to survey, in the light of daily experience the field of medical work covering diseases of the heart, varied as the manifestations may be, without realising the central problem to be failure of the heart to accomplish its work in lesser or greater degree. This work consists in the propulsion of blood through the circle of vessels in adequate quantity to meet the needs of the body in the ordinary and varied circumstances of life. The very essence of cardiovascular practice is recognition of early heart failure and discrimination between different grades of failure. This simple truth is not stated here for the first time, in theory it receives occasional homage from many. It emerges into view for a fleeting moment to retreat and lie concealed beneath a mass of technical, and by comparison trivial detail, it does not dominate cardiac practice as it should. When a patient seeks advice and heart disease is suspected or is known to be present, two questions are of chief importance. Firstly has the heart the capacity to do the work demanded of it when the body is at rest? Secondly, what is the condition of the heart at rest? These questions are answered by the background in most cases of chronic heart disease, they are essentials to sound prognosis and treatment.

CARDIAC FAILURE (GENERAL STATEMENT)

There is but one meaning to the term cardiac failure—it signifies inability of the heart to discharge its contents adequately, and this failure culminates in manifestations with which we are all familiar, namely, dilatation of the heart, general venous congestion, and

dropsy In the progress of failure these phenomena are late, and they mean that the heart has failed in such degree that it has no longer the capacity to do the work required of it while the body is at rest In much earlier phases of cardiac failure, before signs of venous congestion have appeared, the underlying defect shows itself in a lack of reserve by means of the cardinal symptom breathlessness on effort In this book I shall use the words "cardiac failure" consistently to mean failure at any stage of its development, when I desire to indicate the last stages, I shall speak of "failure with congestion"

We may trace the development of the malady as a whole, and correlate its several phases by considering the events as they succeed each other when heart failure comes insidiously and gradually, the reserves being first weakened and the defect displaying itself transiently, the reserves becoming more reduced, and ultimately lost, when the incapacity of the heart is paraded constantly *Breathlessness is not in itself abnormal All people, however healthy, become breathless on exercise The abnormality consists in the occurrence of breathlessness in unusual circumstances The first indication of cardiac failure is a limitation of the subject's activity, he notices breathlessness or unusual breathlessness on undertaking some customary act of vigorous work As failure progresses the same breathlessness continues to be experienced, but it is experienced in response to an amount of work that decreases as time passes This progression may be gradual and spread over months or years, or it may occur in definite steps Cardiac breathlessness is to be graded not by the degree of distress it occasions, but by the amount of physical effort that brings it into being Breathlessness occurs on walking quickly or uphill on walking on the flat on walking slowly short distances, later it is present at rest, these are the grades It is in the last two stages that the early signs of actual venous congestion appear and, as congestion increases, breathlessness at rest becomes more obvious and more distressing Thus the complete course from health to gross heart failure is associated with breathlessness ranging in its degree from normal breathlessness on active exertion to constant distress when the body rests completely Breathlessness is to be ascribed to a deficiency in the flow of aerated blood to the head and neck at first the deficiency is confined to those exercises in which normally the cardiac output is much above resting value, at last there is a deficiency in the physiological quantity of aerated blood expelled by the heart while the body is at rest*

The deficiency, in the later stages, consists not only of inadequate flow, but also of insufficient aeration, a matter not at the moment relevant but discussed later (pages 38 and 165). It is when the output at rest declines that blood begins to collect on the venous side and the patient begins to manifest signs of congested veins and, associated with these, enlargement of the liver, cyanosis, a high coloured and scanty urine, ascites, dropsy of the lower members, and congestion and oedema of the lungs.

Thus cardiac failure may be divided into two stages: the stage of symptoms or *waning reserve*, and the stage of signs or of *circulatory embarrassment and breakdown*. It is a matter of great importance that the full extent of the normal heart's reserve power should be appreciated. What the heart is called upon to do under resting conditions is but a small fraction of that of which it is capable. If the total capacity of the healthy heart for work is taken at ten units, it is a fairly accurate estimate to say that one unit suffices to maintain a normal circulation while the body is at rest, and that the remaining nine form the reserve. Consequently, in considering the development of failure, it is to be realised that by the time congestion of the venous system sets in, nine tenths of the heart's capacity to perform its tasks have been lost. Venous congestion is not an early but a late manifestation of failure, breathlessness on effort is the early one.

BREATHLESSNESS ON EFFORT

The first indication of cardiac failure is to be found in a diminished tolerance of exercise. Of the very numerous tests of cardiac efficiency and inefficiency that have been devised, based as they are mainly upon pulse rate or upon blood pressure or upon both, there is none that approaches in delicacy the symptom breathlessness. For this reason, and because undue breathlessness on effort is the commonest complaint of cardiac patients, it is the symptom that takes first place in our work.

The most important gauge of a reduced respiratory reserve is to be obtained, when dealing with an intelligent patient, from himself. He tells us that some act, previously undertaken without noticeable distress, has recently begun to cause breathlessness or undue breath

In gauging finally, a comparison between a healthy and an unhealthy individual is usually far less satisfactory than between the healthy and unhealthy states of the same subject. In estimating the case with which breathlessness is called forth, a careful interrogation of the patient is to be placed first, not only for the reason given, but for another that is equally important. For this interrogation, while acquainting us with the patient's bodily activities in his daily life, in work and in play, brings us at once into contact with the precise difficulties which the patient actually experiences, in brief it provides just that information which will be required when the control of bodily work for purposes of treatment comes to be considered. The questioning should be close, therefore and should cover a number of activities, so that while full information is obtained, there may be an opportunity of observing consistency or inconsistency. But all patients are not intelligent, neither are all accurate in their statements, and grave inaccuracies are especially liable to appear when the examination is conducted for a public service or for purposes of insurance. It is desirable, therefore, to be familiar with normal reactions to simple forms of test exercise. It is undesirable that tests should become too stereotyped, or that they should be regarded as exact measures: a test that is convenient in one circumstance is inconvenient in another. It is important that the tests employed should be few in number, so that the reactions to them may be the easier to know, and that the exercise performed should be simple and natural. No more than one or two tests are required in any given case. It is a rule that no patient who exhibits breathlessness when standing still after undressing, or while lying at rest, should be tested. The tests in these are indeed wholly unnecessary, since it is already clear that there is no reserve. If this lack of reserve is cardiac in origin the early or later signs of venous congestion will nearly always be found.

SIMPLE TEST

The subject walks briskly up a flight of forty steps, one step at a time, and is examined at once. The equivalent, twenty hops on the right and twenty hops on the left, may be used, the shoulders being raised about six inches at the same time, or the subject may step on and off an eighteen inch chair. But of these tests is so satisfactory as, though customarily used, stairs, since they are a part of the subject's life, and the tests but leading a

reaction, breathing will show no obvious hurry or material increase of depth, questions will be answered without interruption by respiratory movement. The pulse will rise no more than 10 to 20 beats per minute, and the original rate will be resumed within a minute or a little more.

A number of cardiac subjects will show manifestly disturbed respiration on this test. They are subjects who usually complain voluntarily of breathlessness on walking upstairs or briskly on the flat, and belong, when the breathlessness is of cardiac origin, to a class in which the reserves are largely exhausted, but not as yet to the point of circulatory breakdown. The simple test can be employed with safety in the case of any patient who has walked to the consultation, or who shows no breathlessness on undressing.

STRENUOUS TEST

To produce breathlessness is the object of this test. A weight of ten to twenty pounds is raised from the floor, the arms flexing easily and then extending, above the head at the rate of one lift in two seconds. A healthy young man of sedentary habit can accomplish this lift thirty to sixty times, but at the end he will be so breathless that he will want to stop. An athlete in training will continue much longer without complaint. Unhealthy men, with deficient respiratory reserve, will be distressed when far less work has been done, the exercise should be stopped if, during the exercise, the mouth opens in inspiration or if there is any other clear sign of respiratory embarrassment. The number of the lifts is the gauge of the reserve. This test is only to be employed upon subjects who have passed the simple test; its effects should be watched closely while it is in progress, and the exercise stopped as soon as breathlessness is obvious. In these circumstances it can be employed safely, irrespective of what has been found in the heart.

Cardiac cases may be divided, from the standpoint of heart failure, somewhat arbitrarily into four classes:

(1) *Good Exercise Tolerance*—Able to engage in all but the more strenuous forms of exercise and work, passing the strenuous test.

(2) *Fair Exercise Tolerance*—Able to walk on the flat, at a good pace, and upstairs without stopping or distress, become breathless on hurrying and on hills; fail to pass the strenuous test.

(3) *Poor Exercise Tolerance*—Complain of breathlessness on walking upstairs or briskly on the flat. Fail to pass the simple test.

(4) *No Exercise Tolerance*.—Breathless when lying at rest or on the slightest exertion. These patients display signs of venous congestion in greater or lesser degree.

The differentiation of breathlessness that is due to early cardiac failure from breathlessness on effort due to other causes is dealt with later (page 163).

CHAPTER II

CARDIAC FAILURE WITH VENOUS CONGESTION

In the last chapter cardiac failure was dealt with in outline and especially from the standpoint of its early manifestation in the symptom breathlessness. The present chapter describes the symptoms and signs of the late stages of cardiac failure.

SYMPTOMS

Breathlessness is the main symptom and varies much in its degree. In deeper congestion is hurried, the pulse passing 35 per minute.

these sufferers much pressed.

Fatigue and Exhaustion—Fatigue is an early symptom of the late stages of cardiac failure. During walking exercise the patients not only become breathless but their legs feel heavy and on reaching home they are abnormally weary. Another symptom coming on during exercise is a sense of sinking and weakness associated with tremulousness—a symptom often termed exhaustion. Many patients, while lying in bed and presenting congestion of the veins feel fatigue constantly.

Fulness of Head—A tense feeling across the forehead is a usual symptom of advanced venous congestion headache, though not the rule may replace it.

Pain—Many of these patients complain of aching over the precordium and when congestion is deep continuous aching pain of a distressing kind develops in the right hypochondrium or more serious sputum, increases distress.

(4) *No Exercise Tolerance* —Breathless when lying at rest or on the slightest exertion. These patients display signs of venous congestion in greater or lesser degree.

The differentiation of breathlessness that is due to early cardiac failure from breathlessness on effort due to other causes is dealt with later (page 163).

pressure in the right auricle, it is this gauge that is of value. If the needle is passed into the median basilic vein of a normal subject lying supine, the fluid column in the manometer will usually come to rest level with the sternum, or a centimetre or two below it, or less commonly above it. Thus, if the punctured vein lies level with the sternum, the meniscus will come to rest at or about the level of the puncture and, as the arm is depressed, it will not move appreciably relative to the sternum, but the column will reach above the vein by the amount the arm is lowered. The venous reservoir thus behaves much like a simple reservoir (Fig. 1). In this the surface of fluid lies at a given level, and there the pressure is atmospheric or zero. In the depths of the reservoir pressure is greater, and it is




FIG. 1.—A reservoir of water in a wide pipe joined to it. If manometers are attached at *a b c* the measured pressure above these points is represented by the increasing size of the columns *a b c* but all these columns rise to the same zero level (Z.L.). All indicate equally well the height of the water in the reservoir.

greater by the amount of the vertical column of fluid that extends from any given point to the surface of the fluid.

In determining venous pressure by bedside means, attention should first be directed to the level at which the veins collapse.

Venous Swelling and Collapse.—We may use the term zero (or atmospheric) pressure level of the fluid in the venous system in much the same sense as it has previously been employed in speaking of the reservoir, and that level is near the lower border of the manubrium sterni in normal subjects. This point of reference is chosen because it represents approximately the level of zero pressure whether the body is horizontal or vertical, or in any intermediate position. It is likewise true of the flaccid body that the pressure in veins lying at different levels below the sternum is greater, and at levels above the sternum is less, than atmospheric pressure. Thus, normally, all veins which lie higher than the manubrial point are

Vomiting is not uncommon after food has been taken and some times interferes with the oral administration of drugs

SIGNS

The first signs of circulatory breakdown are to be found in the systemic veins. The labouring heart fails to transfer an adequate amount of blood from the venous to the arterial side, the blood collects in the veins, and they swell as the pressure in them rises. It is not in the legs or in the abdomen that the earliest and best evidences of this change will be found, but in the veins themselves. Study of the veins still suffers a most unfortunate neglect, in them are to be found some of the most valuable signs we possess in managing heart cases. A full grasp and working knowledge of the phenomena to be discussed cannot be attained at once, but only by diligent observation and thought.

MEASURES OF VEIN PRESSURE

Direct Method and Principle—Direct measurement of venous pressure will often provide an invaluable guide to a patient's state, repeated readings will often indicate clearly the course of the malady. But the direct measure can never come into general use, moreover, it can be replaced by simple bedside tests that any thoughtful observer may apply. The direct method will be described briefly and mainly with the object of illustrating the hydrostatic principles involved. The apparatus in simplest form consists of a U-tube manometer connected to a wide needle by rubber tubing, the whole system being filled with a solution of sodium citrate to prevent clotting. The needle is driven into a vein, and the height at which the fluid comes to rest in the manometer is observed. A vertical measure, taken from the meniscus to the point at which the vein is punctured, accurately measures the pressure in the vein lying at its then level. If the vein is in the arm and the arm is lifted up, the pressure in that vein will fall, or if the arm is lowered it will rise, correspondingly, thus measuring the actual pressure in the vein has little value, since it is constantly varying in different cir

the same height relative to the heart, whether the arm is raised or lowered. Thus, the height of the meniscus serves as a gauge of

you would think of the level of fluid in a long pan of water, lifted at one end or at the other, watch the rise and fall so that you may recognise the surface and know the general level (Fig 3)

The venous zero level rises to various heights in congestion. Where venous pressure is raised a little the column is seen to stand in the neck of the supine subject as high as the centre of the sterno mastoid muscle or beyond this, it comes unmistakably above the sternal level on both sides of the neck. Where pressure is greater, the veins are swollen to the jaw. Note especially that it is not the anatomical point reached in the neck that matters, but the vertical




FIG 3—A diagram illustrating excessive venous pressure equivalent to about 8 cms of water. In the upright position the zero level may be just a little above the clavicle as the subject is inclined the zero level comes to occupy a higher and higher position in the neck until it reaches to the angle of the jaw. Its vertical distance above the manubrium does not change much.

distance above the sternum. The former is affected by the inclination of the neck to the body.

Where pressure is higher still, the veins remain swollen throughout the neck when the head and shoulders are lifted off the bed. The gauge remains unaltered, it is the vertical height to which the swollen veins extend, or may be raised, above the sternum without collapsing. (The veins are never full in the neck in a normal unclothed person standing easily in ordinary circumstances in the erect posture, in conspicuous congestion they run the length of the neck like cords.)

Veins swollen on one side only point to local obstruction of their immediate outlet and have no value from our present standpoint, often such swelling can be released by a little rotation of the neck,

the vein that has the lowest pressure is alone to be used as an index of general venous pressure

The bedside method here described of estimating the level of venous collapse is not wholly adequate because in many patients superficial veins of sufficient size to stand out from the neck are not found. In such cases, and they are frequent, the required information is often to be obtained by closely observing the venous pulse and its behaviour.

Venous Pulse as Index—Hitherto we have treated the venous reservoir as though the zero level were unvarying. Actually this is rarely so, almost always the level fluctuates a little with the beat of the heart and this fluctuation is often very helpful in gauging the level to which the veins are filled. The venous pulse as we see it in the neck is a normal phenomenon, being caused by alternate expansion and collapse of the vein in which it is occurring, and this expansion and collapse may be regarded, properly, as resulting from the rise and fall of the blood tide in the veins. Though it occurs at lower levels also, the venous pulse has its greatest amplitude immediately at the general level of venous collapse. Veins that are tensely distended do not pulsate appreciably, neither can those that are empty. It follows that pulsating veins are full veins, and that the uppermost point of free pulsation is an index of the level to which they are filled, this index is used, as previously remarked, when the actual point to which the veins are filled cannot be observed directly. Thus it is clearly most important to be able to recognise venous pulsation when it is seen, but this requires very careful study if it is to be accomplished with certainty.

In normal health, the subject being supine, venous pulsation occurs at the root of the neck, often, and especially in young subjects, it is a movement of large amplitude. It is obviously a venous phenomenon when seen in such a superficial vein as the external jugular. In superficial veins it can be studied in relation to the level of venous collapse, at the point of highest filling the vein pulsates freely, the movement being complex and undulatory (as in Fig. 4). The level of pulsation in the vein is seen to alter with respiration, approaching nearer to the heart at each inspiration, for this lowers venous pressure. But there is also a diffusely distributed venous pulsation in the lower neck, occurring along the line of the carotid sheath and subclavian vessels. This is mainly produced in the internal jugular, in the subclavian vein, and in the large superficial tributaries of the latter in the subclavian triangle (lower half of

posterior triangle), and its recognition in these deeper veins is essential and must be cultivated. Most of the pulsation in the neck of such deeper origin and, while it cannot then be seen to arise from veins, can and must be recognised as venous by its characters, its undulatory form, its usual softness, its frequent large volume, and its change of site with posture. Venous pulsation disappears altogether when the erect posture is assumed, because the cervical veins then collapse, it moves higher up the neck when the head is lowered or the abdomen is pressed upon. It extends relatively high in the necks of young people. Such in general is venous pulsation in the uncongested.

In the congested the signs are modified. The field of pulsation is displaced to a higher level in the neck. The largest pulsation comes from the deeply lying internal jugular and, with the patient's head or shoulders supported on pillows, frequently extends to the jaw and moves the lobe of the ear. It is often slow, welling, and sustained, rather than undulatory. When much congestion is present and the subject's head is well raised on pillows, the veins are often too tense to pulsate, tilt the patient into a more erect posture, and the level of zero venous pressure falls in the neck and, as this tide recedes, abundant pulsation appears below the jaw. In these patients venous pulsation may be seen in the neck when they stand, or spreading to the veins of the upper arm or forearm while they lie. Whenever veins pulsate freely above the level of the manubrial line they are overfull, the higher the level of pulsation the greater the overfilling. It is not venous pulsation




FIG. 4.—Venous (or jugular) radial and electrocardiographic curves from a normal subject. The vertical lines mark $\frac{1}{2}$ sec in this and similar figures and any vertical line cuts the three tracings at the same instant in time. Horizontal lines in this and all similar figures mark $\frac{1}{2}$ millivolt. The jugular curve is composed of *a*, *c* and *v* waves. The wave *c* occurs about $\frac{1}{2}$ sec before the radial upstroke. It is actually simultaneous with carotid pulsation and marks the beginning of ventricular systole in the venous curve. The *a* wave lies in presystole and results from the auricular contraction. The summit *v* marks the end of ventricular systole and occurs opposite the bottom of the diastolic notch in the arterial curve. Systole of the auricle is represented by *P* and of the ventricle by *Q S* and *T* in the electrocardiogram. The auricular and ventricular events are later in the mechanical records than in the

that is abnormal it is the high level at which it occurs that constitutes abnormality. The examination must be made thoughtfully thus in many normal subjects if they lean forward as they sit prominent pulsation appears in the jugular veins that is so because pressure on the abdomen in this posture displaces blood into the neck.

Venous is very often mistaken for arterial pulsation especially by those who think the distinction always easy it is often a most serious blunder to make. It is not generally recognised that venous pulsation is much the commoner has usually the greater amplitude and more often moves the lobe of the ear. Pulsation that is maximal in the subclavian triangle is almost always venous pulsation maximal in the carotid triangles often is. An arterial pulsation is usually maximal in the carotid triangle and it increases when the erect posture is assumed a corresponding venous pulse lessens or disappears under the same test. An arterial pulsation that is variable is always palpable it jerks sharply against the finger to a sustained thrust. Venous pulsation is very rarely strongly palpable and only occasionally can be felt the testing finger should be laid as lightly as possible upon it. When as during congestion the venous pulsation tends to lose its undulatory character and becomes more sustained and sometimes plainly palpable the distinction between it and arterial pulsation is not always easy but in such cases the remaining signs of a congested venous system are always manifest.

OTHER SIGNS OF CONGESTION

The Liver—The earliest sign of a congested liver is usually obtained by percussion. I prefer to use it in the mid line and of moderate intensity normally the abdomen is then resonant to the midriff the thin margin of the liver crossing the intercostal angle fails materially to modify the note unless the liver edge has low or light percussion is used. In congestion enlargement is uniform and the firm smooth margin is usually palpable at a varying level between rib margin and navel or lower failure to palpate it is often due to beginning to feel at too high a level. When there is much enlargement the abdomen is usually convex between ensiform and umbilicus and the liver pulsates softly like the veins pulsation is often present though not always easy to detect it has more interest than importance. Tenderness of the abdominal wall over the enlarged liver is frequent and may be associated with rigidity and pain. In very deep congestion the skin becomes jaundiced except where it is

involved in oedema, in such regions the skin shows no stain, owing to dilution of the bile pigment, which passes the capillary wall less easily than the fluid

The Kidneys—The normal quantity of urine passed daily is 30 to 50 oz (1000 to 1500 c cs) it varies much with the activity of the skin and the amount of fluid drunk. Congestion of the kidney gives a scanty and concentrated urine which deposits phosphates and urates and may contain a trace of albumen, a few hyaline casts and red blood corpuscles

The Lungs—Congestion and oedema of the lungs are early displayed by moist crepitations at the bases on taking a full breath, and later by weakness of breath sounds and voiced sounds and by dulling of the percussion note. These signs are associated with cough, and a sticky and sometimes blood stained sputum. A fuller description of passive congestion and oedema than is here required, and of the manner in which it produces breathlessness is given in Chapter XVIII

Cyanosis and Oedema, though frequent accompaniments of venous congestion, are met with in many other conditions. While the former may be a guiding sign leading to the diagnosis, neither can be safely used as an evidence, of venous engorgement. It will be convenient to consider each of these phenomena at length a little later (pages 18 and 36)

Malnutrition is a common accompaniment of long standing venous congestion. Often when the weight has risen owing to accumulation of fluid in the tissue spaces the face, arms, and chest show obvious wasting

Delirium is not infrequent in the terminal stages, though the patients more often sink gradually into a state of deepening unconsciousness and die comatose

DIAGNOSIS OF CONGESTION CORRELATION OF SIGNS

When the full signs of congestive failure are present, neither close observation nor much discrimination is required to know it. The diagnosis of early congestion is more difficult and in the long run much more important. The diagnosis of congestion, whether early or late, should be made finally on the evidence provided by the breathing, by the veins and liver almost exclusively. It is for this reason that these have been emphasised. Experience teaches that to place reliance upon a single sign is precarious. Compare this sign and that, and confident recognition of the patient's state grows as

these signs group themselves together to form a harmonious picture. Thus on finding unquestionable evidence of engorged cervical veins we look at once at the liver and, finding it enlarged, are satisfied that the engorgement is general and not confined to superior caval territory.

Generally speaking, enlargement of the liver in its several degrees goes hand in hand with engorgement of the veins. The liver does not usually enlarge before the rise of pressure in the veins can be detected; neither do the veins swell without the liver edge descending perceptibly. Thus the early diagnosis of congestion turns upon a close examination and correlation of these signs. And here another correlation is most important. Patients who suffer from general venous congestion are without exception breathless, either while lying flat at rest or upon very slight exertion. When early venous congestion is strongly suspected but not quite certainly diagnosed, then the presence of such breathlessness goes far to complete the diagnosis. If, on the other hand, a subject is thought on examination to manifest early signs of general venous stasis, and yet it is clear that breathlessness is not experienced even with quiet or moderate exercise, the opinion first formed must be revised for it is incorrect. Moreover, there is a clear relation, constant within narrow limits, between the degree of breathlessness and the pressure in the veins; it is so clear that if it is established that a patient has no congestion of the venous system and is yet definitely breathless in bed, then that breathlessness is not primarily cardiac in origin. The proper use of this knowledge saves a number of serious diagnostic blunders. It is to be repeated that a full understanding of cardiac failure is obtainable only by studying the manner in which its symptoms and its signs are linked together; the subjective and objective manifestations are part and parcel of one process and dependent upon the same fundamental cause, namely, lowered capacity of the heart for work.

The differential diagnosis of enlargement of the liver from congestion and from other causes rests mainly on an examination of the veins. The differential diagnosis of acute venous congestion is dealt with later (page 75).

Discrepancies — If it is beyond question that the veins of the neck are bilaterally congested and it is equally beyond doubt that the liver is not enlarged, an obstruction of the superior cava must be considered. Its diagnosis will rest upon the discovery of anastomotic veins and upon failure to induce the veins of the neck to pulsate. A

second reason for the same discrepancy is atrophic cirrhosis of the liver in a congested patient, the diagnosis of the liver condition will then turn upon the degree of hardness, and perhaps irregularity, of the liver margin

There is the reverse case an engorgement of the liver has been present for a long time and the venous spaces within have become permanently dilated and its substance a little or much fibrosed In such, even if the signs of increased pressure in the veins greatly decline, the size of the liver may not decrease much or proportionately It is a discrepancy which previous knowledge of the course of the malady explains

CHAPTER III

CARDIAC OEDEMA

FACTORS IN CAUSATION OF DROPSY

THE cause of cardiac dropsy is increased transudation of fluid from the minute blood vessels into the surrounding tissue spaces, and in producing this transudation the most important factor is increased pressure within the small vessels. If the limb of a normal subject is maintained horizontal and a pressure of 30 mms Hg (41 cms water) is thrown upon the veins, recognisable dropsy will appear in the limb within the space of a few hours. If a normal subject stands and allows a leg to hang down, and quite flaccid and still, a considerable hydrostatic pressure gathers in the small vessels of the foot, which becomes recognisably oedematous within a few hours, the same thing happens, though in longer time, to the foot of a subject sitting in a chair provided that the limb remains still. The reason why the feet of normal people do not become oedematous while standing or sitting ordinarily is that the muscles of the limb are never long quiescent, but exert a pumping action on the veins and lymphatics this pumping action greatly lowers the mean pressure in the veins, and tends to empty the tissue spaces of fluid. It is important to grasp that a certain grade of oedema of the feet may be on occasion almost physiological, and that the normal balance between the flow of fluid into and out of the tissue spaces is very easily upset. In cardiac failure increasing dropsy is the rule when general venous pressure rises and is maintained at 10 to 15 cms pressure or more (of water) above normality.

Another factor proved to influence the rate at which dropsy forms in man is temperature, the higher the temperature of the part the faster the exudate occurs, a clinical instance is that in which dropsy of the legs forms quickly in patients while they sit before a fire.

RECOGNITION

In cardiac dropsy much of the fluid collects in the subcutaneous tissues, where it is easy to identify. It causes swelling, and the overlying skin is stretched and becomes as a consequence pale and shiny. If the stretching is very great "lineae atrophicae" develop. Oedema is frequently accompanied by tenderness, which may outlast the remaining signs when oedema disappears. In recognising early dropsy of the subcutaneous tissues it is usual to press upon the part with the finger, the excessive tissue fluids are thus displaced and the skin indented as is moist clay when similarly treated. Firm pressure continued for 5 seconds will often disclose a slight oedema that would otherwise escape notice. Even so elicited, pitting is not a very early sign of the increase in tissue fluids. It is known that the volume of a limb must increase, through the accumulation of tissue fluids by a tenth, before pitting becomes clearly recognisable. Patients who are becoming dropsical begin to put on weight some days before pitting can be found, patients who are losing dropsy continue to lose weight for some days after the last sign of pitting has gone. Advanced dropsy is at once recognised by the relative hardness of the tissues and inability to pick up the skin. The extent of such dropsy can be distinguished easily by simple palpation.

DISTRIBUTION

Cardiac oedema is one form of dropsy that is essentially hydrostatic in its distribution. In patients who spend much of their time upon their feet or sitting, the feet are the first parts of the body to swell and they become most swollen. It is in the instep and ankle that oedema is usually first seen in such patients though the distribution may be modified by the nature of the footgear. Patients with early dropsy who have been up and about often find the oedema forming as a rim over the upper parts of the shoe or boot, swelling of the foot follows removal of outdoor shoes which are then replaced with difficulty. In patients who lie in bed pitting is often found first just behind the malleoli. A continuous and gradual involvement of the limb from foot to thigh, genitalia and trunk occurs in those who spend their time sitting. In bed patients, after the ankles swell the genitalia or sacral region may next show change. These differences depend upon the levels at which the several affected parts lie. If the buttocks sink low in the bed, sacral oedema

will come earlier and be the more in evidence. Parts upon which the patient actually rests show little oedema, the fluid being readily displaced by pressure, when patients are propped on several pillows, and oedema has risen to the trunk, it is common to see a ridge of oedema across the back, between the areas pressed upon by two separate pillows. Oedema spreading on the trunk above the navel is relatively uncommon in purely cardiac cases, occurring only in very advanced stages of failure. Similarly, it is unusual in the hands, unless these are kept for long periods hanging down, or unless, as sometimes happens, thrombosis occurs in the main veins of one limb, oedema being then unilateral. Oedema of the face or scalp is not cardiac, but usually renal.

To retain a clear idea of the influences that affect the distribution of cardiac oedema is important from several points of view. Oedema cannot be decreased by changing the patient's posture, but it can be redistributed. If a patient comes with slight dropsy of the feet and is placed flat in bed, this oedema will often disappear within a day or two because the excessive fluid in the tissue spaces of the legs moves to tissue spaces higher up in the body. Hanging the legs down often enables these limbs to act temporarily as a reservoir and to accumulate fluid, which otherwise might come to lie within the pleural sacs.

DIFFERENTIAL DIAGNOSIS

Dropsy of cardiac origin is recognised by its hydrostatic distribution and by its association with increased general venous pressure. If by careful use of the methods described in the preceding chapter general venous pressure is judged to be normal, then dropsy in the legs can be stated to be due to no weakness of the heart. This very valuable rule is broken only by those instances of cardiac dropsy in which, owing to an unusually rapid fall of venous pressure, as from venesection, the disappearance of oedema lags notably behind. Dropsy of hydrostatic type is a very common affair, and much more often than not it is produced in other ways than through cardiac failure. It may result, rarely, from various forms of local obstruction of the inferior vena cava or its tributaries, as by thrombosis, by pressure from tumour or collections of fluid in the abdomen. It is frequent where the veins are varicose. Most important from the diagnostic standpoint is the fact that hydrostatic dropsy is a very common event in middle aged and elderly people who are heavy, who stand or sit much, but who otherwise are healthy. These

patients are often and erroneously regarded as gravely diseased, and they and their friends are consequently caused much needless anxiety. Hydrostatic dropsy is often present as a late event in diseases in which there is great wasting. The distinction between these forms and cardiac dropsy rests usually upon a thorough examination of the venous tributaries of the superior cava, to exclude a general rise of venous pressure.

EXUDATES INTO SEROUS CAVITIES

These are for the most part late manifestations of cardiac dropsy. Ascites, however, usually comes earlier than material hydrothorax, and occasionally it appears prominently and is long continued before dropsy of the legs is seen. Ascites without dropsy of the legs is chiefly confined to cases of rheumatic heart disease in young people, and is then generally associated with mitral stenosis or mediastino-pericarditis, accompanied by long-standing congestion of the liver and perihepatitis.

CHAPTER IV

FAILURE WITH CONGESTION (CAUSE, PROGNOSIS AND TREATMENT)

CAUSE OF CARDIAC FAILURE

A BRIEF survey of the cause of cardiac failure at this stage will enable an early and broad view to be obtained, and facilitate an understanding of much that is said in more detail upon the same subject in later chapters

In considering the way in which failure of the heart is brought about, the mechanical factors have in the past received undue emphasis. When the heart fails adequately to discharge its contents, it does so because its beat is insufficiently powerful to overcome the resistance it meets therein is expressed the cause of cardiac failure. It will be perceived that failure may happen for one of two distinct reasons, namely, weakness of beat or increased resistance to discharge, or it may happen through a combination of these two.

It is clear that a complete obstruction of any of its orifices can by itself determine failure of the heart, but anything approaching to complete obstruction is rare clinically. An increased resistance to the left ventricle occurs when blood pressure is raised, as in hypertension, or during strenuous bodily efforts, or in aortic stenosis. An increased resistance to the right ventricle occurs in pulmonary and mitral stenosis and in pulmonary emphysema. The output of the ventricles is impeded by certain kinds of pericardial adhesions. The extent to which these increased burdens deplete the reserves of the heart will not be considered here, it is dealt with more fully in speaking of valve disease (page 143) and in other places. It will be enough to say that these reserves are now realised to be very great and difficult to exhaust. There is insufficient reason to believe that any one of the abnormal resistances above named, when acting alone, can determine failure, unless indeed the resistance is altogether exceptional in quantity.

As knowledge of heart failure has developed stress has come to be placed more on the state of the cardiac muscle, for it is quite clear that the muscle may be so damaged that the heart will fail, even at a time when the burden it has to bear is no more or even less than customary. A simple instance is that of failure with congestion arising while the subject is at rest, from thrombosis of a coronary artery killing a substantial portion of the ventricular wall. Another instance is pernicious anaemia, through which nutrition of the heart muscle as a whole may be so reduced that failure supervenes. The acute infectious diseases provide examples of acute heart failure resulting purely from poisoning of the muscle.

In asphyxia from any cause, the heart swells and the veins become more and more engorged until no blood is ejected, an acute failure due directly or indirectly to lack of oxygen, but one in which increased resistance to output may for a time play a minor part. In subacute bacterial endocarditis (page 187) and in hyperthyroidism (page 259) heart failure is chiefly due to change in the muscle, though contributed to by valvular disease or by increased output. Another undoubted and frequent example of the same kind is failure with congestion arising during the course of rheumatic carditis (page 197), the fact that such congestion comes and later disappears completely, shows that it is not due in the cases that display this sequence of events to permanent anatomical changes, such as mitral stenosis or pericardial adhesions for these remain after the congestion has cleared away.

It will be found by actual enquiry or observation that patients who develop failure with congestion, while under supervision for heart disease do so more often in immediate sequence to an infection than in response to a period of unusual or unwise bodily exertion. The infection is not necessarily rheumatic or pneumococcal, but may consist of a febrile cold, influenza, or bronchitis.

As we pass instances of failure with congestion in review, usually substantial evidence is in each case forthcoming that a myocardial weakness is a chief or heavily contributing factor. One of the most frequent determining factors of congestion is the onset of very rapid

It does so in two ways. First it causes the ventricles to expend more energy. At each of its beats the ventricle must raise the pressure within its cavity to arterial pressure before it begins to do effective

work Other things being equal, the waste of energy that occurs in this way increases proportionately to the rate of beating Secondly, rapid beating shortens the time between beats for recovery, and this not only absolutely but relatively Thus, if the ventricle is beating at 60 per minute, systole has a length of about $\frac{1}{3}$ and diastole $\frac{2}{3}$ second, while, if the rate of beating is 120 per minute, systole and diastole each occupy about $\frac{1}{2}$ second Thus the relative length of diastole is reduced to one half its former value, the ventricle enjoys but three quarters its previous total resting periods Weakness of beat follows, and here again is the chief reason for deficient output *when this occurs*

It is essential to the treatment of certain types of cardiac failure, and especially to prognosis in cardiac failure, that ideas as to causation should assume a proper balance, that the condition of the ventricle should be taken into account as well as the work it has to do, and that, because an increased burden is known to exist, this should not come automatically to be regarded as the whole, or necessarily the chief, trouble That weakened muscle is often the sole or chief, and probably almost always a considerable, factor in causing congestive failure, is not merely a view of academic interest, but one that greatly influences the general attitude to heart cases and their management especially it emphasises the need of guarding these patients against infections of any kind

PROGNOSIS

Congestion of the venous system is one of the very gravest manifestations of cardiac disease, but it has been said with some truth, though not with complete accuracy, that no patient dies in a first attack The gravity of the immediate outlook naturally varies with the degree of congestion, and the presence of general anasarca, fluid in the body cavities or congestion and oedema of the bases of the lungs, constitutes at all times a condition of great seriousness But a prognosis cannot often be made upon so simple a basis The manner in which failure has developed requires specially close consideration Sometimes a state of congestion has followed relatively suddenly upon an attack of acute bronchitis or infectious cold, and in such it will usually subside, leaving the patient much as he was Following upon pneumonia or occurring during chronic rheumatic infection the outlook is less propitious Developing during the course of subacute infective endocarditis it is a terminal event, for here

the infection that is its cause is incurable. Failure may be caused by anaemia or by hyperthyroidism which when ultimately cured may leave no cardiac weakness, it may follow an attack of coronary thrombosis and recovery of the former reserves cannot then be expected, though much lost ground may be regained. In all these instances we are weighing the chances that a primary weakness of the muscle may disappear with its cause.

Congestion may develop as a consequence of rapid ventricular action, at the time when fibrillation or flutter comes, or when the action of the ventricle in response to a fibrillating auricle becomes rapid through neglect of proper treatment. Cases of these kinds are frequent and favourable. The depth of congestion cannot be used to gauge ventricular strength unless the precise conditions of beating are known. Of two cases presenting the same signs of failure, the heart is sounder in the case in which it beats under less favourable conditions, it is the probability of obtaining relief from disadvantageous conditions of work or from an increased burden of work, that makes the prognosis more favourable. Thus, given a certain grade of failure, the presence of fibrillation with rapid ventricular action improves the outlook. Similarly, a patient who develops congestion while engaged in heavy manual work has a better prospect than one who begins to suffer similarly in a sedentary occupation. The development of congestion, having no transient or removable cause, by a subject already in bed, is ominous. It will be evident from what has been said that the immediate prognosis in congestive failure requires careful thought, and must take into account the probable response to treatment.

The place taken by cardiac failure, whether early or late, in a general prognosis that attempts to estimate the duration of life, will be found in other parts of this book, and especially in its last chapter.

TREATMENT

The treatment of failure with congestion is described in the first place without reference to the underlying cause of muscular weakness or to the burden the muscle has to bear. Most cases of failure with congestion are chronic, and the principles and measures of treatment are in general the same for all. It is convenient to deal first and in detail with cases of relatively severe congestion presenting normal rhythm, and then briefly with less severe cases, usually presenting auricular fibrillation. This will form a sufficient basis, and

such modifications and additions as are necessary in special cases will be discussed in other parts of this book

Indications for the management of patients suffering from the grades of failure that precede congestion will be found under After Care (page 32) and later and especially in the last chapter of this book

SEVERE CONGESTION

Rest—Failure with congestion is one of the few manifestations of heart disease that calls for treatment in bed and it calls for this imperatively. The watchword throughout the treatment of congestion is indeed rest—rest for the body, rest for the mind, rest for the heart. Heart failure that has reached the stage of showing deep congestion requires most skilful handling of the patient by day and night nurses. The patient must be guarded against effort of all kinds. He should lie still in bed, should not lift himself or change his own position, should not wash, dress, or even feed himself. He may be moved a little when uneasy. He must be lifted on and off the bed pan, the nurse therefore must be strong and will sometimes require help. The patient should be discouraged from talking beyond indicating his immediate wants and from attempting to assist the nurse who moves him. These precautions against effort may be relaxed appropriately in cases of more favourable type.

The motions should be kept loose by means of salines, so that straining may be avoided, two actions daily of the bowels, or one if full, is adequate, purging such patients even if they are oedematous is not good treatment.

Posture—Congested patients are uncomfortable unless the head is well raised. When the head is lowered the subject may be seen to become increasingly cyanotic in the face and breathlessness appears, or in the orthopnoeic case increases to the point of severe distress. Attacks of breathlessness at night are often due to slipping down the bed and off the pillows; this should not be permitted to occur.

The author's bedstead (Fig. 5) facilitates nursing in cardiac failure. The wire mattress of this bedstead is in three pieces, hinged together and controlled so that they can be inclined at any desired angles. The patient can be propped up to sitting positions by raising the back rest (A). Lifting the thigh rest (B) prevents his slipping and this device is more satisfactory than the simpler alternative of tying a bolster across the bed beneath the patient's thighs. The two

adjustments described, combined with a lowered leg rest (C), convert the bed into a chair

Sleep—Congested, and especially very breathless, patients who fail to obtain a number of hours' sleep in each twenty four hours rapidly lose ground through fatigue and weakness. The hours of sleep should be reported as a daily routine, and, in general, hypnotics are used when the hours of sleep fall below six. Hypnotics of the barbitone series are of the greatest general value, there is a choice between many. Luminal (phenyl ethyl barbituric acid) or luminal-sodium is given in doses of $\frac{1}{2}$ to 1 grain (exceptionally 2 grains),




FIG. 5.—The cardiac bedstead (See *British Medical Journal* 1928 ii p 977)

medinal (barbitone sodium) in doses of 5 to 10 grains, and dial (diallyl barbituric acid) in doses up to 5 grains. A single powdered tablet is placed on the tongue and swallowed with a little fluid an hour before sleep is desired. Potassium bromide and chloral (of each 20 grains) is a well tried and excellent hypnotic mixture, it occasionally induces nausea or vomiting, and should not be repeated in this circumstance. Paraldehyde (1 drachm in $\frac{1}{2}$ oz. of brandy, or in capsules) is effective, but its taste and smell are very objectionable. Where sleeplessness is associated with pain, morphia should be used ($\frac{1}{8}$ to $\frac{1}{4}$ grain), where it is associated with irritating but ineffective coughing, heroin hydrochloride ($\frac{1}{16}$ to $\frac{1}{8}$ grain) or codeine phosphate ($\frac{1}{4}$ grain) may be given by mouth.

Diet—Patients with grave congestion have little desire for food and often become distressed while masticating. Distension of the

stomach by food is clearly undesirable. The food therefore should be small in quantity, of high nutritive value and easily taken (see Diet I page 236). The fluid intake should be restricted. A diet of about 800 calories can be maintained for a few days or for a week without appreciable loss of strength; it may be increased gradually as congestion subsides or as other circumstances dictate.

Venesection—Venesection is a most potent remedy and should be used in most cases in which there is material venous engorgement (8 or more cms pressure). Only obvious anaemia contra indicates its use.

To ensure relaxed veins and ample flow the arm including the hand should be thoroughly warm. The pneumatic cuff of a sphygmomanometer is placed as high as possible on the upper arm and distended at a pressure of 60 mms Hg; a rubber band wrapped around the arm at a tension adequate to distend the veins strongly but insufficient to decrease the strength of the pulse is of equal service. The obstruction to the veins should be maintained during the period of bleeding. This may be accomplished by dissecting out a vein and opening it under local anaesthesia but is better done by thrusting a needle of large bore through the skin into the vein so as to meet the current of blood with its point towards the hand. In the latter case flow can be expedited from time to time by applying suction with a syringe. In an adult of medium weight (140 lb) 10 to 20 oz (250–550 ccs) of blood can be taken according to the severity of congestion. Rapid bleeding lowers venous pressure promptly and profoundly by the simple act of withdrawing fluid from the venous reservoirs. A decrease in the fulness of the cervical veins is almost always and in the size of the liver is often to be ascertained. Very soon however fluid is taken up by the blood from the tissue spaces and the blood becomes diluted and therefore less viscous. This reduction of viscosity is probably a chief beneficial after effect of bleeding because it increases the flow of blood through the tissues including the heart and the brain. Patients will almost always speak of relief both from breathlessness and from the sense of tension in the head; sleep becomes easier to obtain.

In cases that react most favourably venous pressure remains low after bleeding; oftener it is found to be raised again though not to its original level on the succeeding days. If the rise in level continues another bleeding of 10 oz a week or ten days after the first may be required and in the most obstinate cases three or four venesections may be needed within four to six weeks.

In this stage of failure the patient should be visited at least once a day. At these visits and during subsequent progress the most important enquiries will concern breathlessness and the hours of sleep. The signs to watch are not in the heart, but in the veins and liver, in the legs, and at the bases of the lungs.

Oxygen—It is useless to give oxygen through a funnel held near the patient's nose; the oxygen content of the inspired air is insufficiently raised thereby. Delivered through a nasal catheter, passed well back, a much higher content is obtained, and this method sometimes affords a little relief from breathlessness. The gas should be first led as a continuous stream of bubbles through warm water, a stream of dry gas is not tolerated. In recent years a few cases of congestive failure have been treated in enclosed chambers in which 40 per cent oxygen is maintained, some relief seems to be afforded, but the precise value of the method is still unknown.

Cardiac Tonics and Stimulants—The most emphatic action of digitalis and its allies is in the case of auricular fibrillation, to be considered later (page 86). The use of drugs calculated to strengthen the heart beat is open to the theoretical objection that this would increase the work of the heart, but we know so little of such drug actions upon the failing muscle that the matter must be treated empirically. There is no evidence that digitalis, given in small doses (up to 20 minims of tincture daily) recognisably influences the progress of uncomplicated congestion, either at a late or at an early stage. Its administration to heart cases, merely because they are heart cases, is to be discouraged. Occasionally, however, cases of chronic congestion that present regular heart action, and that have been treated by all the usual methods without success, seem to respond to full doses of digitalis (single doses of 2 drachms of tincture or repeated doses amounting to 1 drachm daily (see page 88)). The remedy usually comes to be tried when others have failed.

I have seen no patient respond to injections of strychnine or camphor. Alcohol is sometimes useful, not as a stimulant, but in overcoming restlessness.

Pain over an engorged liver or over an enlarged heart is to be treated first locally by the ice bag, or by irritants (a mild capsicum plaster or antiphlogistine). The application of a dozen leeches to the hypochondrium will sometimes completely relieve when other remedies have failed, morphia is the best final remedy for precordial pain.

Cough—Where cough is the result of an associated bronchitis, the expectoration of stiff mucus is helped by alcohol and stimulants such as ammonium chloride (see page 255), where it is the result of commencing congestion and oedema sedatives (as for example codeine and heroin) are more suitable. The object is in both cases to decrease the effort of coughing.

Oedema, Drainage—In cardiac failure pitting confined to the feet or lower parts of the legs usually disappears within a few days of the beginning of bed treatment, this may result simply from redistribution of the excess fluid, it may result from rapid improvement of the circulation. In cases of early dropsy, no special remedies are called for, though it is wise to restrict the fluid intake, and saline diuretics (15 to 20 grains of both potassium citrate and acetate, three or four times daily) may be given. In only a proportion of cases does dropsy call for further treatment, for dropsy is an end result of cardiac failure, and as failure is relieved dropsy disappears. Treatment should be directed, therefore, first of all to the circulation, but special remedies for dropsy should be started promptly in cases in which the accumulation of fluid is spreading to the trunk threatening the serous cavities or in which congestion is responding little or slowly to treatment. In such cases, while the fluid intake is reduced as closely as possible to 20 oz (570 ccs) a day, the measure of urine passed daily should be watched. The most potent diuretics of the xanthine series are theobromine (dimethyl xanthine), which is given in 10 grain capsules, and theocin (its isomer) in 5 to 10 grain doses, three or four times a day, the latter is the more generally used but sometimes gives gastrointestinal symptoms, headache and visual disturbance. In both cases a reaction, if obtainable is to be expected within the first few days, if it does not occur the drug should be discontinued, to be tried again later. Caffeine (trimethyl xanthine) and diuretin (the sodium salicylate compound of theobromine) are less useful. In recent years mercurials have been introduced as diuretics, and some of these are very powerful. It is said to be best to give ammonium chloride (100 grains in 24 hours) first to induce acidosis, I have not found it so. *Novasarin* is so often followed by adverse symptoms that it has given place to the equally potent but less disturbing substance salyrgan, the latter is supplied in ampoules of 1 or 2 c.c. of 10 per cent solution, the content of the smaller ampoule is injected intravenously, great care being taken that it does not enter the subcutaneous tissues, and the result observed, the dose may be

repeated as often as 10 or 20 times at intervals of three or more days. This diuretic usually gives an extremely copious flow of urine, and the flow once started, frequently continues until all dropsy has gone. It is a most valuable drug.

When dropsy of the legs has spread up the trunk, the patient should be kept in a sitting position, so that the fluid may collect in

converted into a chair (Fig. 5). If in these circumstances the legs become very large, drainage must be considered. The number of cases treated by drainage should be small, the chief indications for

I have never seen serious sepsis follow an inch long incision into the subcutaneous tissue of the dorsum of each foot, or the insertion of Southey's tubes. With abundant sterile dressing both methods are safe, but as the fluid escape is freer with incisions, this method is to be preferred. Drainage may be continued for days or a week or more, it is brought to an end because the dropsy has subsided, because the flow ceases and dropsy though decreased is stationary, and not infrequently because it begins to cause weakness of the patient.

Tapping the chest in cardiac failure is more often required for unilateral, inflammatory effusions than for bilateral accumulations, which can generally be displaced by posture with or without drainage of the feet. When aspirating, the needle should be inserted in the line of the scapula about the 8th interspace.

Tapping the abdomen in cardiac failure is called for in those occasional cases of rheumatic heart disease in young people, in which fluid collects in the peritoneum rather than in the legs. It may be repeated as it is required.

CASES PRESENTING AURICULAR FIBRILLATION

The general principles underlying the treatment of failure with congestion, and many of the details of treatment, are the same whether the auricles beat regularly or are fibrillating. Most of the cases actually belong to the latter class, and in certain respects the presence of fibrillating auricles usually calls for modified treatment. The work of recent times has shown that the chief service of digitalis is to control ventricular rate in cases of this type. By

slowing the ventricle it decreases the expenditure of energy and by slowing it increases the length of diastole relative to that of systole. This action of digitalis in conserving the heart's energy and prolonging its rest is in harmony with the chief principle of treatment of cardiac failure and justifies digitalis being called a cardiac hypnotic. The action of digitalis in auricular fibrillation is so potent that in cases of failure with rapid ventricular action rest of the body in bed combined with slowing of the ventricle by digitalis usually suffices to ensure rapid disappearance of the signs of failure. The response to digitalis in these cases can be looked for with so much confidence that venesection unless congestion is conspicuous initially can usually be avoided. Special measures for the relief of oedema, the use of the stronger diuretics or actual drainage should await the full response to digitalis and they then usually prove to be unnecessary.

Since digitalis benefits congestion in this class of case indirectly and the treatment of auricular fibrillation by this drug is the same whether congestion is present or not, the details of this treatment will be given at a later stage (page 86).

AFTER CARE

Many cases of severe failure with congestion and most cases of less severe type including those presenting fibrillation of the auricles respond well when treated along the lines previously indicated. The necessary treatment will naturally be less drastic from the beginning in the cases that were originally of mild type and in the severer cases a rigid system of rest and other parts of the regime may be relaxed as recovery happens. But there should be no thought of ending treatment by rest either in the severer or milder types of cases when the signs of increased venous pressure disappear. The heart has recovered no more than to regain its capacity to carry on a circulation adequate to the needs of the resting body; it still altogether lacks reserve. This reserve becomes re-established in greater or lesser degree by enforcing a further period of almost complete bodily rest and the rate of recovery is slower in cases originally of the severe type. No patient who has shown definite signs of venous congestion should leave his bed for several weeks after all these signs have vanished. A total period of six to eight weeks in bed is rarely too long; it is often too short to ensure maximal benefit. Patients who recover slowly or imperfectly must often lie quietly for three to six months or even longer. Cases that are allowed up too

soon quickly slip back to their former state, haste in treatment does not make for real speed. Prolonged rest should be rendered less irksome by arranging mental occupation, such as reading or being read to, wireless telegraphy, simple games or handwork, by permitting the patient to do more and more for himself as he recovers strength, by transferring him to a couch or chair for some hours each day, by employing massage especially to the legs, and later passive and gently resisted movements, and by an early return to fuller dietary. The steps taken towards increased liberty of action are to be small and not too frequent. At no stage of treatment should tobacco be allowed.

Return to Exercise Supervision —The guide to a return to bodily activity is the response to effort. A patient has recovered normal venous pressure and has been for a further period of several weeks confined to bed, at the end of this time he washes himself, sits up, and lies down without a trace of breathlessness. Such a patient may be permitted to walk across his room to a couch and, after some days, to walk a few minutes up and down the room. A schedule of increasing exercise cannot be laid down beforehand, but for weeks nothing must be done that gives a noticeable increase of respiratory rate or fatigue. Walking in the room may be replaced by short walks on the flat out of doors, the bed being arranged downstairs, or the patient carried up and down the stairs each day for the latter purpose a special carrying chair having two large wheels and long handles fore and aft is useful. As tolerance of exercise improves, exercise may be extended, until perhaps a few miles are walked on the flat at a moderate pace. Few patients who have experienced severe failure will advance beyond this point, the milder cases will, and should be encouraged to do so. However much the most favourable case improves, it is never good practice to allow these patients to call forth their full reserves or to press exercise to the point of bodily discomfort. Strenuous acts (page 285) should be forbidden for all time. The patient should adopt strictly healthy habits (page 283), he should report himself at frequent intervals, or at once if he feels unwell.

The hours in bed should be longer than is customary, and any tendency to fatigue may be counteracted by the patient lying down

chair,
■ more

DISEASES OF THE HEART

CH I

REFRACTORY CASES

The rule that confines patients to bed until after all signs of venous congestion have vanished has sometimes to be broken. The instance may be that of a patient already treated several times for congestion and responding very slowly to treatment. In the latest illness prolonged treatment brings the patient to an improved condition but one in which the veins and liver remain definitely engorged the condition remains stationary for perhaps another three or four months and the patient becomes restless under his confinement. These are very serious cases but some increased liberty of action has often to be allowed them.

CHAPTER V

SKIN COLOUR AND CIRCULATORY RATE

PALLOR

THE blood coloration of the skin is due mainly to blood in the net works of capillary venules in the subpapillary layers of the dermis. The cause of pallor is reduction of the amount of haemoglobin held within these vessels, and this happens when the haemoglobin content of the blood is low or when the amount of blood held in the vessels is decreased.

In the heart disease of adults persistent and conspicuous pallor, especially when the skin has a yellowish or subicteric tinge, is almost always due to streptococcal infection of the heart valves. Conspicuous pallor in children is often the result of active rheumatic infection. These pallors naturally involve both the skin and mucous membranes for they are associated with actual anaemia of the secondary or hypochromic type. Heart disease is occasionally accompanied by secondary anaemia caused by local infection or arising from haemorrhage. These are the main associations of heart disease and actual anaemia. Pallor in cases of heart disease should not be dismissed lightly, and must never be attributed to aortic regurgitation (see page 126) it is an invaluable guide to active heart disease or local infection. Dilatation of the heart in pernicious anaemia may be mistaken for heart disease with infection.

A bloodless condition of the skin is not infrequent in the advanced mitral stenosis of young girls, it is also frequent in the terminal phase of failure with congestion in these it does not involve the mucous membranes, and is evidently due, not to general anaemia but to actual emptiness of the minute cutaneous vessels.

Transient pallor occurs in two sets of circumstances. Firstly when there is a simple drainage of blood from the skin in syncope (page 96), such pallor may be intense, it is seen especially in the face of the standing or sitting subject, and is inconspicuous in

dependent parts. Secondly, it occurs in the rare attacks of vaso constriction described on page 60. In these the skin generally is affected and the hands and feet become pale although they lie below the level of the heart.

CYANOSIS AND ITS INTERPRETATION

CAUSE OF CYANOSIS

The cause of cyanosis is an increase in the amount of reduced haemoglobin, in any form, within the small vessels of the skin or mucous membrane. These reduced pigments appear blue through the skin. The condition may result when the blood pigment as it is brought to the integument, in which cyanosis is observed, is less oxygenated than is normal haemoglobin, or it may be due to an unusual loss of oxygen by the blood in its actual passage through the skin or mucous membrane. In some instances each of these two factors is in part responsible. We shall consider the pure examples first.

Cyanosis arising Centrally—In this variety of cyanosis, the blood as it leaves the heart contains haemoglobin in forms other than oxyhaemoglobin. It occurs in any condition in which air is prevented from reaching the alveoli in sufficient quantity, as in asphyxia or in capillary bronchitis, but these are usually impure forms of cyanosis, being associated with cardiac failure, and are dealt with later. The most familiar and striking example of a condition of pure central cyanosis is that of congenital heart disease in which the main lesion is a gross septal defect, allowing a quantity of venous blood to pass freely from the right heart into the aorta, thus short-circuiting the lungs and remaining unaerated. Another example is that in which methaemoglobinaemia or sulphhaemoglobinaemia occurs. Pure examples of cyanosis arising centrally are usually easy to recognise. Such cyanosis is often first suspected because it is conspicuous, because, although conspicuous, it is accompanied by little or no breathlessness while the subject is at rest, or because it is unassociated with signs of venous congestion. It is a general cyanosis, affecting not only the skin of face and hands, but the mucous membranes even within the mouth and the skin of the trunk, the last, however, often contains too little blood to yield an obviously cyanotic tint. A more important statement than any that precedes, because applying exclusively to this form of cyanosis, is that it occurs in superficial skin and mucous membrane that is

spontaneously warm or hot. Warmth or heat implies a rapid flow of blood through the skin that displays it, and cyanosis of central origin remains, however fast the blood travels through the skin.

In the form of cyanosis considered, the skin in which cyanosis is observed is usually warm, but if it is suspected in a skin that is cool or cold, it is my habit to provoke an increased flow of blood. Thus, if the lobe of the ear is deeply cyanotic and cool it can usually be made warm by local massage, preferably it should not only become warm but should display capillary pulsation (page 40) after the massage. Where cyanosis is central in origin the ear lobe remains cyanotic even though it becomes hot, it will become a little brighter in colour if the cyanosis in part originates centrally and in part peripherally. Another method is thoroughly to warm a cyanotic hand in water at 35°C (95°F), then to stop its circulation for seven minutes and finally to observe the tint of the reactive hyperaemia (or flush) which pervades the skin when the bandage is released. Normally and in cases of peripheral cyanosis presently to be considered, the colour of the quickly appearing flush has the bright redness of arterial blood, but where cyanosis is of central origin the flush appears bluer than normal. By using the signs and tests described cyanosis that is purely central can always be recognised at the bedside. These methods lead up to the diagnosis of many cases of congenital heart disease or of more occasional abnormality of blood pigment. An examination of the blood pigments spectroscopically differentiates between the former and the latter, so does an examination of the heart when this uncovers clear signs of congenital disease.

Cyanosis arising Peripherally—This variety happens when the rate of flow through the skin diminishes sufficiently. In its slow transit the blood is robbed by the cutaneous tissues of an unusually high percentage of its oxygen. On the one hand, slowness of flow may be confined to superficial tissues like the skin; on the other hand, it may result from failure of the general circulation. We shall consider the former instance first. Slowness of blood flow mainly affecting the skin is usually a response to low surface temperature. In interpreting cyanosis in various conditions, the temperature of the skin in which it is occurring is of very great importance. Cyanosis is generally noticed in the regions of skin that are most highly coloured, namely in the face and hands, and these are just the parts that vary conspicuously in temperature in response to changed environment. It is familiar knowledge that in cold weather these

parts of the body frequently become cyanotic in people otherwise regarded as normal, slight but distinct cyanosis of the skin is in fact a normal accompaniment of skin temperature between 15° and 25° C (59° and 77° F), and in many subjects this discoloration is exaggerated. The cyanosis is due to diminished rate of flow, and this in turn is due to constriction of the arterioles of the skin in response to the low temperature. Cardiac patients are just as prone as normal people, if they are not more prone, to display this purely local change. The unreliability of cyanosis as an evidence of general circulatory efficiency arises largely out of this fact and out of the difficulty of distinguishing in the skin between slowness of blood flow that is local and that is general, there is no constant relation between the blood flow to superficial and deep structures.

Cyanosis of Failure with Congestion—It is erroneous to believe that cyanosis in cardiac failure results simply from deficient aeration of the blood in the lungs, this is probably never the case. It is true that in advanced failure and very deep cyanosis the blood may leave the aorta in an imperfectly oxygenated condition, consequent upon its passing through alveoli that are airless or relatively so, owing to congestion or oedema of the lungs. But in the same cases the flow of blood through the skin is slow, and the visible cyanosis thus receives a peripheral contribution which is substantial. In states of lighter cyanosis the peripheral factor is the chief or the only one concerned. Cyanosis in cases of cardiac failure with congestion is found in skin or mucous membrane that for the most part is cool or cold.

When in a case of failure with congestion, cyanosis is found in a skin that is cool or cold, it is not very useful to try and ascertain if this cyanosis arises in part centrally. It is of greater interest to consider whether the blood flow has become slow locally in response to coldness of the skin or whether the slow circulation is primary and the cold skin has followed as an effect. In the latter case when the skin cools, its arterioles will likewise respond by contraction to the lowered temperature and the cyanosis will tend to be a little deeper for a given level of temperature than in the former case. But tint and temperature cannot be gauged by the unaided senses with nicety, and there are wide individual variations and so it is frequently found to be impossible to decide clinically between these alternatives, only one of which signifies a cardiac defect. In cardiac cases the evaluation of minor grades of cyanosis in facial or digital skin that is in fact cool or cold presents this recurring difficulty.

Though recognition that a patient seen for the first time has cyanosis may provide the earliest hint of cardiac disability, it will be apparent from this discussion that it is quite precarious to judge the general state of the circulation from this phenomenon. To use cyanosis when present as a sign of venous congestion is to blunder into accepting indirect and imperfect evidence, when the direct evidence in the veins, described in an earlier chapter, is available. Cardiac patients may develop cyanosis of central origin and yet display little systemic venous congestion, this is specially the case in bronchitis (page 255) and in oedema of the lungs (page 167). For similar reasons it is unwise to draw conclusions as to the general condition of the circulation from the absence of material cyanosis. Many patients, undoubtedly and severely congested, present very little obvious cyanosis. This is in part due to simultaneous pallor of the skin for the appearance of cyanosis is very dependent upon the presence of a full amount of blood pigment in the skin. Relative emptiness of the small vessels that give the skin colour is frequent in failure with congestion and thus the tint of the blood is concealed similarly it is hidden when, as is not infrequent, actual anaemia is present. On the contrary, full vessels or a polycythaemic state emphasise a deficiency in the aeration of the blood, as this is judged by skin tint.

MALAR FLUSH

A patch of colour of unusual depth over the region of the malar process has been thought to be a sign of mitral stenosis. It is not related causally to this disease. In very many young people who have been exposed frequently to cold wind, a bright malar flush is present, in many others the vessels in this region, though less conspicuous, possess less tone than do those of other parts of the face. The skin of this region of the face is therefore particularly prone to be involved in flushes of any kind amongst other causes venous congestion brings this region into prominence, rendering it bright and a little cyanotic. Because in young people failure with congestion is so often accompanied by mitral stenosis, the flush in question is often seen in them. Such value as this sign possesses is in directing notice to the cardiovascular system, and especially to the veins, in young people.

CUTANEOUS BLOOD FLOW

Cyanosis is a reliable sign of slow blood flow through the skin, provided that the entering blood is fully arterial. Actually a distinct

cyanosis in these circumstances indicates a very greatly reduced flow. And yet the flow may be very slow and the skin bright red, this is so when the skin is unusually cold (below 10° or 15° C, 50° or 59° F), the reason is that at low temperatures the blood can give up little of its oxygen. If the skin of the hand is blanched by pressure and the blood returns slowly, it is clear that rate of blood flow in the skin is slow, but a rapid return cannot be interpreted in the reverse sense, since it happens when the blood flow to the limb is arrested provided that the skin is already turgid with blood. Whether the skin is cyanosed or not, the depth of its colour is an uncertain gauge of the amount of blood flowing through it. In general the pale and not the flushed areas of the facial skin have the greatest flow of blood through them. The best gauge of blood flow in the skin is not colour, but temperature. Spontaneous warmth of the skin shows it to be possessed of good circulation, irrespective of tint or depth of colour.

Capillary Pulsation, so called, or visible flushing and paling of the skin with each heart beat, is a sign of locally dilated arterioles. It can be induced in any young and normal subject by heating the skin of face or finger. It is present in any circumstance of local or general vasodilatation, thus it is seen in inflammatory reddenings, large and small, and when the body is overheated, and in exophthalmic goitre. Its occurrence in aortic regurgitation is partly due to the high pulse pressure but chiefly to the vasodilatation commonly prevailing in this condition (page 126). It occurs only in warm or hot skin or mucous membrane. It has no clinical value apart from the evidence it provides for the state of the arterioles, consequently it should not be elicited by rubbing or warming the skin, but should be looked for as a spontaneous phenomenon, as such it is much more frequent than has been thought. It occurs less frequently under the nail than in the pulp of the finger and in the facial skin.

CHAPTER VI

CARDIAC ISCHÆMIA CORONARY THROMBOSIS

CORONARY ARTERIES AND CARDIAC ISCHÆMIA

THE CORONARY ARTERIES

THERE are usually but two coronary arteries, the right and left, supplying blood to the heart, these lead off directly from the aortic wall and although anastomosing in a limited way with each other, they anastomose scarcely at all with other arteries in the body. The coronary territory is an almost isolated territory, and such connections with other vascular territories as exist are minute and occur in the parietal pericardium and in the walls of the main vessels. The heart is really dependent upon two arteries for its supply, and obstruction of these is an event for which there can be little compensation by collateral circulation. It is difficult to believe, though there are some that do, that the muscular walls may be nourished by a backward stream through Thebesian veins. The two coronary arteries must supply the heart, whatever the amount of blood it needs. They must supply not only the normal sized, but also the greatly hypertrophied, organ, they must supply the muscle adequately, not only when its work is suited to the needs of the resting body, but when the heart is called upon, as it often is, to do a greatly increased quantity of work.

Pathological Anatomy—Disease interfering with the blood supply of the ventricular muscle occurs either in the length of the coronary vessels, or at their mouths. In the first instance the disease is a part of an atheromatous change in the vessels of the body generally, though the coronary vessels may be affected most severely, the narrowing results from gradual thickening of the vessel wall. Obstruction at their mouths is usually due to disease of the aorta itself, and this disease is atheromatous or, more commonly, the fibrosis of syphilitic aortitis.

Coronary vessels are also liable to sudden obstruction. Thus on occasion a free embolus or the end of an aortic valve vegetation is driven into an orifice and blocks its mouth. A much more frequent source of sudden obstruction is thrombosis; this occurs in the length of the artery and only in such as are already diseased. The changes in the muscle consequent upon coronary thrombosis are described on page 44. Coronary arterial disease is often accompanied by diffuse fatty degeneration or diffuse fibrosis of the ventricles (see page 250).

ANGINAL PAIN AND MUSCULAR ISCHAEMIA

It is known that when a coronary artery is acutely obstructed pain results; this pain is anginal in character and distribution. Pain of similar character is produced in the muscles of a limb that are worked when a tourniquet stops the flow of blood to them. It is probable that anginal pain is caused by a specific chemical or physico-chemical change in the musculature of the heart and that this is usually if not always consequent upon defective blood supply. Almost all if not all the phenomena relating to anginal pain are explained upon the basis of this theory; many of the phenomena have been explained as yet in no other way. While it is desirable to eliminate theory from this book as far as possible, the conception that muscular ischaemia, absolute or relative, is the cause of anginal pain, so simplifies an understanding of the circumstances in which this pain comes and goes, that it is advantageous to model description around this idea. When a coronary artery is blocked, a corresponding area of muscle loses its blood supply; there is a state of absolute ischaemia. When the supply to an area of muscle is reduced, or when it is incapable of increasing sufficiently to bring an adequate supply to muscle called upon to do an unusual amount of work, there is a state of relative ischaemia. That the first state gives rise to anginal pain is certain; that the second does is unproved but highly probable, especially from recently accumulated evidence. Theoretically, pain will occur when relative ischaemia affects the ventricular muscle as a whole or more usually in part. To be complete, the conception of ischaemia includes not only change in the structure of the coronary vessel, but the state of tone of coronary artery and arteriole; it allows that a deficient supply may be caused or contributed to by undue coronary tone and that a deficiency may be made good by coronary vasodilatation.

The long lasting effort to form a simple but complete classifica-

tion of patients who complain of anginal pain is not yet successful, but knowledge in this subject is being gained rapidly. Our ideas of angina have changed much recently, and will change more, because the symptoms of coronary thrombosis have been recognised and because this malady is no longer confused with allied maladies. There remains a number of comparatively rare and diverse types that are improperly understood and require further study. The chief types only can be described here.

CORONARY THROMBOSIS

Thrombosis of a coronary vessel is an accident occurring chiefly in men in their fifties or sixties, it happens occasionally in women, it is rare before the age of forty. The subjects usually present signs of arterial disease and, more often than not, high blood pressure. Coronary thrombosis is comparatively rare in cases of syphilitic aortitis. Although it not infrequently happens in patients who have already presented or are showing signs of failure, it is rare in rheumatic heart disease, and very rarely occurs in cases of auricular fibrillation. A considerable number of those who suffer have complained of earlier anginal symptoms, but some are affected without warning while in robust health. The malady is not infrequent in father and son or in two brothers.

PATHOLOGY

The cause of coronary thrombosis is to be found in disease of the vessel affected, such disease is usually but a part of a general arteriosclerosis. No coronary vessel is exempt from thrombosis, but the descending branch of the left coronary, which runs in the anterior interventricular sulcus, is affected much more frequently than any other, probably because of its relatively exposed position. The vessels next, but far less prone, to be affected are the circumflex part of the left coronary or the last part of the right coronary artery. The coronary arteries anastomose, though not very freely, with each other. If such a vessel as the descending branch is tied in a dog, a part of the ventricular wall loses its blood supply, becomes dark in colour and bulges with each systole of the heart. In many instances, and within a few minutes of ligation, the ventricle may begin to beat irregularly owing to the occurrence of ventricular extrasystoles, single or grouped, and in a number fibrillation of the ventricle comes and at once kills the animal. In others the heart continues to beat,

though less efficiently owing to part of its wall being thrown out of action. In a few instances where the territory is small the circulation is re-established through collateral coronary channels and the action of the muscle becomes restored.

In the human subject similar events undoubtedly happen. When the blood supply to a part of the ventricle is permanently arrested and the subject survives an area of muscle probably smaller than that originally supplied undergoes a fatty necrosis or 'myomalacia cordis'; becomes soft and fragile and may rupture. Ultimately the necrosed tissue is replaced by scar tissue which is firm in about eight weeks. The position and size of the muscle area involved naturally depends chiefly upon the position and size of the vessel that is blocked. In the usual instance of thrombosis of the descending branch the area affected is mainly in the front and apical regions of the left ventricle and is from 3 to 6 cms in diameter; it involves a corresponding area of septum. The wall of the heart is always more affected on its endocardial than upon its pericardial surface; the latter often escapes. Upon the former it is usual for blood clot to form, sometimes in an extensive and deep mass. The necrosis of the wall, the endocardial thrombosis and involvement of the pericardium are each responsible as we shall see later for special symptoms or signs.

SYMPTOMS AND SIGNS

Coronary thrombosis usually occurs when the subject is at rest sitting in a chair or lying in bed or actually during sleep; it also occurs at other times. The symptoms may be so slight and few that the patient seeks no advice; usually they are distinct or severe. The usual symptoms and signs may be divided into those occurring as immediate effects which include sudden death, pain and the manifestations of collapse and cardiac failure and into those that are delayed.

Death—In a considerable proportion of the cases death occurs very rapidly; there may not even be time for a complaint of pain. Many subjects are found dead. These sudden deaths are due in all probability to fibrillation of the ventricles.

Pain and Collapse—Pain is not invariable. When the pain comes it may be slight and may remain so; more often it rises rapidly to its culminating point and usually from then on is severe, often agonising; it is a continuous pain in the sense that it is unfluctuating. It is gripping or burning in quality; generally it starts over the

sternum, it may remain there or it becomes more diffuse as it strengthens and spreads to involve the left or right or both arms, but especially the left, or to the neck and jaw, or to the interscapular region, or to the abdomen. The lower sternum or the epigastrium is more frequently the starting point than in angina of effort. During the period of pain the patient may rest immobile, but frequently shows agitation and is restless or rolling about in agony, unlike the pain in the angina of effort, movement does not seem to increase it appreciably. Though its duration varies, the pain is characteristically long lasting, a feature which renders the malady so distressing. It lasts usually for one or more hours at its height, and declines gradually over a long period of many hours or several days.

The pain is usually accompanied by a state bordering on collapse, the patient being pale, haggard, anxious, weak, and often sweating profusely. When this pallor is associated with cyanosis the face assumes a light blue grey colour that has been thought to be characteristic of the malady. In this state of collapse nausea is frequent, and vomiting and movement of the bowel may occur. The pulse is small or it may be impalpable, the blood pressure is lowered, and much lowered in cases where it was originally high, the urine is scanty. The heart rate is usually increased but may be normal or slow in rate. This group of symptoms often resembles closely that of vasovagal attacks (page 98), though with quickened pulse the resemblance becomes less distinct.

Manifestations of Cardiac Failure—Symptoms and signs of cardiac embarrassment are frequent but by no means invariable, their degree naturally varies with the extent of myocardial infarction and the state of the remaining muscle. Breathlessness is almost invariable, often severe, and sometimes it is the only prominent symptom. Cyanosis is common. Signs of venous congestion may be relatively inconspicuous. They may appear quickly in any degree or may increase gradually. The heart sounds are indistinct, gallop rhythm is frequent, a systolic murmur may appear in the apical region, the heart may exhibit extrasystoles, paroxysmal tachycardia or auricular fibrillation, the last two, if present, will increase the circulatory embarrassment. Rarely, when the ventricular septum is involved, a third sound may be present. The signs in the lower extremities would be expected from the suffering of the patient. Cheyne Stokes' breathing is common. Signs of pulmonary congestion and oedema often pre-

dominate notably over those of systemic venous congestion, there is persistent cough with frothy and sticky, or serous and blood stained, sputum and râles and rhonchi appear over the lungs sometimes this oedema develops very rapidly and kills the patient

Delayed Signs (Fever Leucocytosis Ruptured Heart, Embolism, and Friction)—As is usual when a portion of a parenchymatous organ loses its blood supply and dies there is a febrile reaction. This fever is low amounting to 1 or 2° F. The temperature rises usually on the second day of the illness it may rise earlier it continues for a number of days and occasionally for as much as two or three weeks there may be no fever. Associated with the death of the muscle and its




FIG. 6-4. Diagrams of ECG at different stages of the disease.

c
e
c
of more leads

disintegration there is usually a leucocytosis also developing within a day or two and continuing with the fever. The white blood cells are from 12 000 to 15 000 exceptionally 20 000 or 25,000 per cubic millimetre.

The weakness of the ventricular wall which is at its height during the second week after the accident is responsible for the special accident of rupture of the wall haemopericardium and rapid death. The accident is rare before the fifth day and after five weeks.

As has been stated, clot usually forms on the affected portions of endocardium, and this clot may become detached in small or large pieces and give embolism. The emboli may enter and plug any artery in the body, but are more common in the systemic than in the pulmonary system. Embolic accidents give rise to numerous

complications, for example, hemiplegia, loss of circulation to a limb, splenic infarction and so forth, the presence of albumin or red blood cells in the urine consequent upon small emboli is often valuable for diagnostic purposes

When the pericardium is involved in the infarction and the area is on the front of the heart, friction sounds appear over the central and lower portions of the precordium. This pericarditis usually appears on the second or third day of illness and lasts a few days. Occurring in perhaps a third of the patients, it is a most important diagnostic sign and therefore should be sought for closely and daily where the diagnosis is in doubt.

DIAGNOSIS

The picture in its typical form is too characteristic to be missed by those who know of it. The central feature is pain of anginal type that continues long beyond the expected time and fails to yield to nitrites. These cases used to be diagnosed 'status anginosus'. When the pain is over the lower sternum or in epigastrium the trouble is apt to be mistaken for acute abdominal states, especially for perforated ulcer, tenderness and rigidity of the upper abdomen and signs of collapse with vomiting occur in both. The age of the patient will often be helpful in differentiating, for ulcer commonly occurs in the young, a previous history recognisable as clearly anginal on the one hand or gastric on the other is often to be elicited. In thrombosis the patient will often speak of pain as present over the middle or upper sternum also. Abdominal cases though pale are not cyanosed as are those of thrombosis, neither do they present the evidences of oedema of lungs or early signs of congested veins, either of which, like the breathlessness accompanying each, may be of great value in differentiating. The severity of the pain may be responsible for a diagnosis of gastric crisis, a mistake which should not occur, because in this the pain is spasmodic and not continuous. The same statement applies to biliary colic.

When pain is less severe or mild, acute indigestion is apt to be diagnosed, and the true state may not become manifest until embolism or heart rupture occurs, or angina later develops.

If pain is mild or absent, and breathlessness is the chief feature, the diagnosis made is of an attack of cardiac asthma (page 166), acute cardiac failure of unknown origin, or an anomalous form of pneumonia, and the fact that brief and even severe pain has been present will not always prevent the first and last diagnosis. Success

in diagnosis will depend mainly upon having the possibility of thrombosis in mind and in watching for the delayed signs of the disease, especially for the subdued fever, the leucocytosis red cells in the urine, and pericardial friction. The sudden onset of symptoms or signs of cardiac failure in a middle aged patient, unexposed to heavy work and afebrile at the time, should always arouse suspicion of coronary thrombosis. When there is coronary thrombosis but the diagnosis is in doubt, electrocardiographic examination will often resolve the doubt (Fig 6) but patients should not be taken to the instrument.

COURSE AND PROGNOSIS

The course of coronary thrombosis is very variable. The mortality in the stage of the acute illness is generally estimated at about half the patients. In a number death occurs at the onset or abruptly after a day or two, and presumably from ventricular fibrillation, others succumb to rapidly deepening failure of the systemic circulation or are suffocated by a rapidly developing oedema of the lungs, some are killed by rupture of the heart, others by rarer accidents such as the syncope of complete heart block, or gross embolism of brain or limb.

Those that recover may be severely crippled, in comparatively good health, or in any intermediate state, the ultimate condition depending largely on the condition of the remaining heart muscle and of the vessels supplying it. Patients previously suffering from heart failure or from angina who survive are usually severely damaged, failure is increased, congestion becomes established and progresses with varying fluctuations in intensity slowly but surely to a termination. Anginal cases are in general less tolerant of subsequent exercise. Curiously however, some such patients lose their pain after thrombosis, a possible explanation is that the original anginal pain arose from an area of muscle supplied by a narrowed artery that this vessel becomes thrombosed, and that the pain giving muscle then dies. Many of the survivors are left with very limited capacity for exercise owing to anginal pain. This may be *a remnant of the past, or may appear during convalescence or at any subsequent time*. A few patients are known to recover to good health, leading active lives or relatively active lives. The prognosis in these is uncertain owing to the development of angina, and owing to the not very infrequent recurrence of coronary thrombosis. Originally regarded as a hopeless malady, further observation,

especially more accurate diagnosis of the milder cases and the finding of scars of infarction in subjects unsuspected of the disease, has lightened the gloom of the previous outlook. Of those who convalesce well, some return to active lives for periods of five or more years and to live on in comparative activity for periods up to ten or occasionally fifteen or more years.

TREATMENT

Treatment of this grave condition is necessarily slight. The prominent symptom is pain, and for this morphia is the best remedy, the doses often need to be heavy ($\frac{1}{4}$ or $\frac{1}{2}$ grain, repeated until 1 grain has been given in four hours). Morphia is also useful in inducing the rest that is so desirable. Even morphia, however, frequently fails to give relief. Opium is sometimes more successful (1 to 3 grains). In cases of very severe continuous pain that is unrelied by opiates, light anaesthesia may be induced. Nitrites are generally useless. The patient is nursed in bed as a case of grave cardiac failure. Rest in bed should be continued for at least eight weeks, even in the milder cases, to ensure firm cicatrization of the ventricular wall, during the whole of this period the patient is to be guarded by day and night nursing, and helped in every way to avoid voluntary movement or effort. A number of patients have lost their lives, and especially those who have early recovered from symptoms, by neglect of these precautions. The after care is similar to that of cardiac failure or of angina with comparable symptoms.

CHAPTER VII

ANGINA PECTORIS

HEBERDEN'S ANGINA PECTORIS ANGINA OF EFFORT

THE malady originally described by Heberden under the term "angina pectoris" is one in which pain of characteristic type occurs during effort. It is by far the commonest form of malady in which anginal pain occurs. The pain may be interpreted as resulting when the blood supply to the heart or part of the heart, is limited and consequently inadequate when the heart is called upon to do work at a certain increased rate—a condition of relative ischaemia.

CLINICAL ASSOCIATIONS

It is essentially a malady of middle aged or elderly men though it occurs exceptionally in women of the same age period and occasionally in younger males. It is commonest in the sedentary and well fed classes, general arterial disease is frequent and about a third of the cases display high blood pressure. Disease of the coronary vessels, either narrowing of their mouths or thickening of their walls with

but none of these is the rule. A large fraction of the cases show no physical signs in the heart, and many present only a few distinctly abnormal signs, such as those of slight but definite cardiac enlargement, gallop rhythm, bundle branch block, and pulsus alternans. Some of the patients suffer from cardiac asthma. The malady is rarely associated with mitral stenosis. It very rarely occurs in cases of fibrillation of the auricles or of failure with congestion.

SYMPTOMS

Angina of effort not infrequently follows, immediately or later, upon an illness attributable to coronary thrombosis. Usually, how

ever, this is not the case, and the first complaint is of discomfort or of actual pain on effort. This is felt first of all, and in mild cases, on undertaking some accustomed act, such as running, or walking briskly uphill. The sensation appears usually over the middle or upper part of the sternum, though it may be over lower sternum, over precordium, or more generally over the front of the chest. It consists of a feeling of pressure, of gripping or of burning; it is smooth or continuous in the sense that it does not fluctuate in intensity from moment to moment or throb, though there may be slow changes in its intensity. When high in the chest and at the base of the neck it gives a sense of constriction. Its origin is usually in the lungs, but it may be in the heart or in the aorta. It is usually accompanied by dyspnoea, and is often associated with other symptoms of heart disease. It is usually relieved by rest, and by the use of nitroglycerine. It is sometimes relieved by the use of digitalis. It is sometimes relieved by the use of morphine. It is sometimes relieved by the use of strychnine. It is sometimes relieved by the use of arsenic. It is sometimes relieved by the use of iron. It is sometimes relieved by the use of quinine. It is sometimes relieved by the use of cod-liver oil. It is sometimes relieved by the use of cod-liver oil.

cases suffer very little pain because they become still at the first warning of its appearance. More often than not there is no radiation. If the subject presses forward the sensation usually grows in intensity, and may become more diffuse, spreading often to neck and jaw, to one or other arm to the abdomen or scapula. In the arm, and it is usually the left, the pain generally follows the inner aspect of the limb, descending to the elbow or rarely as low as the little finger. Numbness in the fingers or arm is often experienced at the time or subsequently. In rare cases the pain starts in jaw or arm and spreads from this to the chest. Some patients speak of being able, by continuing to exercise, to rid themselves of pain, but this is uncommon.

In severer cases the same discomfort and pain is felt but is elicited by smaller amounts of work, as by walking briskly or at a moderate pace on the flat the pain when it comes is apt to become more severe and to subside more slowly. In these and the milder cases the amount of work necessary to induce pain varies little sometimes astonishingly little, as when a man walks out of his house and is brought to his first standstill day after day at the same place.

countenance, which may be pale and moist. The blood pressure is little raised, the pulse rate is as usual or is a little increased or decreased. Salivation may be free. These attacks last a few minutes or, recurring quickly, are drawn out for a period of a half an hour or more. They leave the patient weak and tremulous, and the skin

of the chest, the intercostal muscles, pectorals, and sternomastoid often tender to touch

In attacks of pain produced by moderate exercise, it is common to hear that they occur more easily in cold air than in warm, walking out of a warm house into night air, or sitting down in a draught, may suffice to induce an attack. The excitement of animated conversation or anger provoked by dispute, is an adequate stimulus. In some cases exercise is particularly prone to induce attacks when it is taken directly after food, this and the frequency of flatulence in the same cases may lead diagnosis to the stomach rather than to the heart.

Some writers distinguish between the pain and the sense of tension or gripping in the chest. I am not satisfied that they are separate phenomena. Few patients speak spontaneously of both, and many state that the pain consists of a sense of violent pressure or tightness, or that a sense of tightness grows until it becomes a severe suffocating pain; many others will not allow that gripping occurs.

Expectation of death in the severer attacks is not often referred to unless the question is raised. Such a fear is a common association of many new, painful, or disturbing experiences and is usually irrational. There is no reliable evidence that a given sensation is responsible for it, or that it is more than an interpretation of a very unpleasant and new sensation or of an intolerable pain, it has at present little value in diagnosis. Much depends upon the mentality of the patient, and something upon what he has read or heard previously about his ailment.

DIAGNOSIS

The diagnosis of angina of effort turns, in subjects of suitable age and sex, almost exclusively upon the history, this should be elicited with the greatest care and will often require close and prolonged questioning. Though pain may appear first in one of many places, a history of pain over the sternum has more value than any other. It is a serious error to imagine that anginal pain must be severe and that slight pain occurring on effort cannot possess grave significance. In enquiring as to the type of pain, a most important point to elicit clearly is that the pain is uninterrupted, anginal pain does not throb or stab. Sometimes it is important that final answers should be deferred until, in the next attack, the patient has paid special heed to the particular points raised. An extremely important matter is consistency in the history of the response to effort. It is the rule that there is a given tolerance of work, constant within narrow

limits. Patients with this malady do not relate that on one day the attack is brought about only after walking briskly, and that on the next it occurs spontaneously while the patient is resting quietly and undisturbed. Single inconsistencies of this kind in the history may be explained occasionally by differences in external temperature, or excitement or fulness of stomach, but when inconsistencies are obvious and frequent, the case is rarely one of Heberden's angina pectoris. In angina of effort the pain of a given grade recurs with repetition of a particular act. Isolated severe attacks of pain are not characteristic of this malady, but more of coronary thrombosis or of acute gastric affections.

Minor inconsistencies are introduced when the attack is associated with large changes of pulse rate and blood pressure, a different type described in the next section, and major and frequent inconsistencies occur in cases in which the fault is not primarily in the heart.

The presence of signs of definite heart disease will help to establish the diagnosis in cases of doubt, but it should be understood that in most cases gross evidences are not forthcoming. From this standpoint, some physical signs that help the identification of cardiac pain, and which sometimes stand alone, are described in Chapter XXV.

A SPECIAL ANGINAL SYNDROME

Anginal pain in resting subjects is usually an indication of grave disease. Thrombosis of a coronary vessel may be responsible for a single severe attack, a second thrombosis may cause a similar attack. In patients who suffer from Heberden's angina pectoris, and in whom the disease progresses far enough, pain may be aroused eventually by such small stimuli that the patient takes to bed and still suffers attacks of pain.

The commonest type of grave spasmodic angina met with in bedridden patients is different and has certain special features which make its separate description desirable. The cases are usually young or middle aged male subjects suffering from aortic regurgitation, females also suffer, and the same syndrome is also found apart from aortic disease. The aortic disease may be rheumatic or syphilitic. The heart is greatly enlarged. The coronary vessels may or may not be obstructed. Some of these patients give a clear history of having suffered from the angina of effort. They are for the most part bedridden on account of pain and other symptoms. At the best they are capable of only slight exertion, for attacks of pain

are easily induced by walking. They experience severe attacks of pain while resting in bed. This pain is of the usual type, and has the usual and varying location and distribution of anginal pain. It may or may not be associated with a feeling of intense anxiety. The attacks may occur at any time, but are especially prone to occur in the early hours of morning, often waking the patient from sleep. In the day they may be provoked by food or by the thought of food. They last a few minutes to half an hour or more. The patients usually state that the pain is accompanied by violent throbbing of the chest and neck and a feeling of tenseness in the head. The face in the attack is usually flushed, sometimes deeply, sweating is often profuse, rapid breathing is the rule, but there is neither cyanosis nor râle in the chest. The heart rate is raised to 100 or 140 beats per minute, the systolic blood pressure is raised from normal limits to very high points, usually to 200, sometimes to 250 mm Hg or more, these phenomena distinguish the attack. The pain, usually severe, is almost always relieved by amyl nitrite, though blood pressure does not always fall. In the attacks the rise of pulse rate and of blood pressure begins before the pain starts, evidence that the attack originates in a disturbance of the cardiac accelerators and the vasomotor nerves, whereby increased work is thrown upon a defective heart. But the relation between extra work and pain is inconstant, and suggests that the vasoconstriction while raising blood pressure involves not only the abdominal but also the coronary vessels.

DIAGNOSIS

The presence of clear signs of serious heart disease, and the severity of the attacks at rest, makes the diagnosis of a grave form of angina pectoris certain. The associated disturbance of pulse rate and blood pressure may be suspected from the patient's story of violent throbbing, but it should be confirmed or otherwise by observations in the actual attacks. This malady is not to be confused with Nothnagel's syndrome (page 60).

COURSE AND PROGNOSIS

The subjects who display Heberden's angina are for the most part middle aged or elderly when the malady first displays itself. The malady is associated with processes in the vessels that are degenerative and usually progressive. The usual course therefore is towards greater limitation of exercise by pain. To avoid pain, less and less

exercise is taken as time goes by, eventually pain occurs on such slight effort that the patient is driven to his chair, his couch, or his bed and is never secure from attacks of pain, which now tend to be of increased severity. A few pass into an anginal state and die in it, though this state more frequently results from coronary thrombosis. The average life though very variable, is about six or seven years from the beginning of the attacks. In very many cases the progress of the disease is slow, and men who experience their first symptoms in the middle forties live on to reach the middle fifties or sixties. The course of angina is not always steady in its progression. Years may pass with little or no perceptible change of condition. In a number of instances the patients acquire signs of cardiac failure, with or without fibrillation of the auricles. Breathlessness rather than pain now curbs exercise. These patients as a rule never suffer afterwards from anginal pain, nevertheless the onset of failure with congestion in them is a very serious event and terminates life usually within a year. The fact that anginal patients are the subjects of general arterial disease and are prone to its various accidents, renal and cerebral, should be remembered. The end does not always come through failure of the heart.

The prognostic significance of Heberden's angina in individual cases should be regarded first from the standpoint of the amount of exercise needed to induce pain. Cases that have mild pain on walking briskly uphill, pain that ceases soon if they come to rest, are in general favourable cases. Those in whom pain of longer duration is induced by walking slowly on the flat are in general much more serious. When the patient becomes sedentary in habit and experiences pain at frequent intervals the end is not far distant. Patients who have taken no care of themselves, have worked excessively, or have recently been subject to acute infection, are apt to hear a worse prognosis given than they deserve. Judgment should not be hasty, as tolerance of exercise changes appreciably from time to time in some cases, and many react well to treatment. Secondly, prognosis should be based upon the rate of progress as this is elicited in the history or by observation. The severity of pain on effort is not often of much value, apart from its indicating the carelessness with which the patient nurses his malady. An element of great uncertainty is introduced by the occurrence of coronary thrombosis, this accident cannot be foretold, but, though it may happen at any time, most of these patients escape it. Sudden death will come to many of these patients whether they acquire a throm-

be guarded against undue excitement, mental or physical, and from coldness in and out of doors. The food should be small in amount, easily digested, nutritious and taken dry (page 286, Diet II) a rest of an hour should follow the heavier meals. The use of tobacco should be forbidden, and not much tea or coffee should be drunk. It is important to treat flatulent dyspepsia, for this is often responsible for attacks of pain or for their easier provocation. If attacks of pain are easily induced, as by very quiet walking exercise, a period of several weeks' rest on couch or bed benefits. Usually the pain disappears in such patients from the time they go to bed or within a few days, simply as a result of rest. It is customary to find after several weeks' rest and when the patient is up and about again, that more exercise can now be taken painlessly than previously. It is as though there were originally some cumulative process in the heart, as though the muscle were in a sense tired, and recuperated with rest. If attacks of pain are occurring apart from effort, very close enquiry should be made to ascertain the cause the trouble may arise out of a concealed source of irritation or worry, sometimes the patient will not recognise the cause until he is helped in his search for it.

Many of those patients who work or exercise during the day, become unduly fatigued or a little exhausted in early evening. Rest and a glass of wine often comforts and revives them, but they should do less.

Amyl Nitrite capsules (2 to 5 minims) should be carried always by those suffering from Heberden's angina though they should not often require to use them, the proper remedy is to cease exercising at the first warning of pain the nitrite is rather for the emergency, for an attack of pain that is provoked unexpectedly or is of disturbing severity. Alcohol may be used in similar emergency, but the objection to its regular use in attacks will be obvious. Those who are frequently forced to use capsules should not be walking. Attacks in resting people may be treated fully by nitrite and, when severe and prolonged a number of capsules may be broken and inhaled in succession, until the paroxysm of pain ends or it is clear that the nitrite fails to take effect. Often a single inhalation that flushes the face well suffices to stop an attack. Throbbing under the drug is sometimes uncomfortable, headache often follows its use, patients whom it relieves never refuse it on these grounds. The action of the drug upon the vascular system is fleeting, but its effects on pain outlast this action. Its potency is probably due mainly to its

powerful dilatation of the coronary vessels rather than to its effect in lowering blood pressure. It fails to relieve in cases where the coronary circulation cannot be increased as when there is thrombosis in the vessel supplying the affected area.

Nitroglycerine is a very valuable remedy in the stage when a progressive angina of effort has brought a man to his chair or couch. It is then used to ward off attacks of pain that would otherwise occur or to lessen their severity. The dose is 1/100 grain and a tablet when chewed and swallowed begins to act within one or two minutes and its action lasts for very many minutes. This dose may be taken at intervals of a half hour or even less for periods of several hours if necessary. The transaction of urgent business or the visits of friends are sometimes rendered possible only by its use. It should not be employed with the idea of simultaneously allowing greater bodily activity. The drug like any nitrite may occasion throbbing giddiness and headache.

Theobromine (in 10 grain capsules) or *euphyllin* (tablets containing 1½ grains) may be given three or four times daily with benefit in cases with frequent attacks at rest. These drugs are also vaso-dilators and are more prolonged but less certain in their action than the nitrites. There are patients whom they suit particularly well.

Ammonium bromide (15 or 20 grains) taken thrice daily for periods of a week or ten days with intervening periods of rest is used similarly in the resting case with frequent attacks. This method of dulling the nervous system is especially valuable in restless and sleepless subjects.

Luminal (1 or 2 grains) twice a day may be used for the same purpose. In the agonal state when severe attacks recur rapidly morphia is used and on occasion light anaesthesia is required.

Surgical Interference in the form of removal of the sympathetic ganglia at the root of the neck (left inferior and middle cervical ganglia) has been proved to abolish the pain of angina in a number of instances. It is an operation sufficiently dangerous to render its legitimacy doubtful. Its recommendation is possibly justified in intractable cases in which severe attacks of pain occur daily despite all other remedies so also is the injection of alcohol into the immediate neighbourhood of the upper thoracic posterior root ganglia an operation only to be undertaken by those skilled in the procedure.

SPECIAL CASES

This and the preceding chapter describe the chief types of case in which anginal pain is the main symptom of a grave cardiac malady. If these types are fully understood few cases of grave angina will be overlooked and most of those patients in whom the heart is sound or relatively so will be separated from these.

Neurasthenia —There is a group of patients containing more women than men and well known to all medical men they present the picture of long drawn and constant ill health. These people never feel really well they are depressed often irritable and in a state of constant fatigue. They present a great variety of other symptoms many of which are trivial and when all efforts to arrive at a more precise diagnosis have failed become classed under the term neurasthenia. Some of these patients are actually suffering from unrecognised and chronic local infections in others no such lesions are ever discovered. The condition of the nervous system is such that any pain is poorly tolerated it may be that the threshold for pain is low in these people. Be that as it may some complain of spasmodic pain referred to the chest and the state of the heart comes under discussion. Grave angina in women under forty five years of age should be diagnosed infrequently and very rarely in younger women or much younger men who present no clear signs of cardiac disease such as manifest enlargement of the heart or aortic disease. Symptoms of an irritable nervous system of constant ill health between attacks of pain will naturally tend to deflect attention from the heart. An incorrect diagnosis is often saved by detailed consideration of the pain which is more frequently referred to the precordium than to the sternum or is constituted by continual soreness or is stabbing in quality. Often respiration is rapid in the attacks and the patient very restless. Usually too there is no relation or but a very imperfect one between the pain and effort.

Tobacco and Angina —The occasional heavy smoker is affected by attacks in which there is a sense of constriction in throat or chest they may be accompanied by pallor nausea giddiness and weakness. Occurring in early adult life and ending when tobacco is discarded they are probably to be ascribed solely to smoking. An oppression in the chest or slight sense of sore constriction is not uncommon amongst heavy smokers when they take active exercise. In elderly men it is probable even when tobacco is evidently provocative that there is an underlying predisposition to angina. It is

suspected, but not proved, that heavy tobacco smoking can lead to degeneration of arteries including the coronary vessels, it has also been supposed, though upon insufficient evidence, that excessive smoking can cause spasm of the coronary arteries

Tachycardia and Angina —Occasional cases of paroxysmal tachycardia, in which the rate of beating is high and paroxysm prolonged, complain of characteristic anginal pain in the attack. This example is interesting in that the heart is overburdened, not by any rise of arterial pressure, but merely by an excessive rate of beating

Severe Anaemia, Hyperthyroidism —Attacks of anginal pain occur sometimes in any severe anaemia and disappear when, under liver extract or otherwise, the anaemia is cured. Here there seems a definite relation between anginal pain and nutrition of the heart. It is not clear in this instance, or in the last of tachycardia, whether there is or is not an underlying disease of coronary vessels and a consequent predisposition to attacks of angina. A similar statement applies to cases of hyperthyroidism, in which anginal pains may continue until the thyroid has been treated adequately

Vasoconstriction provoking Thoracic Symptoms —When adrenaline is injected intravenously into healthy subjects, pallor appears in the skin of the body generally and the pulse rate and blood pressure are raised. In conspicuous reactions there may be strong palpitation and throbbing in the neck, a sense of constriction in the chest or actual pain across the sternum, closely resembling the anginal phenomena. A corresponding group of symptoms and signs develops spontaneously in rare cases of paroxysmal hypertension, and has been described in association with suprarenal tumour; the attacks are thought to be due to natural adrenaline released, intense headache, nausea and vomiting may be added

Nothnagel described under the unfortunate term "angina pectoris vasomotoria" patients with similar manifestations. The malady is a very rare one, so rare that no cases of precisely the same kind have been described since 1867, and it is here discussed chiefly for historical reasons. Nothnagel's cases were not heart cases, they were chiefly men, otherwise healthy, and in their thirties and forties, in whom for a limited period attacks were provoked by exposure to cold or by bouts of drunkenness. The attacks, which lasted a quarter to one hour, started with blanching of the skin, the hands and limbs being involved conspicuously and becoming numb and cold. The pulse was narrow and small, its rate unchanged or a little slow. These signs preceded anxiety or actual fear, and severe

palpitation. In one case, the only elderly subject of the series, pain of frankly anginal type appeared. A few cases of a similar kind have been described with anginal pain but raised pulse rate.

The real importance of the cases here grouped together is not that they confuse diagnosis, but that they suggest that vasoconstriction, by throwing a considerable burden on the undiseased, though not necessarily wholly efficient, heart, can determine secondary symptoms of considerable thoracic distress, or actual angina.

CHAPTER VIII

MINOR PULSE IRREGULARITIES

SINUS ARRHYTHMIA

THE normal pulse may be definitely irregular in rhythm, this is frequent in childhood and adolescence owing to what is called sinus arrhythmia. All the chambers of the heart are involved in the irregularity, which though often inconspicuous may be pronounced. It is nearly always related to respiration, the pulse quickening during inspiration and slowing during expiration (Fig 7). Occasion-




FIG 7—Simultaneous electrocardiographic, radial, and respiratory tracings. The irregularity is of sinus origin and accompanies natural respiration. The whole heart participates in the irregularity. The heart rate rises during the phase of lowered intrathoracic pressure and falls during that of raised intrathoracic pressure.

ally the relation to respiration is indistinct, in such cases the nature of the irregularity may be tested by asking the patient to breathe more deeply, sinus irregularities then become more distinct and become obviously respiratory. The significant point of this test is that the original irregularity is completely displaced and not that respiratory irregularity appears, for it is the rule for it to appear when young people breathe deeply. These sinus irregularities result from

inconstancy of vagal tone, and are almost always associated with relatively high vagal tone, and therefore with relatively slow action of the heart

In the days before irregular heart action was analysed and the various types distinguished, all irregularities of the heart were regarded as of grave significance, and sinus arrhythmia, though occurring normally in childhood came to be regarded as particularly associated with meningitis and other grave maladies. This example, one of many in clinical medicine, clearly illustrates how seriously signs may be misinterpreted until properly investigated. To day the importance of recognising this normal phenomenon, sinus arrhythmia, lies almost exclusively in its possible confusion, during simple palpation of the pulse, with other forms of irregular heart action, and especially with fibrillation of the auricles, in the latter, when the ventricular action is slow the irregularity is often inconspicuous.

rate is raised in exercise in fever, or by drugs such as amyl nitrite. Sinus arrhythmia very rarely occurs with pulse rates as high as 100 per minute

INTERMITTENCE COUPLING AND GROUP BEATING

Certain kinds of irregular heart action are relatively simple in form a given phase being repeated over and over again, there is order in the arrangement of the irregularity. The simplest disturbance of this class is the intermittent pulse

Intermittent Pulse—An occasional pause of pronounced length, interrupting an otherwise regular pulse, may result from an extrasystole, or from a dropped beat due to heart block. The former is common and the latter rare

The extrasystole is a disturbance consisting of a premature beat. This beat may arise in the auricle, in which case the premature contraction will replace the natural beats of both auricle and ventricle that are nearly due (Fig 8) or it may arise in the ventricle, in which case the ventricle only will be involved, its premature beat replacing the rhythmic response of the ventricle that is shortly

expected (Fig 9) In either case the early beat of the ventricle will expel a smaller quantity of blood than usual into the aorta, and the




FIG 8—Venous radial and electrocardiographic curves showing an auricular extrasystole. The extrasystole is well shown in the radial curve as a premature beat and in the early waves *a* *c* in the venous curve. In the electrocardiogram the early beat is represented by deflections *P* *R* *S* and *T*. The beat arises in an unusual auricular focus; it spreads abnormally through the auricle and consequently causes an abnormal *P* deflection. Arriving from the auricle it spreads in normal fashion throughout the ventricle and thus the ventricular deflections are of usual form.




FIG 9—Venous radial and electrocardiographic records showing a ventricular extrasystole. The


of the record

corresponding pulse beat will be weak or will fail. When the early beat fails the pulse intermits; a movement is detected over the

heart and a first but not a second, sound is heard the aortic valves have not lifted This is the extrasystolic intermission

In heart block, intermittence is produced by failure of the ventricle to respond to a rhythmic auricular beat (Fig 10) There is no premature impulse to detect the heart is motionless and silent during the whole pause Thus the intermittence resulting from extrasystole is easily distinguished from that of heart block by attentive examination

Coupling and Group Beating—The pulse may intermit occasionally or at regular intervals If the regular intervals are close together, the pulse beats arrange themselves in simple groups Thus, if every third rhythmic beat is replaced by an intermittence, the pulse presents a coupling of its beats if every fourth beat is



Intervals the pulse intermits

replaced the pulse beats occur in groups of three Coupling of the ventricular beats will appear if an extrasystole replaces alternate beats of the ventricle and the pulse will also display coupled action in the same circumstance if the premature beats are strong enough to yield pulse waves but if the premature pulse beats fail the pulse rate will be halved

It will be clear that very similar groupings of the pulse beats and halving of pulse rate can also occur when the ventricle fails to respond to the auricle at regular and frequent intervals

In distinguishing between these two mechanisms at the bedside we depend on being able to recognise the extrasystole when it occurs, by remarking its prematurity and, when it is weak enough, by noting its failure to produce a second sound In cases presenting

difficulty the heart rate should be raised by exercising the subject, when the irregularity will generally disappear, a stethoscope is now placed on the impulse and the form of the first returning irregularity is noted. In extrasystole it begins with a premature beat, in block it begins with an unusually long pause.

Palpitation from Extrasystoles —Symptoms may accompany intermittence or group beating when these result from extrasystoles, but not when they result from heart-block. In a large number of the subjects the extrasystole passes unnoticed, on the other hand it is a common cause of what patients term "palpitation." The symptoms are more prominent in the young, in women, in people of nervous temperament, or in general ill health or fatigue. With the extrasystole there is a feeling of uneasiness, as of a void, in the chest, while the succeeding beat is accompanied by a shock to the chest wall, and often by a feeling of gripping in the throat. By calling attention to the heart they awaken anxiety. When numerous they may cause much discomfort; occasionally they bring actual distress, the skin becoming cold with sweat and the subject faint.

Palpitation arising from occasional extrasystoles can generally be identified in a patient's account of himself, if he can clearly describe his sensations. When extrasystoles are more numerous, the sensations are difficult to distinguish from those in paroxysmal fibrillation (page 82).

Associations of Extrasystoles —Extrasystoles are a very common phenomenon, occurring at all ages, but especially in the later periods of life, it is probable that they occur in most people who live into middle life or beyond it. A few extrasystoles may occur each day, or from time to time, or they may persist for hours, days, or years. Numerous or persistent extrasystoles are frequent in cases of chronic rheumatic heart disease, hypertension, and chronic gastrointestinal disorders. They may occur temporarily during the exanthemata or as associations of local infections. They are often provoked by excessive smoking, sometimes by large doses of digitalis or its allies. In those subject to them, extrasystoles are particularly prone to occur during rest after exercise, after heavy meals, and at night shortly after going to bed. In other subjects they occur only in the upright posture.

Prognostic Significance of Extrasystoles —That extrasystoles are found in cases of serious heart disease and that the mortality amongst heart patients displaying them is high, are facts which provide no evidence that extrasystoles have grave significance. The belief that

extrasystoles are due to a disturbance in the myocardium rather than in the nervous system is also to be left out of account in judging them. To assess the prognostic significance of any phenomenon it is important not to be misled by associated phenomena or by ideas of causation. Prognostic significance is to be gauged purely upon an experiential basis with carefully controlled evidence: the prognostic significance of many clinical phenomena is still misjudged because these simple rules are broken. If a random series of subjects having extrasystoles and a random series without them are selected and the two compared it will be found that the death rate in the extrasystolic series is the greater. Such evidence is worthless in gauging the significance of extrasystole because amongst the extrasystolic series will be a greater number of patients having serious cardiac disease or other malady and such disease and not the extrasystole will possess the real significance. To judge the extrasystole it is necessary to compare with the first series a series of selected cases presenting all the phenomena of the first series and presenting them with equal frequency namely enlargement valvular disease infection etc (only excepting the extrasystole itself) or a selected series of patients having extrasystoles but apparently otherwise in full health may be compared with a series of normal controls of like age etc. When such comparisons are made the significance of extrasystoles is found to be so slight that its use in determining the prospect of life becomes practically negligible.

When extrasystoles are found the heart amongst other organs should be scrutinised closely. If no further signs are detected the extrasystole can be neglected; if other signs are found the prognosis will be based on these: the extrasystole has served its purpose and can again suffer neglect.

There are rare and special cases in which extrasystoles are persistently present in long groups of successive beats: it is sometimes clear that such groups increase the energy expenditure of the heart sometimes obviously embarrassing its action: previous statements do not refer to cases of this kind: the prognosis of which is less precisely known but which is very probably nearer to that of auricular fibrillation.

Whether extrasystoles will persist or not is difficult to state in the individual case. Many people are temporarily affected by extrasystoles which do not reappear or reappear at rare intervals: in far less numerous instances the disorder may be present constantly for long periods of years. The answer to the question here raised must

take any factor known to be provoking the beats, and the prospect of its removal, into consideration

Treatment of Extrasystolic Palpitation —When extrasystoles produce no symptoms they require no treatment Their presence should rarely limit a patient's ordinary activities Restrictions are called for occasionally by symptoms that are more than slight The patient if anxious, should be reassured, this encourages indifference to the beats Sometimes the provocative cause, namely, excessive smoking, overwork, flatulence constipation, or infection, can be detected and removed The only drug that frequently stops extrasystoles is quinine given in doses of 2 to 5 grains twice or thrice daily Its use should be restricted to cases that are much disturbed by the beats Digitalis should not be used Ammonium bromide given in doses of 15 to 30 grains or more a day will often lessen or abolish the symptoms of extrasystole, this remedy is particularly valuable when patients who have extrasystoles are much upset or kept awake by palpitation

Heart block is dealt with further on page 92

CHAPTER IX

TACHYCARDIAS

SIMPLE TACHYCARDIA

SIMPLE tachycardia is a term used to indicate a natural action of the heart at an increased rate. Its cause is increased rhythmicity of the sino auricular node (Fig 17, page 93), the heart's pacemaker. In physiological circumstances this rhythmicity is increased (a) by altered innervation by loss of vagal or increase of sympathetic tone, (b) by change in the chemical environment of the heart, as when the blood becomes more acid or when there is an increase in its adrenaline content or (c) by alteration in the physical environment, a notable example of which is the reaction to increased temperature. Thus simple tachycardia is not primarily cardiac in origin.

In the human subject simple tachycardia (Fig 11) is seen in a variety of conditions and the exact manner of its production is not




FIG 11.—Electrocardiogram from a young woman who showed persistent tachycardia of an known origin. The rate at rest was 130 per minute rising to 180 with slight exercise. It is a simple tachycardia, each cardiac cycle being represented by normal auricular and ventricular deflections. Time in $\frac{1}{4}$ sec.

always known. As a physiological variation, the heart rate may be persistently above the average (exceptionally as high as 90) in the healthy subject, the natural rate is faster in children than in adults. It increases during exercise (to as much as 180), during mental excitement, and when heat is applied to the body. It is increased by fever, tachycardia is present in all acute febrile diseases, though to a less extent in those affecting the central nervous system and in

typhoid fever. Rapid heart action is prominent after hæmorrhage and in shock, it is a natural consequence of low blood pressure. It follows lesions of the vagi, especially the right nerve. It is a special feature in exophthalmic goitre. Simple tachycardia often accompanies local infections, whether fever is present or not, one of the most notable instances being pulmonary tuberculosis, but small and often hidden foci, such as tonsillar infection or pyorrhoea, may be accompanied in afebrile subjects by a conspicuous increase of heart rate (to 130 resting). It occurs in poisoning by atropine (or bella donna), this and nitroglycerine (the latter taken by soldiers in the form of cordite) are notable examples, it is seen in acute and chronic alcoholism. It is a chief feature of what is termed the "effort syndrome" (page 158). Amongst valve affections high rate of beating is frequent in mitral stenosis and in aortic regurgitation.

As a temporary disorder simple tachycardia is often encountered in those who present signs of nervous irritability, in these it is provoked by exercise or emotion. Transient tachycardia occurs similarly in many who are convalescent from acute illnesses, and in many of the instances already given, in which the heart rate is already raised, a physiological cause of acceleration often provokes an exaggerated response.

In most of the instances cited, not only is the rate of beating increased, but the action of the heart is more forcible and its output increased, a state sometimes termed "augmentation."

SYMPTOMS


Augmentation of the heart beat is one of the most frequent causes of 'palpitation' as patients use this term. The unpleasant throb in chest and neck may be accompanied by restlessness, agitation, and occasionally by nervous prostration.

SIGNS ASSOCIATED

When the heart's action is augmented the impulse becomes diffuse, or very diffuse, giving rise to the false idea that the heart is dilated (see page 109). Systolic murmurs frequently appear at base or apex and strengthen the erroneous impression of cardiac disease. The diffuse impulse and the systolic murmur are expected manifestations of a heart overacting in any circumstances.

PAROXYSMAL TACHYCARDIA

Paroxysmal tachycardia is caused by the heart acquiring temporarily a new rhythm usually arising from an abnormal focus in the heart and uncontrolled by the heart nerves and by other physiological factors that influence the natural heart rhythm. The nature of the new rhythm is unknown. It is a pathological type of beating occurring in attacks that begin and end quite abruptly (Fig 12). In the attack the ventricular rhythm is quite regular. Different



paroxysms vary in rate between the wide limits of 110 and 220 per minute but the usual rates are between 160 and 200. They may last a few seconds, a few hours, a few days, a week, or a little more. The usual duration is an hour or two, or a day or two.

CLINICAL ASSOCIATIONS

Paroxysmal tachycardia may occur at any age but its highest incidence is in the twenties and thirties. It often occurs in subjects who present no signs of disease or relevant history of past illness. Of associated conditions, rheumatic heart disease is the most frequent. Thus, paroxysmal tachycardia is relatively common in mitral stenosis and in enlargement of the heart of rheumatic origin. It appears to be associated from time to time with chronic local infections in other cases with flatulent dyspepsia.

In those predisposed, paroxysms are often provoked by exercise or emotional disturbance, sometimes by flatulence, rarely by the assumption of the erect posture. In a given patient the length of the paroxysm is fairly consistent. In some cases they are infrequent and of long duration; in others they are frequent and of short duration.

SYMPTOMS

The symptoms proper to paroxysmal tachycardia are those resulting from the rapid and abnormal action of the heart. They vary chiefly with the rate of the heart beat and with the reaction of the nervous system. When the attack is brief and the heart rate is not extreme, and especially if the patient is of phlegmatic type, it is not unusual for the attack to pass unnoticed. The first symptom is usually precordial discomfort, beginning abruptly, and described as slight or violent palpitation. A sense of flutter in the chest and throbbing in the neck are common. In sensitive or nervous patients there is alarm, the subject becomes pale, sweats, and feels tremulous and weak. The end of the attack is marked exceptionally by a sharp stab of pain in the chest or by one or more forcible heart beats, but as a rule the patient speaks only of quick relief from such symptoms as have been present.

CARDIAC REACTIONS

The reaction of the heart varies with the rate of beating, the length of the attack, and the original state of the heart's reserves. Rapid action causes the heart to expend more energy and decreases its rest period (see page 23). When the attack is brief or the heart rate not extreme, and the heart's reserves are relatively normal, there are no serious disturbances. The X rays usually show the heart to diminish a little in size when the attack begins. This is the common type of case. The most serious symptoms occur when the cardiac reserves are already diminished, for a paroxysm of rapid tachycardia then exhausts them. This leads to the rapid development of acute failure with congestion. The veins in the neck begin to swell, the cardiac impulse moves perceptibly outwards, the impulse becomes diffuse, the liver margin begins to descend, the colour of the patient alters, the countenance becoming cyanosed, the eyes sunken, the expression haggard, the patient is breathless. Within a half-hour the signs may be conspicuous. Breathlessness becomes
 tion is seen in it. The abdominal muscles become rigid. Aching pain

progress until extreme In the latter case the cardiac impulse continues its outward movement until it reaches the mid axillary line In long lasting paroxysms exhaustion is prominent In the last stages of steadily progressing attacks a cough accompanied by frothy and sometimes blood stained sputum develops with signs of engorgement and oedema of the lungs These steadily progressing attacks occasionally terminate in general dropsy delirium and death Almost all attacks end in a resumption of normal beating in a very few sudden death occurs When the attack stops the patient experiences immediate relief breathing becomes free and the remaining symptoms soon disappear the signs of engorgement disappear with remarkable quickness The heart resumes its original size within a few beats the liver recedes more slowly in the space of hours Flatulence and weakness are often present for several days after an attack and a sense of exhaustion and soreness of the chest may persist

In rare cases and these usually known to be predisposed a paroxysm may start an attack of characteristic anginal pain slight or severe and radiating

AURICULAR FLUTTER

This remarkable condition in which the rate of auricular beating is usually as high as 260 to 320 per minute is closely akin to auricular fibrillation It is believed to be caused by a wave of contraction circulating regularly and continuously through the auricle in this rapid rhythm (circus movement) it is due to a fault in the heart itself the rate is so high that the ventricles rarely respond at more than half the rate (Fig 13) The usual ventricular rate is 130 to 160 that is to say half the auricular rate There are rare cases in which auricular flutter occurs in short paroxysms and it may then be confused with simple paroxysmal tachycardia Usually if untreated it continues unchanged for months or years The clinical associations and the symptoms of this persistent condition are similar to those of lasting auricular fibrillation (see pages 81 and 82) A history of fainting attacks is not uncommon in flutter

DIAGNOSIS AND DIFFERENTIATION OF TACHYCARDIAS

Cases of conspicuous tachycardia present themselves to medical men in one of several ways

A patient comes complaining of attacks of palpitation and the

action of the heart is at the time normal. The distinction between simple tachycardia and paroxysmal tachycardia which is necessary in such a case can often be formed on the basis of history alone. The most important point in evidence of paroxysms is a perfectly clear history of abrupt onset and offset in the attack with continuous rapid and regular beating during it; such histories can be obtained from observant patients. On the other hand a statement of gradual ending must be received cautiously since it is sometimes given by patients whose long paroxysms end in a series of quite short ones or in a declining number of extrasystolic interruptions. Another helpful point is the circumstance in which the attack occurs although paroxysms of tachycardia may be provoked by exercise.




FIG. 13.—Venous radial and electrocardiographic curves. The record is from a case of auricular flutter in which the auricle was beating at 250 and the ventricle at 175 per minute. The ventricular beats are recognized by the deflections *R* in the electrocardiogram, one of which precedes each radial beat. These deflections *R* spring from a zigzag line; the up-strokes of this line start at quite regular intervals and each complete phase corresponds to an auricular cycle; alternate auricular deflections are increased in size because *T* deflections fall with them. The venous curve consists mainly of very rapid waves but alternate *s* waves are distorted by *c* waves being superimposed upon them.

or by emotion usually they come unexpectedly and remain unexplained whereas in simple tachycardia definite acts usually bring them about or they recur regularly at given times such as meal times or on lying down at night. The general picture will often point distinctly to simple tachycardia as when the complaint comes from a nervous woman at the time of the menopause. But there will remain instances of doubt and in these the doubt will usually be resolved if the patient is seen in an actual attack.

A patient is noticed to have a rapid regular heart action. When ever the heart rate in a patient at rest is found at 140 or over the enquiry as to whether the tachycardia is of simple type or not

should be made. The normal heart rate may be as high as 140 in an adult lying down and fully atropinised, occasionally and very temporarily it rises to this rate under emotional stress, more frequently to this or even higher rates in high fever and exophthalmic goitre. But such rates are not long continued, they fall as rest continues. When, under resting conditions, the heart rate is fixed continuously above 140, essential abnormality of rhythm is usually responsible. When the rhythm of the rapidly beating heart is normal, the rate rises when the erect, and falls when the reclining, posture is assumed, it responds by rising still higher when the subject exercises, and falls when he rests, it rises higher in response to a whiff of amyl nitrite. None of these changes influences the rate in paroxysmal tachycardia, it remains at a fixed and high level in a great variety of circumstances. Very simple tests on patients, who are fit to submit to them, will thus distinguish the two states, but any test employed should be undertaken deliberately and repeatedly. The beginning or the ending of an attack if witnessed should always be decisive, for the change in the paroxysm is abrupt.

A patient is seen for the first time in an attack of rapid heart action, and the picture is that of acute failure with congestion. It may be the first attack, in which case the help provided by accounts of previous incidents will be lacking. The congestion being recognised, the condition is apt to be termed "acute dilatation" of the heart, if the attack has begun during exercise, or while lifting a weight it may be called "heart strain". Neither of these diagnoses is adequate, both are based on and lead to misunderstanding, they do not appreciate the significance of the rapid heart rate and its consequences. When in elderly subjects congestion and oedema of the lungs develop during the attack, dullness and crepitations at the bases may be mistaken for pneumonia. Abdominal pain, anginal or hepatic, with signs of collapse and running pulse in the attack, have been confused on many occasions with acute abdominal symptoms. Neither of these blunders will occur if the ventricular rate is accurately counted and the veins of the neck are carefully examined. Forceful beating in the veins of the neck with distension of the jugular bulb on the right side, may closely simulate innominate aneurysm on occasion.

Flutter—When the ventricle is responding regularly and at half the rate of auricular beating, the tests that differentiate flutter are the same as those that differentiate paroxysmal tachycardia from simple tachycardia, for the rates in flutter are also remarkably

constant in very varying circumstances. The differentiation between paroxysmal tachycardia and flutter is not always easy in bedside tests. In the former the rate is usually at or above, and in the latter at or below, 160 per minute. A long continued regular ventricular action of 130 to 160 per minute in middle aged or elderly subjects should always suggest flutter. If such tachycardia persists for two or more weeks and no change of rate occurs with rest or exercise, the diagnosis is practically certain. Frequent repetition of precisely the same high rates in the pulse over periods of days, weeks, or months is an important sign. On occasion, and usually as the result of unusual emotion or exercise the ventricular rate rises temporarily to the full auricular rate and syncope results. It is often possible in flutter to slow the rate of the ventricle for a few seconds by pressure on the carotid sinus (see page 79) the original rapid rate being quickly resumed; this test is often valuable. Digitalis given in adequate doses will always slow the ventricle and render its action irregular.

When the original ratio of response is higher, for example, when the auricle is beating say at 300 and the ventricle at 75 (4 to 1 response), flutter is more difficult to recognise. Occasionally the very rapid movements of the auricle are displayed in the neck. Sudden and precise doubling of heart rate (70 to 140 or 80 to 160) on slight exercise is significant.

When the response of the ventricle to a fluttering auricle is rapid and irregular (Fig. 14) the condition is usually confused clinically with fibrillation, unless exercise is found to make the pulse more rapid and at the same time regular. The error is unimportant, for the treatment of the two conditions is much the same.

The distinction between simple tachycardia, paroxysmal tachycardia, and auricular flutter can always be made with certainty electrocardiographically; this method when available should be used in cases of doubt.

PROGNOSIS

Simple Tachycardia—The prognosis of simple tachycardia obviously depends upon its cause, and requires no further comment here.

Paroxysmal Tachycardia—In considering the immediate prognosis of the severe attack several aspects require emphasis. The patient's symptoms are much influenced by the reaction of the nervous system, nervous subjects, especially women, awaken undue

anxiety The burden imposed on the heart depends upon the rate of its beating and upon the duration of the paroxysm the time that a paroxysm will last can often be judged from the length of previous seizures The manner in which the increased burden is borne depends upon the cardiac reserve As a prognostic sign the strength of the pulse has little value Anxiety is apt to increase when after several days congestion is progressively increasing respiration is becoming embarrassed and dropsy is appearing But even when circulatory failure has become advanced the paroxysm almost always ends favourably and the patient passes in a few minutes from a state of distress and of apparent danger to one of comfort and safety Confidence in recovery is almost always justified by the event namely, an abrupt cessation of the attack Attacks lasting ten days are very rare thus, as each day passes the end of the attack becomes more inevitable A fatal termination is a rare event except in those




FIG 14. Electrocardiogram of auricular flutter The auricular rate is 310 per minute the ventricle responds irregularly Time \equiv .1 sec

already gravely ill before the attack begins from cardiac failure pneumonia or other cause

The future outlook of patients who suffer from paroxysmal tachycardia is naturally very varied It is based firstly upon an estimate of the soundness of the heart and secondly upon the severity and frequency of the attacks The first factor is formed from the condition of the heart between the attacks The basis of prognosis is the same as in a similar case where there are no attacks it being remembered that the attacks themselves indicate a defect in the muscle and that very severe attacks temporarily threaten the patient's life The reaction of the heart to a paroxysm of tachycardia often itself provides valuable evidence of the heart's reserve When the heart is healthy a simple increase of rate (140 or more) causes its volume to decrease whereas a defective heart reacts by dilating In the first case the circulation is carried on effectively and in the second it becomes impaired

There is no actual evidence that paroxysms of tachycardia leave damage behind them though it would seem probable on general

grounds that frequent and severe attacks in subjects in whom there is little reserve must exercise an injurious effect. The possibility of unexpected death in a paroxysm, though remote, necessitates some, but not undue, caution in the general prognosis.

In general the most favourable, and incidentally the commonest, cases are those in which the heart is sound between the attacks and the paroxysms are infrequent, relatively short, and not severe. Such paroxysms rarely shorten life. Young subjects, and especially patients who are passing temporarily through a period of ill health, frequently lose their paroxysms altogether in later years. The least favourable cases are those in which there is evidence of cardiac damage between the attacks and in which the attacks themselves are frequent and severe.

Auricular Flutter —Instances are known in which the auricle has continued to beat without cessation at 280 per minute for a period of ten or more years, the ventricle following suit at half that rate, and the patient meanwhile showing little change. Prognosis follows much the same lines as it does in simple paroxysms and in fibrillation. The reaction of the heart to flutter depends upon its reserve, and upon the nature of the burden that the flutter imposes. It follows the lines laid down for fibrillation (page 85) more closely because both flutter and fibrillation are essentially persistent conditions, and because they arise in the same class of patient and present very similar ventricular rates that are in general below those of paroxysmal tachycardia. In making a prognosis in flutter it is important to allow for the results of treatment, to which flutter when properly handled is very amenable.

TREATMENT

Simple Tachycardia —The treatment of simple tachycardia obviously depends upon its cause. The rapidly beating organ is but a gauge telling of disturbed innervation, or of altered physical or chemical environment. The discovery of an enlarged thyroid, signs of intoxication, a focus of infection, or instability of the nervous system forms the guide to appropriate treatment. Bromides are most useful in treating attacks of simple palpitation, especially those associated with the menopause.

Paroxysmal Tachycardia —The treatment of cases of paroxysmal tachycardia resolves itself into remedies applied during the crises and general care of the patient during the course of the malady.

There is no unfailing remedy for the attack. A few patients are able to stop the seizures by assuming a particular posture, for example, by sitting and placing the head between the knees, or by inducing nausea. Deep pressure applied successively on the right or left side over the carotid artery below the angle of the jaw, the pressure being sustained for ten or twenty seconds, may terminate attacks. The termination is then due to a special reflex, set up from the specially sensitive wall of the carotid sinus, an expansion of the common carotid artery at and above its bifurcation, the nerve impulse travels to the medulla through the ninth, and returns through the tenth (the vagus) nerve. The procedure, though uncomfortable, is without danger, and is so often successful that it should always be tried. The application of an ice bag to the precordium, which usually affords relief, will sometimes stop the attack. More often than not all these remedies fail. Of drugs, the only one that is certainly known to have effect is quinidine, which may be given by the mouth in the doses prescribed for auricular fibrillation (page 90). Quinidine has also been used intravenously with occasional success, but is not without danger. It is given very slowly and well diluted as a single dose $1\frac{1}{2}$ to 3 grains, the patient's heart being most closely watched. In most cases treatment ends by being purely palliative. The patient rests in the posture that gives most ease, usually lying well supported on pillows, though sometimes preferring to sit or stand. The diet should be that given in failure with congestion (page 27) when attacks are severe or prolonged. The drinks should be iced. The ice bag, or an irritant, applied over the precordium or liver will relieve local pain. For severe pain morphia may be used, though this drug is not often needed. Hypnotics are used to induce sleep in long continued paroxysms. Grave congestion may be met by letting 8 to 12 oz. of blood flow from the veins, but the occasion for this does not often arise.

The general treatment of the patient is largely governed by the state of the heart between attacks. Enquiry may reveal exciting causes of the paroxysms. Scrupulous attention should be paid to general health, especially to oral and pharyngeal sepsis, the diet must be orderly, dyspeptic troubles remedied, and the use of tobacco moderated or banned. The prescription of 3 grains of quinidine, taken two or three times daily, will sometimes prevent attacks occurring or reduce their number, such dosage may be continued for months if necessary.

Auricular Flutter —The most important remedy is digitalis or an

allied drug. A rapid ventricular rate can always be reduced by giving full doses of digitalis, and the slower rate can be maintained so long as the drug is given. The treatment to this point is similar to that employed in fibrillation, under which it is described in detail (page 86).

It is often possible in cases of flutter to go further. Digitalis is given in stiff doses (1 drachm of tincture daily), and this dosage is maintained until the ventricular rate falls to 70 or less per minute. Continuing or increasing this dosage may now force the rate as low as 50 per minute. If this can be accomplished, without other toxic symptoms, it is not unusual for flutter to give place to fibrillation, the heart's action becoming grossly irregular though still very slow. Once fibrillation has appeared the drug should be stopped. If fibrillation persists it is treated as such. But it is not unusual for normal rhythm to return when the drug is omitted, and in that case no further treatment is required. This method of treating flutter by restoring normal rhythm is a little difficult to accomplish without frequent graphic records of the heart's action, but in practice it will be found that if flutter cases receive continuous and heavy doses of digitalis conspicuous improvement will follow, either as a result of simple ventricular slowing, or from the resumption of a normal rhythm. In either case signs of venous congestion, when previously shown disappear, and breathlessness and other discomforts are relieved.

Quinidine has also been employed in flutter, as in fibrillation, to restore normal rhythm, the method used being that described under fibrillation (page 90), the remedy, however, enjoys no great success.

CHAPTER X

AURICULAR FIBRILLATION (IRREGULAR TACHYCARDIA)

INTRODUCTORY

It is believed that auricular fibrillation, like flutter, is caused by a wave of contraction circulating continuously through the auricle (circus movement) but differing from that of flutter by circulating over a variable path and in a much faster rhythm. Like flutter it may be regarded as an intrinsic cardiac fault.

Auricular fibrillation is a condition in which the normal quick and strong systolic contraction of the auricular walls is lost. These remain in a position of diastole, and only fine tremulous movements, which have insufficient power to expel blood from the cavities, occur in them. The auricular systole, though mechanically ineffective, stirs the ventricles to rapid and wholly irregular action, the rate of response, when the conducting tissues are quite healthy, being as high as 180 or even 200 a minute. The usual rates of response are between 100 and 140 a minute. In special circumstances the rate may be lower.

Auricular fibrillation is one of the commonest forms of disorderly heart action in man. Essentially it is a persistent condition, once established it persists for the rest of life unless interrupted by treatment. A few cases are paroxysmal in type.

CLINICAL ASSOCIATIONS

In man auricular fibrillation has been noted from childhood to old age, but it is extremely rare before the age of twelve years. It is a frequent sequel to rheumatic fever or chorea, and is very often associated with mitral stenosis. Thus the rheumatic group is the largest, and has its heaviest age incidence in the twenties and thirties. A second group of cases of fibrillation display degenerative changes of heart and arteries, the incidence being chiefly in the fifties and sixties. Although fibrillation accompanies mitral stenosis

alised drug. A rapid ventricular rate can always be reduced by giving full doses of digitalis, and the slower rate can be maintained so long as the drug is given. The treatment to this point is similar to that employed in fibrillation, under which it is described in detail (page 86).

It is often possible in cases of flutter to go further. Digitalis is given in stiff doses (1 drachm of tincture daily) and this dosage is maintained until the ventricular rate falls to 70 or less per minute. Continuing or increasing this dosage may now force the rate as low as 50 per minute. If this can be accomplished, without other toxic symptoms, it is not unusual for flutter to give place to fibrillation, the heart's action becoming grossly irregular though still very slow. Once fibrillation has appeared the drug should be stopped. If fibrillation persists it is treated as such. But it is not unusual for normal rhythm to return when the drug is omitted, and in that case no further treatment is required. This method of treating flutter by restoring normal rhythm is a little difficult to accomplish without frequent graphic records of the heart's action, but in practice it will be found that if flutter cases receive continuous and heavy doses of digitalis, conspicuous improvement will follow, either as a result of simple ventricular slowing, or from the resumption of a normal rhythm, in either case signs of venous congestion when previously shown disappear, and breathlessness and other discomforts are relieved.

Quinidine has also been employed in flutter, as in fibrillation, to restore normal rhythm, the method used being that described under fibrillation (page 90). The remedy, however, enjoys no great success.

CHAPTER X

AURICULAR FIBRILLATION (IRREGULAR TACHYCARDIA)

INTRODUCTORY

It is believed that auricular fibrillation, like flutter, is caused by a wave of contraction circulating continuously through the auricle (circus movement) but differing from that of flutter by circulating over a variable path and in a much faster rhythm. Like flutter it may be regarded as an intrinsic cardiac fault.

Auricular fibrillation is a condition in which the normal quick and strong systolic contraction of the auricular walls is lost. These remain in a position of diastole and only fine tremulous movements, which have insufficient power to expel blood from the cavities, occur in them. The auricular systole though mechanically ineffective, stirs the ventricles to rapid and wholly irregular action, the rate of response, when the conducting tissues are quite healthy, being as high as 180 or even 200 a minute. The usual rates of response are between 100 and 140 a minute. In special circumstances the rate may be lower.

Auricular fibrillation is one of the commonest forms of disorderly heart action in man. Essentially it is a persistent condition, once established it persists for the rest of life unless interrupted by treatment. A few cases are paroxysmal in type.

CLINICAL ASSOCIATIONS

In man auricular fibrillation has been noted from childhood to old age, but it is extremely rare before the age of twelve years. It is a frequent sequel to rheumatic fever or chorea, and is very often associated with mitral stenosis. Thus the rheumatic group is the largest, and has its heaviest age incidence in the twenties and thirties. A second group of cases of fibrillation display degenerative changes of heart and arteries, the incidence being chiefly in the fifties and sixties. Although fibrillation accompanies mitral stenosis

frequently, in aortic disease it is rare. At least 70 per cent of all cases of failure with congestion display fibrillation. It is a common complication of hyperthyroidism (page 258)

PAROXYSMAL FIBRILLATION

Though fibrillation of the auricles may be spoken of as essentially a chronic malady, it occurs in some patients in a form of attacks lasting a few hours or a few days. Only in the very rarest circumstance is a paroxysm continued for longer than ten days. The cases are usually rheumatic heart disease or hyperthyroidism, in others no abnormalities are to be found between the attacks. Isolated attacks of auricular fibrillation sometimes supervene during the course of infection especially acute infection, such as severe tonsillitis, acute cholecystitis, appendicitis, pneumonia and infective endocarditis.

SYMPTOMATOLOGY

Fibrillation of the auricles gives rise to no symptoms directly, but by stirring the ventricle to rapid and irregular action, as it usually does, it results in "palpitation" or "fluttering" in the chest. The symptoms at the onset of auricular fibrillation are similar to those already described for paroxysmal tachycardia and are similarly modified. There may be no symptoms or these may be severe. In intelligent and observant patients will sometimes describe the beating of the heart as irregular. As a rule in persistent fibrillation, palpitation is not felt continuously but only when the ventricular rate rises high with effort or under emotion.

CARDIAC REACTIONS

When the auricles fibrillate the heart works at a disadvantage. In the first place the auricles are virtually paralysed, this, however, does not seem greatly to influence the general circulation. In the second place the ventricle beats irregularly and more rapidly. Owing to the rapid action the muscle enjoys less rest and it expends more energy (see page 23) the irregularity exaggerates these troubles because many of the beats are abortive in that they fail to raise the semilunar valves. A relatively healthy ventricle can tolerate the increased burden, a damaged ventricle does not, it dilates. It should be clear from what has been said, here and previously, that the changes following the onset of auricular fibrillation,

whether this is the beginning of the chronic condition or whether it is to be a paroxysm, are in every respect almost identical with those described under paroxysmal tachycardia (page 72), and it is unnecessary to repeat them. There are the same variations in the intensity of the symptoms, according to the rate at which the ventricle beats and according to the amount of the heart's reserve. For a given rate of ventricular beating the disturbance is more

usually a smaller capacity of the heart for work. Auricular fibrillation, when it comes, sometimes produces no increase of breathlessness on exercise, or only a slight limitation of respiratory reserve, but these cases are rare. It is usual for the patient to develop persistent breathlessness, or a persistent increase of breathlessness, or shortly to be plunged into a state of failure with congestion, which will continue, if the state is unmodified by treatment, until death supervenes. In cases that already show congestion before fibrillation comes, it will often immediately endanger life. In elderly subjects the rate of ventricular beating is not usually very high, and so the change of mechanism more often passes unnoticed or without much disturbance.

Cases of auricular fibrillation are relatively exempt, though not entirely so, from anginal pain.

RECOGNITION OF AURICULAR FIBRILLATION

Auricular fibrillation is responsible for two series of signs, the one is due to virtual paralysis of the auricle, the other to rapid and irregular action of the ventricle. The auricular signs comprising loss of the auricular venous wave and corresponding electrocardiographic deflection, and the appearance of small waves derived from the continuously fibrillating muscle of the auricle, are well displayed in graphic records (Fig. 15), by means of which auricular fibrillation is at once identifiable. The only clinical signs that are produced directly from the auricle concern the effect of fibrillation upon the heart sounds in mitral stenosis, under which valve disease they are described (page 134). The diagnosis of auricular fibrillation without the aid of graphic records depends almost always upon study of the ventricular action. Characteristically the pulse is turbulent, or grossly irregular, it consists of a medley of beats, varying in force

and most unevenly spaced. It should be noted especially that phases of irregularity are never accurately repeated and that when small groups of specially rapid beats occur these do not necessarily end in an unusually long pause. Numbers of the beats fail to reach the wrist but this is the habit of weak beats in general as for example extrasystoles. It is the rule for rapid beats to be weak and, as rapid beats are numerous in fibrillation many naturally fail to yield arterial pulsation. The loss of beats in transit to the artery is unimportant to diagnosis but makes it necessary that counts of ventricular rate should be taken from the heart itself.

One of the most reliable clinical signs is based on the association of irregular and rapid action of the ventricle. If the heart is beating




FIG. 15.—Venous radial and electrocardiographic curves taken simultaneously from a case of auricular fibrillation. The arterial pulse is completely irregular, each beat being accompanied by waves

irregularly and the ventricular rate is 120 or over, fibrillation is usually present. The higher the rate in these circumstances the more definite becomes the diagnosis. Sinus arrhythmia disappears when the rate surpasses 100, extrasystoles almost always when it surpasses 120, and the irregularity of auricular flutter almost always at 130 or 140. Irregularity at higher ventricular rates is practically unknown apart from fibrillation. As the rate is lower and it is often below 120 in the resting subject, so the sign becomes less reliable but reliability may be obtained by raising the rate artificially by exercise. Briefly, the association of irregularity with a pulse of 120 or over has the same significance whether this rate is spontaneous or induced. There are, of course, patients in whom acts of exercise

that are at all strenuous are undesirable, but in auricular fibrillation the rate lifts far more easily than it does when normal rhythm prevails. Amyl nitrite may also be used on occasion to raise the rate in tests.

The association of irregular heart action with venous congestion is in itself highly suggestive but has not the same value as the signs described. Auricular fibrillation is the only disorder that gives persistent irregularity of the heart's action, other irregularities disappear occasionally, so that from hour to hour or day to day quite regular action is noted.

In cases where the rate of ventricular response is low (below 80 or 90 per minute) the diagnosis is more difficult. These are usually patients under the influence of drugs such as digitalis, in such the presence of fibrillation will become more obvious as the effect of the drug lessens. There are however, cases in which the rate tends to be persistently low. Very close attention to the arrangement of the pulse beats is then required, and the relation of the rapid beats to deep inspiration should always be investigated to distinguish fibrillation from sinus arrhythmia. It is consoling to realise that although diagnosis becomes increasingly difficult as the ventricular

first time

PROGNOSIS

The prognosis of the attack in paroxysmal fibrillation that is untreated is similar to that of paroxysmal tachycardia but attacks of fibrillation can be brought to an end with greater certainty by giving quinidine by the mouth. If fibrillation has started and continued for more than ten or fifteen days it should be regarded, not as a paroxysm, but as the chronic condition. Untreated this lasts as long as life, the normal rhythm can be restored by quinidine in a number of patients, but in only a few of these does it persist. Normal rhythm is more likely to persist when the fibrillation was originally associated with hyperthyroidism and this has been treated (page 260).

The presence of persistent auricular fibrillation influences the general outlook in one of several ways. No patient who displays it can safely be regarded as possessing a sound muscle. It may constitute the only sign that the muscle of the heart is affected and

should always be so interpreted. Auricular fibrillation loads the already defective muscle with an extra burden in considering this extra burden rate of ventricular response is of chief importance. High rate is not long supported by damaged ventricular muscle but in this connection response to treatment is to be taken into account. It is not the rate naturally prevailing but the rate prevailing under such treatment as the patient can obtain that matters. A considerable reduction of the burden can be anticipated if adequate doses of digitalis are maintained for this drug influences the ventricular rate in a remarkable fashion. Nevertheless the onset of fibrillation in any patient is a serious matter it is a particularly grave event in those already suffering from cardiac symptoms at the least it means continuous medication with the heart rate often ranging much above its previous average. In most cases it heralds or actually brings venous congestion few survive the event for more than five or ten years. There are however rare but well authenticated instances of patients surviving for longer periods.

TREATMENT

Paroxysmal auricular fibrillation is treated upon the lines laid down for paroxysmal tachycardia (page 78). Digitalis or its allies tends to prolong fibrillation and should not be given unless there is extreme urgency dosage should then be arranged to yield a speedy reversion. After the tenth day if fibrillation still persists it may be regarded as permanent and treated accordingly. In the paroxysm a course of quinidine should always be given (see page 90) it is often successful. Doses of 5 grains twice a day if continued tend to ward off further attacks. From the standpoint of treatment the possibility of hyperthyroidism should be held in mind in all paroxysmal cases.

Speaking of chronic auricular fibrillation there is probably no type of cardiac disease that benefits so greatly from active interference than the well managed case of this disease. It is treated in one of two chief ways by digitalis or an allied drug or by quinidine.

DIGITALIS THERAPY

It is to their striking effects in cases of auricular fibrillation that drugs of the digitalis group almost exclusively owe their well founded reputation. In fibrillation it is usual for the ventricular rate to be increased digitalis can be made to control this rate as it can also in auricular flutter by decreasing the power of the auriculo ventricu-

lar bundle to conduct impulses from auricle to ventricle. By slowing the ventricle it brings the heart its much needed rest from an unusual burden of work. There are few cases in which the desired slowing cannot be obtained and control of rate maintained by continued administration. The higher the original rate of beating, the more certain is the response to treatment. The drug may be given beneficially whenever the heart rate is found higher than 90 per minute while the patient is at rest. In treating cases of auricular fibrillation the guide is ventricular rate, as this is counted at the cardiac impulse and not at the wrist.

Patients treated for the first time should go to bed and remain there until the reaction to the drug has been investigated fully, or for so long as other circumstances dictate. As a routine, tincture of digitalis should be used and, if heavy initial doses are to be employed, the tincture should be one that has been standardised biologically (U S A tincture) or by clinical experience. In the latter case the stock of tincture drawn upon should be a large one and it should be known when the stock is replenished. A usual and safe dosage for an adult is 60 minims of the tincture (U S A or B P) daily by mouth. This may be given as a single dose or divided into two or three doses. It is continued until a reaction in the form of the desired slowing of the heart rate, or until nausea or diarrhoea appear in response to the drug.

It may be said that in the average a patient will use up or excrete 20 minims of tincture daily, and that a reaction is not usually obtained until 5 drachms have been accumulated in the body so that the reaction is not usually seen before seven days. Many patients require more and a few require less. All must be watched closely. As a rule ventricular slowing is the first clinical sign of the drug's action. The object is to obtain a reaction including a fall of rate to 70 or 60 per minute. If the remedy is maintained too long, the rate may fall to 50 or even 40 and coupling of the beats may occur. Neither digitalis coupling (Fig 16) nor profound slowing should be allowed to continue, both are evidence of overdosage and call for immediate withdrawal of the digitalis. Many patients at this stage seem to be in a most satisfactory state of convalescence, they are actually in danger. The patient should be kept at complete rest with strict avoidance of emotional excitement until the heart quickens again. I have seen more than one sudden and avoidable death result from neglect of the signs named. In connection with slowing it is here to be observed that an allowance is to be made

for the effects of fever upon the heart rate, the rate of the heart in this should not be forced lower than 80 or 90 per minute. Occasionally coupling at high rates interferes with adequate treatment, in such digitalis and quinidine may sometimes be combined usefully.

In urgent cases, a more rapid reaction can be obtained by giving an initial massive dose of $1\frac{1}{2}$ to 2 drachms (to an adult of 100 to 140 lbs weight) under which the heart rate will usually begin to fall within eight hours. If there is little slowing, the dose may be repeated next day and smaller doses given subsequently. This method offers the prospect of rapid improvement, but is not to be used except upon patients under very close observation. It particularly suits cases displaying very rapid action of the ventricle (150 to 200); it is safe in the hands of those who thoroughly appreciate the behaviour of auricular fibrillation and have experience of its reaction to digitalis.





FIG 15—The radial pulse curves from a case of fibrillation of the auricle exhibiting coupled beats arising from overdosage with digitalis. The intervals between the large and small beat of each pair are uniform but the long intervals which follow each couple of beats are very irregular in length.

Another plan used in cases of the same type and with similar reservations is to administer the newly discovered and potent crystalline glucoside digoxin (Burroughs Wellcome). Either a dose of 0.5 to 0.75 milligram is given in 10 c.c. of saline, slowly and intravenously, in which case great care must be used that none of the drug escapes outside the vein, or a dose of 1.0 to 1.5 milligram is given by mouth. The first begins to slow the heart rate almost at once, and appreciably within two hours; the second takes obvious effect within four or six hours. Either may be followed by regular doses of digitalis at the rate of 1 drachm of tincture a day, starting at the 6th hour from the dose of digoxin.

There are patients in whom adequate slowing cannot be obtained by digitalis before gastrointestinal symptoms are induced by it. Vomiting due to congestion of the stomach must be distinguished, because it often disappears under continued digitalis treatment, which clears away congestion; vomiting is rarely produced by digitalis before a drachm of tincture has been taken. Digitalis may be continued for a day or two advantageously in cases in which it is causing nausea and occasional vomiting despite these symptoms,

for the effects of fever upon the heart rate, the rate of the heart in this should not be forced lower than 80 or 90 per minute. Occasionally coupling at high rates interferes with adequate treatment, in such digitalis and quinidine may sometimes be combined usefully.

In urgent cases a more rapid reaction can be obtained by giving an initial massive dose of $1\frac{1}{2}$ to 2 drachms (to an adult of 100 to 140 lbs weight) under which the heart rate will usually begin to fall within eight hours. If there is little slowing, the dose may be repeated next day and smaller doses given subsequently. This method offers the prospect of rapid improvement, but is not to be used except upon patients under very close observation. It particularly suits cases displaying very rapid action of the ventricle (150 to 200); it is safe in the hands of those who thoroughly appreciate the behaviour of auricular fibrillation and have experience of its reaction to digitalis.



110 10 The tidal pulse curves from a case of fibrillation of the auricle exhibiting coupled beats arising from a regular rate. The intervals between the large and small beat of each pair is uniform but the long intervals which follow each couple of beats, are very irregular in length.

Another plan used in cases of the same type and with similar reservations is to administer the newly discovered and potent crystalline glucoside digoxin (Burroughs Wellcome). Either a dose of 0.5 to 0.75 milligram is given in 10 c.c. of saline, slowly and intravenously, in which case great care must be used that none of the drug escapes outside the vein, or a dose of 1.0 to 1.5 milligram is given by mouth. The first begins to slow the heart rate almost at once, and appreciably within two hours, the second takes obvious effect within four or six hours. Either may be followed by regular doses of digitalis at the rate of 1 drachm of tincture a day starting at the 6th hour from the dose of digoxin.

There are patients in whom adequate slowing cannot be obtained by digitalis before gastrointestinal symptoms are induced by it. Vomiting due to congestion of the stomach must be distinguished, because it often disappears under continued digitalis treatment, which clears away congestion, vomiting is rarely produced by digitalis before 5 drachms of tincture have been taken. Digitalis may be continued for a day or two advantageously in cases in which it is causing nausea and occasional vomiting despite these symptoms,

for often the desired low rate follows soon upon these symptoms, but dosage cannot long be maintained after they appear, and thus they occasionally prevent fully successful treatment. The nausea and vomiting can be relied upon to disappear soon after digitalis is omitted, both symptoms are largely due to effects on the central nervous system, but not exclusively so, they can be avoided sometimes by diluting the tincture and by giving it by mouth after meals only, or rectally.

Strophanthus is a much less certain remedy than digitalis and, owing to the difficult absorption of its active principles, must be given in doses which contain relatively large quantities of these if the drug is to be effective. It has the danger too of variable absorption, and is very apt to cause diarrhoea, headache, yellow vision, and sometimes more serious symptoms. Considering these undesirable characteristics and that it scarcely ever, if ever, succeeds where digitalis fails, I am now inclined to recommend its complete disuse. In view too of the recent isolation of digoxin in pure form, I recommend the disuse of the less reliable strophanthin.

The initial object of treatment is to reduce ventricular rate to about 70 or 60 per minute, the second object is to maintain the rate between about 70 to 90 beats per minute in circumstances of rest and light exercise. When the rate has fallen on the initial heavy dosage, the quantity of the drug should be decreased and the amount that will maintain proper heart rates, as the patient is allowed more and more liberty, ascertained by trial. Usually it will be found that the maintaining dose in patients out of bed is 30 minims of tincture of digitalis in the twenty four hours, sometimes it is 20 sometimes it is 40, doses of 15 minims or less are nearly always useless for adults. It is usual for the patient to continue such doses for the rest of life and to be seen every few weeks. Continuance of the drug is governed almost entirely by heart rate, enough is to be given to maintain rates that are little in excess of normal, cumulative effects do not occur provided the rate is not allowed to become excessively slow. Intelligent patients quickly learn to regulate their own doses by their sensations, taking the minimal quantity that frees them from breathlessness or palpitation, they can be allowed to do so provided that a maximal dosage is named, which may not be exceeded. Business men find it convenient to carry the small phials of Nativelle's granules. They are very potent, one or at most two of the white granules a day should suffice to maintain proper rates for those already under the influence of the drug.

CHAPTER XI

BRADYCARDIA SYNCOPE, AND SUDDEN DEATH

BRADYCARDIA

SIMPLE BRADYCARDIA

SIMPLE bradycardia is a term used to indicate a natural action of the heart at a decreased rate, the whole heart participates in the slow action and it is caused by decreased rhythmicity of the pace maker (Fig 17) The heart in such is usually healthy In the human subject this slowing of the heart occurs in a variety of circumstances Although the rate of about 70 per minute has long been accepted as the average for the normal subject at rest, lower rates (65, 60, 55, or even 50) are not infrequent in perfectly healthy people The resting heart rate is often slower during adolescence (Fig 18) and is frequently much below the average in athletes The rate becomes slower with advancing years It is often decreased in convalescence from acute illness It may fall sharply after brief exercise and may be retarded by exposure to cold Gross cerebral lesions, such as tumour, apoplexy and meningitis, are often associated with slow action of the heart likewise myxoedema and, so it is stated, forms of insanity such as melancholia In jaundice the pulse is usually slow, often notably so In general the pulse tends to fall when blood pressure is raised Slow action of the heart is a usual accompaniment of common fainting attacks (page 98), of vomiting, and of attacks of giddiness of middle ear origin, in all these it is vagal The slow heart in jaundice and the relatively slow beat in typhoid fever are due to influences other than those of innervation

HEART BLOCK

Heart block is an abnormal mechanism in which the ventricle fails to respond properly to the beats of the auricle It is responsible for regular slow action of the ventricle when the latter responds

to each second beat of the auricle (2:1 heart block), or when it fails to respond at all and the ventricle assumes an independent slow rhythm (complete heart block) (Fig 19)

Heart block is caused by reduced power of the auriculo-ventricular node or bundle, which normally convey impulses from auricle to ventricle, to conduct. It may be produced experimentally, (a) by direct interference with these structures, as by pressure or cooling, (b) by inducing the vagus, and especially the left nerve, to




FIG 17—A diagram of a human heart. The walls of inferior vena cava, right auricle and right ventricle have been partially removed to expose the septa. The cut surfaces are stippled in the diagram. The position of the sino-auricular node in which the heart-beat commences, is shown as are also the position of the auriculo-ventricular node and the course of the auriculo-ventricular bundle and its branches. The last-named structures convey the contraction wave from auricle to ventricle.

overact upon them, and (c) by changing the character of the blood that supplies them, for example, by asphyxiation or by the injection of such substances as digitalis, large doses of adrenaline, or diphtheria toxin.

In man heart block can happen at any age. It may be congenital. It often results from a lesion invading the auriculo-ventricular bundle. These lesions are mostly inflammatory, acute, subacute, or chronic, commonly the inflammation is rheumatic, sometimes it is

siphilitic in later life degenerative processes with or without calcareous deposits are more responsible. Rarely tumours starting in, or invading the bundle have been recorded. The lesions that produce heart block are not often confined to the bundle.

Minor grades of heart block such as are responsible for intermittence of the ventricle (page 65) are commonly associated with infections especially subacute rheumatism, diphtheria, influenza, and pneumonia. The minor grades are often brought about also by vagal influences as by pressure on the sensitive carotid sinus (page 79) vagal block is usually temporary and is relieved by atropine. The same grades of block often occur during heavy administration of such drugs as digitalis, strophanthus and squills, especially when these are administered to young subjects suffering from rheumatic heart disease.

The only special symptoms associated with heart block are due to profound slowing of the ventricle or to sudden cessation of its beat. The resultant syncopal attacks are not infrequent in heart block of high grade. They are described on page 100.

DIFFERENTIATION OF BRADYCARDIAS

Differentiation of various forms of slow action of the ventricle may usually be accomplished by bedside signs. The diagnosis can always be made absolute by means of graphic records. In simple bradycardia the ventricular rate at rest is rarely less than 50 per minute. In 2:1 heart block rates of 40 to 50 generally prevail. In simple bradycardia the ventricular rate rises gradually with exercise, excitement, and such drugs as atropine and amyl nitrite. In 2:1 heart block these influences frequently raise the ventricular rate abruptly to double, and subsequently it falls abruptly to its original point. 2:1 block is usually unstable, and abrupt changes of rate with or without intervening periods of irregular action are frequent. Thus in differentiating simple bradycardia from 2:1 block such simple tests as exercise and the inhalation of amyl nitrite are very valuable.

The diagnosis of complete heart block is suggested by regular ventricular action below 35 beats per minute, a constant rate of 30 per minute almost always indicates complete heart block, especially if this so if there is a history of syncope. In complete block the rate of the ventricle is usually uninfluenced by exercise, atropine, or amyl nitrite. The diagnosis can usually be made absolute by noticing the characteristic variations in the heart sounds, which occur in-

dependently of respiration. The breath being held, first and second sounds are heard with each ventricular systole but these sounds vary greatly in intensity or show inconstant reduplication from cycle to cycle. Sometimes muffled auricular sounds may be heard over the right auricle, or at the impulse, or in the epigastrium. The




FIG. 18.—A single-lead electrocardiogram showing the exceptionally slow heart rate of 31 per minute. The bradycardia is a simple one of unknown origin; each cardiac cycle being represented by normal auricular and ventricular deflections. Time in $\frac{1}{4}$ sec.

occurrence of numerous and regular waves in the jugular veins, some falling in the long intervals between ventricular systoles, is also diagnostic.




FIG. 19.—Venous, radial and electrocardiographic curves taken simultaneously from a case of complete heart block. The ventricle is beating at 47 and the auricle at 5 per minute. The relations of auricular and ventricular systoles are clear in the electrocardiogram; the ventricle being represented by deflections R, S, T and the auricle by summation P. The auricular and ventricular rhythms are each regular but independent. A radial pulsation and a venous wave c occur with each ventricular complex (R, S). A venous wave a and a minute radial a wave occur with each P wave. When in the venous a and c fall together a large r wave results. In the electrocardiogram auricular and ventricular deflections superimpose accurately when they fall together; thus the last cycle P falls upon R and S consequently R is faded and S is less deep.

PROGNOSIS AND TREATMENT

Simple Bradycardia is without direct prognostic or therapeutic significance.

Heart block occurring under the influence of drugs should not influence prognosis occurring otherwise even in its milder grades.

(dropped beats) it is to be regarded as a serious manifestation. It is to be read as usually indicating a diffuse affection of the muscle; it should not be dismissed as a mere affair of the bundle. Almost always the underlying process is widely diffused throughout the more silent areas of the myocardium and other evidences of a failing muscle are forthcoming. In rheumatic fever, pneumonia, influenza and other infectious diseases its occurrence is significant of a severe infection and it may be the only outstanding sign of cardiac damage. In most of such infections however the block is transient there being recovery from much or all of the original damage. Most patients who present persistent heart block die within a few years whether the block is of milder or severer grade and whether syncope occurs or not. The majority never have syncopal fits but die of failure with congestion. There are occasional cases in which heart block even of high grade continues for very many years; these are exceptional instances in which the driving muscle has suffered little and remains sound.

Strictly speaking there is no treatment for heart block. Treatment is directed to the cause or to the symptoms. Where partial heart block is of recent origin it is an index of active mischief such as an intoxication or an infection and the subject of it should be searched thoroughly for the cause. He should lie at rest in bed until all signs of block have gone or until it becomes clear that the block is permanent. The higher grades of heart block are usually persistent; the habits of these patients are governed according to the proved capacity of the heart to do work. Many such patients are able to pursue usual duties though they are unable or they are unwise to be very active. Where there is any evidence that the lesion is progressive rest and close attention are necessary. Syphilitic subjects should be treated thoroughly with mercurials or arsenical preparations (page 226).

The prognostic significance of the syncopal attacks of heart block and the special treatment of these seizures are given more conveniently under syncope (page 102).

SYNCOPE AND RELATED PHENOMENA

An attack of faintness or actual fainting is a frequent reason bringing patients for advice. It is a popular belief that such attacks are often manifestations of serious cardiac disease or precursors of apoplexy. Although there is some ground for these beliefs the great

majority of the attacks referred to have far less significance. It is a matter of much practical consequence that syncope and related phenomena should be understood thoroughly. The chief forms of the attack referable to the cardiovascular system will be described, and subsequently contrasted with some phenomena from which they must be differentiated.

By syncope we mean loss of consciousness caused by a deficient flow of blood from heart to brain. The defect may be vascular, the input of blood to the heart being diminished, or cardiac, the pump failing to do its work adequately. When it is vascular there is a preliminary fall of general blood pressure. A fall to 90 or 80 mm. Hg may be maintained without effect on consciousness. If the systolic pressure falls to such levels as 60 or 50 mm., the pulse becomes extremely feeble or imperceptible. Loss of consciousness occurs when systolic pressure falls to these levels or somewhat lower. Syncope resulting from a decreased return of blood to the heart whether due to haemorrhage or to vasodilatation, is associated with loss of pulse-beat.

POSTURAL FAINTNESS

If a hutch rabbit is held for a period by the ears its blood accumulates in the abdominal viscera, the heart becomes empty and the rabbit shortly loses consciousness. When the human subject passes from the supine to the erect posture there is ordinarily a small fall of blood pressure which is quickly and in large part compensated. In many people this phenomenon is exaggerated and leads to temporary giddiness or transient loss of consciousness on rising. The complaint comes usually from middle aged or elderly subjects, and the symptom is particularly frequent on rising from a chair after a full meal in an overheated room. On rising the vision of the subject becomes dim, he is momentarily unsteady, puts out a hand to support himself, or braces his muscles and so recovers. In some cases the effects are more profound and definitely alarm those who experience them. In rare instances the effect is not transient but continues and the subject is unable to maintain both the erect posture and consciousness. The phenomenon is due to an accumulation of blood mainly in the abdominal vessels, there is faulty distribution. A closely related event occurs after stooping to lace the boots, or at the end of a strenuous lifting movement. In both these acts blood tends to be driven out of the abdominal vessels by pressure or muscular action, and sudden resumption of the erect posture in the first, or relaxation in the second, instance leaves a

large potential reservoir for blood in the abdomen. This then fills, meanwhile preventing a natural return of blood to the heart.

VASOVAGAL ATTACKS

These are the commonest form of fainting attack, they occur in both sexes, especially, but not exclusively, in the young. A condition of poor health predisposes, thus they are more frequent after recent illness or in chronic infection, fatigue, a period of fasting, confinement to an overheated, overcrowded room are not unusual attendant circumstances. The attacks are not confined to people presenting no signs of disease, they also occur in cases of chronic heart disease and in fact form the usual type of fainting in aortic regurgitation. It is insufficiently realised, as this example illustrates, that transient disturbances yielding symptoms in relatively healthy folk also occur in cardiac patients, and it is necessary to be on guard not to interpret such symptoms as belonging to the major disease. The vasovagal attack is a probable cause of fainting in response to certain forms of pain, such as that experienced in coronary thrombosis in the manipulation of disused joints, and in gallstone and in renal colic. The attacks, however, are not confined to people who are unwell, but also occur in robust health.

Nervous agitation, emotional distress, are commonly provocative, for example an abrupt order or threat, an unexpected and important encounter, the receipt of unusual and stirring news, the witnessing of a distressing accident. The provocation may be adequate without arousing feelings either of dread or disgust, there is usually an element of surprise, as when the subject sees his own blood flow into a syringe, or a sphygmograph lever begin to move on his own wrist.

The central feature of the attack is syncope, loss of consciousness may occur without warning and result in a heavy fall and injury, more usually there are warning symptoms, and then the subject sits or lies down deliberately, or slips or falls more slowly, the warning is a feeling of instability or uncertainty, a dimming of vision or giddiness. Sense of rotation does not occur. The attacks occur rarely, if ever, while the subject is lying down, but almost always while he stands or sits. A peculiar sinking feeling in the abdomen, generally accompanied by nausea, sometimes by retching, and occasionally by a feeling that the bowels will be moved, are often early symptoms. Precordial distress or constriction do not occur, and real alarm is very rare.

From the beginning of the attack the blood pressure falls progressively and is almost always accompanied by a steep fall of heart rate, the pulse often becoming imperceptible and the rate falling to 50 or 40 (exceptionally to 30) beats per minute. Pallor, rapidly increasing in intensity, and ultimately extreme, accompanies these changes. The pallor is of the type associated with simple drainage of blood from the skin, being most in evidence in the face and much less so in dependent parts such as the hands. As cerebral anaemia increases the subject is seen to become unsteady, confused in speech, restless, and yawning. There is decreasing response to questions and physical stimuli, loss of consciousness usually follows with flaccidity, some dilatation of the pupils, and lost conjunctival reflexes. Slight clonic movements of face, upper limbs, and trunk may occur, respirations become slow, deep, and sighing. Sweating, first of face then of body, is usual and often extremely free.

The attack is almost always brief, lasting from two to ten minutes, rarely half an hour or more. Recovery is slower than the onset, leaving the patient limp and inert, and the skin pale and wet. Weakness, tremulousness, and sometimes severe headache may last for hours.

The slowing of the heart has been proved to be due to increased vagal tone, the slowing but not the loss of consciousness is relieved by atropine. The low blood pressure is an independent effect, presumably due to dilatation of the vessels in the splanchnic area, it is the vasodilatation that is responsible for loss of consciousness. The double term *vasovagal* aptly describes two effects, proved to be dominant features of the attacks. It is believed that the attacks are caused by reflex disturbance of a central nervous mechanism, which is normally set in unexaggerated motion by stimulation of either the carotid sinus (page 79) or the depressor nerve.

CARDIAC SYNCOPE

Postural faintness and vasovagal attacks are common. Those instances of syncope now to be described are rare and consist of instances in which the heart fails to transfer incoming blood from the venous to the arterial side. This is cardiac syncope properly so called, and results when the ventricles cease beating or beat very slowly, and when the ventricles beat very rapidly or fibrillate. It has no special association with any particular valve lesion.

Ventricular Arrest—Simple slowing of the heart fails to produce a material lowering of mean blood pressure, it raises systolic and

lowers diastolic pressure the heart accommodates itself to the increased quantity of the blood returning to it in diastole, throwing it out at each systole. Great slowing of the ventricles alone gives rise to symptoms. It is found in man that if the rate becomes as slow as 20 to 8 beats per minute, unconsciousness results. These rates are equivalent to cycles lasting about three to seven seconds. It has been shown that cessation of the heart beat for a single period of five to ten seconds is sufficient to produce unconsciousness. Arrest lasting from fifteen to twenty seconds causes obvious distension of the veins, occasionally cyanosis, and deep breathing, a little twitching of the face and upper limbs is added in pauses of longer duration.

There are very rare cases in which the whole heart stops beating for a few seconds; these are probably often of vagal origin. In others, block between auricle and ventricle comes suddenly, normal response




FIG. 20—Auricular and radial curves from a patient who exhibited many brief attacks of syncope. The record shows the syncope attacks to be due to temporary heart-block; the auricle beats throughout but the ventricle stops first for five and then for three seconds; the ventricular slowing resulted in loss of consciousness. At the end of the attack normal action is at once resumed. Time 1 sec.

of the ventricle ending abruptly in a pause (Fig. 20). Neither these varieties of ventricular arrest nor transient ventricular fibrillation (page 103) which is known occasionally to induce syncopal attacks, is of sufficient frequency to be of much practical importance.

When arrest of the ventricular beating is responsible for repeated syncopal attacks, this arrest almost always occurs in cases of chronic heart block. In patients who suffer from high grades of partial heart block or in whom there is complete block, attacks of giddiness, of momentary loss of consciousness, of longer unconsciousness with congestion and epilepsy are common manifestations. In these fits the ventricular action is extremely slow or ceases for shorter or longer periods; the auricles continue to beat, as is shown by the waves that appear in the veins of the neck. The severity of the attack is governed by the degree of slowing or by the length of time over which the ventricular beats lapse. Death results when the ventricular arrest lasts from one to two minutes. In some instances of heart-block the attacks may be absent or infrequent, occurring at intervals of months or years, in others they are more frequent, coming weekly

or daily, or occurring in quick succession over periods of several

biting in the fits are extremely rare. Sweating is not a feature of the attacks, though occasionally present in them and in the stage of recovery.

These cases, in which persistent slow action of the ventricle is accompanied by syncopal attacks, are the cases to which the term "Adams Stokes syndrome" is peculiarly applicable.

Rapid Action of the Ventricles—Syncope may be caused by very rapid action of the ventricles. It is usually necessary for the rate to rise to 250 or more. At such rates diastole is so short and the beats are so weak that little or no blood is forced into the arteries and the blood pressure falls profoundly. The attacks to which patients suffer from auricular flutter are prone to be of this nature (page 76).

Fibrillation of the ventricles (page 103) is to be associated with sudden death rather than with syncope from which there is recovery. Very rare cases have been described of brief loss of consciousness arising from a disorder which, if not ventricular fibrillation, is very closely akin to it.

DIFFERENTIAL DIAGNOSIS OF FAINTING

It is often necessary to differentiate between fainting that arises from the heart or general vascular system—syncope as it is termed—and loss of consciousness that is primarily due to a cerebral condition such as epilepsy. Syncope is always associated with loss of pulse beats either because the pulse becomes too weak to be perceptible, or because the ventricle fails to beat. Thus loss of consciousness, in which the pulse rate departs little from normal in rate or strength, cannot be the result of diminished blood flow to the head and neck; it is a form of fainting that is cerebral in origin. For similar reasons pallor is constant and conspicuous in syncope; it is not a feature of cerebral faints. Clonic movements occur in syncope occasionally, but are not violent so tongue biting and evacuation of the bladder which distinguish many attacks of epilepsy, do not occur in syncope. A sense of rotation never occurs in syncope, a fact that is very helpful in differentiating it from auditory vertigo.

The peculiarity of vasovagal attacks, that they rarely occur in a

patient who is lying down, is often crucial in distinguishing these both from epilepsy and from cardiac syncope. Profuse sweating favours and is almost necessary to a diagnosis of vasovagal seizure, but does not exclude cardiac syncope, in which, however, sweating is rare. The presence of heart block between these stamps these as due to ventricular disease.

perpetuated in the aged or elderly. Ascribed to cerebral arterial disease, and regarded as attacks premonitory to an aneurysm, cerebral arterial disease usually presents itself in a variety of ways, which may consist of a gradual loss of consciousness.

The attack always be recognised by suitable enquiry. The chief points differentiating the attacks of cerebral arterial disease from vasovagal attacks are absence of profound pallor or heavy sweating and unaltered pulse in the former. In attacks due to cerebral arterial disease, nausea may be present. It is frequent the associated symptoms are not observed. The attacks are usually of a few minutes' duration.

PROGNOSIS AND TREATMENT

Postural Giddiness is without any serious significance, it is occasional and not as a rule persistent. It may be relieved or abolished by the use of a broad and firm abdominal support, which should be advised when attacks are frequent or sufficiently severe to threaten or induce loss of consciousness. Otherwise treatment consists in attention to general health, to reduction of obesity and to measures (exercise and massage) that increase the tone of the abdominal muscles.

Vasovagal Attacks are not serious. They are occasional events experienced once, or a few times, and often in special circumstances that can be avoided. In the attack the patient should be placed supine, preferably with the head low, and, as in syncope from whatever cause, the clothing around the neck and chest should be loosened. In almost all cases these simple measures suffice and recovery occurs within a few minutes. In more prolonged attacks, atropine will relieve pulse slowing and stop sweating. Spirituous liquors should not be poured into the mouth of any unconscious subject, but may comfort or revive a faint person. Subcutaneous adrenaline (5 minims of 0.1%) is also recommended in prolonged attacks.

In the general treatment of these cases no special remedies are used, the basis of treatment is close attention to and observance of

general rules of health, especially recommending open air exercise when long hours of sedentary work is the habit

Ventricular Arrest—Fits that are due to ventricular arrest are usually so brief that no remedies can be employed In single long attacks it is clearly useless to give cardiac stimulants by mouth or subcutaneously, for the remedy will not reach the heart The injection of adrenaline (5 to 10 minims) directly into the heart

one or more doses of adrenaline (2 5 minims) given intravenously, the veins of the arm being massaged towards the heart after the injection to expedite the passage of the drug to the heart Ephedrine given three or more times daily in doses of $\frac{1}{4}$ to $\frac{1}{2}$ grain has also been used successfully Barium chloride in doses of $\frac{1}{2}$ grain daily by mouth is stated to have beneficial effects in some cases Strychnine, digitalis preparations, and amyl nitrite are useless The general treatment, when attacks are occurring hourly or daily, should confine the patient to bed over the period of disturbance and for some weeks afterwards, gastrointestinal or other disturbances that predispose to attacks should be enquired for and corrected when possible All those who suffer from syncope should be warned that it is dangerous to run the risk of attacks in situations in which they are exposed to accident Not a few lose their lives by falling heavily or in traffic Another and more frequent manner of death is failure of the ventricle to renew its beat this ending is not infrequent in patients who experience many severe attacks, or enter a state of prolonged unconsciousness The lives of those who have these seizures are very unstable

SUDDEN DEATH VENTRICULAR FIBRILLATION

In cardiac cases, anginal or otherwise, that die suddenly, coronary arterial disease is the commonest lesion to be found In many of these cases and whether there is fatty change in the muscle of the heart or not (see page 250), death is probably due to fibrillation of the ventricles Where actual thrombosis of a coronary artery is discovered, death is almost certainly produced in this way in most cases, as experiments on animals have clearly indicated, occasionally rupture of the necrosed ventricular wall is responsible, especially in the aged

Fibrillation of the ventricles is a condition in which co ordinate systole is suspended and the muscle of the ventricle exhibits a minute but ineffectual twitching movement. Recovery from this state is a rare event. Unhealthy or poisoned heart muscle is much more prone to fibrillate than normal muscle (for death under chloroform see page 274). Sudden death in cases in which the auricle is already fibrillating is usually attributable to this disorder. It is the cause of death when this occurs abruptly under heavy doses of digitalis when ventricular action is slow and possibly coupled (Fig 16 page 88). The catastrophe occurs although convalescence seems advanced and may be provoked by sudden movement or emotion. Perhaps the patient sits up in bed to fall back suddenly the pulse has gone there are a few gasps cyanosis rapidly develops and the patient is still. No lesion to account for death is found subsequently.

Vagal overaction has also been regarded as possibly producing sudden death but death occurring in this way has not been proved and it is doubtful if permanent arrest of the heart beat can be brought about through the vagus nerve. Ventricular arrest occurs in heart block and is responsible for sudden death in these cases.

A ruptured aorta a large thrombus plugging aorta pulmonary artery or narrowed mitral orifice are occasionally responsible for sudden death in heart disease.

In hypertension although sudden death is most often due to apoplexy coronary disease is not infrequently responsible.

CHAPTER XII

CARDIAC ENLARGEMENT

THE size of the heart is variable in healthy people, the weight of its muscle is related to that of the somatic muscle. The taller and stronger the man, the bigger and heavier is the heart. The heart is heavier in those who engage in heavy work or exercise than in those who live sedentary lives. These are physiological variations.

CAUSES OF PATHOLOGICAL DILATATION AND HYPERTROPHY

Pathological enlargement of the heart may be due to dilatation of its cavities or to hypertrophy of its muscle.

CAUSE OF DILATATION

Pathological dilatation is caused by inability of a well supplied heart properly to discharge its content. Putting on one side exceptional cases of extreme obstruction to the outlet, inability to discharge is due to inefficient beating. Inefficient ventricular beating may result from very excessive rate, as in flutter or fibrillation of the auricle. It may be due to weakness of the muscle as a consequence of its poisoning, its inadequate nutrition, or its structural disease. Examples of poisoning are to be found in the acutely dilated heart of such diseases as pneumonia, subacute rheumatism, diphtheria, and asphyxial conditions, of inadequate nutrition in coronary arterial disease, severe anaemias, and probably in instances of low mean blood pressure in the aorta to which rate of flow through the coronary arteries is sensitive. The pressure before which the heart dilates is that within it while it fills; it is a low venous and not an arterial pressure. Thus applies to general dilatation, we must exclude distension of the cardiac wall where it is locally fibrosed—thus in the case of the ventricle a fibrosed area may gradually distend to form an aneurysm of the wall, it distends under

systolic ventricular pressure—and we must exclude the events in rupture of the aortic valve discussed on page 152

CAUSE OF HYPERTROPHY

The recognised cause of hypertrophy is increase in the work of the heart sources of increased work sufficient to produce hypertrophy are persistent high blood pressure defects of the valves or septa and in certain instances pericardial adhesions

Increased load falls on different parts of the heart unequally Thus mitral stenosis imposes its burden upon the left auricle and right ventricle and these are the chambers that show corresponding hypertrophy most conspicuously Aortic regurgitation and aortic stenosis burden the left ventricle and cause most conspicuous hypertrophy in this chamber High systemic blood pressure is a most potent cause perhaps the most potent of left ventricular hypertrophy The distribution of these hypertrophies however is not so simple as it is often thought to be Thus in half the cases of mitral stenosis the left ventricle is increased quite as much as the right and in half the cases of aortic disease the right ventricle is increased quite as much as or more than is the left In both diseases it is the rule for the two ventricles to participate in the enlargement although one may do so more than the other Simple adherence of the visceral and parietal pericardium produces little or no enlargement of the heart the burden of the heart may be increased however when the heart's action is impeded by adhesions between it and surrounding structures There are many cases of hypertrophy and of great hypertrophy of the heart in which during life and after death no source of increased work can be discovered The degree of enlargement found in many hearts in which there is a valve defect such as aortic regurgitation or mitral stenosis is sometimes out of all proportion to the apparent increase of burden There is still much that remains to be explained it is clear that there must be hidden sources of increased work or the conclusion that increased work is the cause of hypertrophy needs revision

SYMPTOMS

There are no known symptoms of hypertrophy of the heart Such symptoms as might be ascribed to dilatation have been described already under failure with congestion

MEASURING THE HEART'S SIZE IN LIFE

The most accurate method of estimating the size and position of the heart during life is by projecting its outline with X rays. When the chest is heavily covered with fat or the lungs are emphysematous, it is often the only method that has any value. If the X ray picture is taken with the tube close behind the chest and the plate resting on the sternum, the outline is inaccurate. A light held in the hand causes an enlarged shadow of the subject's body to be projected upon a neighbouring wall, the enlargement depends upon the distance of the body from the candle on the one hand and from the wall on the other. If the body is near the candle and far from the wall, the distortion will be great. In X ray work similar conditions prevail. To obtain a radiogram of the heart with sufficient accuracy, the plate should be against the front of the chest and the tube 2 metres from the back. When the tube is within a short distance of the chest wall, error is large. The distortion is due to the divergence of the rays which emanate from the small cathode, and separate from each other more and more widely as they pass the margins of the heart.

The Orthodiagraph overcomes this error of measurement. The X ray tube is encased except at one point and moves easily upon a special arm, only one small beam emerges from the protective case and passes through the chest, and this beam is carried around the heart's margin, so that its paths through the chest are always quite parallel to each other. A complete and undistorted outline of the heart is recorded by a special device as the beam moves. In this way, each part of the outline being recorded in a similar phase of the respiration and in diastole, an outline of the heart is produced that is as accurate as any that can be obtained. As a measure however, it is still imperfect, for the record measures the heart in a single plane only. To obtain a volumetric measure of the heart it would be necessary to record the silhouette in a number of planes. The coronal record is, however, in general a sufficient gauge of size, and this has been used to check bedside methods of estimating the size of the organ. Statements of the values of methods now to be discussed are based chiefly on these comparisons.

The Maximal Impulse — In the healthy chest of an adult standing erect the cardiac impulse is usually confined to a small area and is found in the 5th intercostal space 4 inches (8 to 10 cms) from the middle line. $3\frac{1}{2}$ inches, the value usually given, is insufficiently

elastic

in large

nipple l - -

The impulse does not move much when the subject lies down, occasionally there is movement, and it is then generally towards the sternum. The impulse is not only visible, it is palpable. A finger placed upon it clearly feels systolic movement of the underlying heart muscle. It is not an almost impalpable flicker but a definite and more or less sustained thrust. The circumscribed area




FIG. 21. Orthodiagram 21 is and all other orthodiagrams are exactly one sixth natural size. Tail of 16 years weight 100 lb. Maximal impulse (M) standing 6th space to right 13½ cm from mid line. Percussion only (P) 19 cm from mid line. It is also marked in 2½ cm on lying down. This and subsequent orthodiagrams were taken in the erect posture. The right border is composed of right auricle (RA), ascending aorta (AA) and superior vena cava (SVC). The left border is composed of ventricle and basal vessels. The knob (K) representing the transverse aorta as it passes backwards.

over which this is felt is in most men about 1 inch in diameter, and its outermost limit is the best clinical guide that we have to the heart's size provided that this organ is not displaced. It usually corresponds accurately with the left limit as displayed in the orthodiagram, in hearts that are not enlarged or not greatly enlarged. When the heart is considerably enlarged the impulse is often in the axilla. The movement is now to the side, and is not a forward thrust, for the heart lies to the right of and not behind the part of the chest that displays the impulse. The anterior surface of the heart is no longer responsible for the movement of the chest wall, but, as the heart enlarges and the impulse travels back into the axilla, its left surface becomes more and more responsible

for the impulse. The impulse can no longer be related to the X-ray silhouette although it still remains an index of the heart's size.

A well defined maximal impulse, the outer border of which is situated 4½ inches or more to the left of the mid sternal line, or quite clear of the nipple line in an adult male, is to be taken as a definite sign of the heart's enlargement in the absence of cardiac displacement. The sign is all the more emphatic if it is discovered in the 6th interspace, it is less emphatic if found in the 4th.

elastic The nipple line, also used as a guide is sound inasmuch that in large men the nipple is more distant from the sternum but the nipple has an inconstant relation to the outline of the chest wall The impulse does not move much when the subject lies down occasionally there is movement, and it is then generally towards the sternum The impulse is not only visible, it is palpable A finger placed upon it clearly feels systolic movement of the underlying heart muscle it is not an almost impalpable flicker but a definite and more or less sustained thrust The circumscribed area




FIG. 1.—Normal Heart. This and all further orthodiagrams are exactly one-sixth natural size. Last 10 years well marked maximal impulse at standing the space between the ribs was from 10 to 12 cm. The heart is 1/3 and 1/4 of the normal size. The impulse is visible on the lower part of the chest. The right border is composed of right auricle (P.A.) and right ventricle (V.C.). The left border is composed of ventricle and basal vessels the knob (K) representing the transverse aorta as it passes backwards.

over which this is felt is in most men about 1 inch in diameter, and its outermost limit is the best clinical guide that we have to the heart's size, provided that this organ is not displaced. It usually corresponds accurately with the left limit as displayed in the orthodiagram in hearts that are not enlarged or not greatly enlarged. When the heart is considerably enlarged the impulse is often in the axilla. The movement is now to the side, and is not a forward thrust for the heart lies to the right of and not behind the part of the chest that displays the impulse. The anterior surface of the heart is no longer responsible for the movement of the chest wall, but, as the heart enlarges and the impulse travels back into the axilla, its left surface becomes more and more responsible for the impulse. The impulse can no longer be related to the X-ray silhouette although it still remains an index of the heart's size.

A well defined maximal impulse, the outer border of which is situated $4\frac{1}{2}$ inches or more to the left of the mid sternal line or quite clear of the nipple line in an adult male, is to be taken as a definite sign of the heart's enlargement in the absence of cardiac displacement. The sign is all the more emphatic if it is discovered in the 6th interspace, it is less emphatic if found in the 4th.

The idea that percussion gives a projected image of the heart could only apply to small hearts percussed on the front of the chest. When the heart is enlarged percussion wanders farther into the left axilla, and the stroke is now towards the side of the heart and not towards its front, it is not a margin of the heart that is being percussed. Margins of the heart are purely imaginary, its walls merely project certain distances from the mid line in given planes. It is incorrect to speak of the border of the heart. All we can expect to do by percussion is roughly to map out the region of the chest wall that is approached rather nearly by the heart and by the great vessels. An increase of the cardiac dulness to the left is an important diagnostic aid, but should never be allowed to outweigh the evidence of a well defined maximal impulse. Thus the chief value of percussion to the left is in bringing confirmation when the impulse is ill defined and in ascertaining the approximate extension of the heart towards the axilla, when no distinct impulse is available.

In marking out the position of the heart by percussion, moderately firm percussion should be used. The left border of dulness is ascertained by percussing the 5th sometimes the 6th, intercostal space, starting well out in the axilla, and working forwards and inwards in the chosen space. The outermost normal limit is $4\frac{1}{2}$ inches (11 cms) from the mid line.

In marking out the right border of dulness the upper border of the liver to moderate or rather lighter strokes is first ascertained in the right nipple line. Percussion is now conducted inwards immediately above this liver dulness. Dulness corresponding to the right auricle is generally encountered $1\frac{1}{2}$ to 2 inches (3 to 5 cms) from the mid line, in the 5th (or sometimes the 4th) interspace. The chief information to be obtained from the right border is that the heart is or is not displaced. Increased dulness to the right occurs in cases of auricular enlargement, especially in mitral stenosis.

It is usual to percuss the upper border. The method has little value and it is open to the objection that the upper border is obliquely set. It is of much more practical value to percuss as a routine the 2nd left space from left to right and to cross the sternum. The 3rd left space can be percussed similarly if desired, and should be in special cases. The 2nd space should be resonant throughout and its percussion will disclose any material extension of the cardiac dulness upwards. Simultaneously it reveals the dulness of a dilated aorta, which under the ordinary routine of percussion is often overlooked.

when the maximal impulse is in its normal place. Simple augmentation of the heart beat though producing a diffuse impulse in the spaces rarely moves ribs or cartilages appreciably. Prominence of the precordial ribs and of the lower sternum is frequent when the heart has become much enlarged during the period of growth before the chest wall has hardened to its final shape.

Epigastric Pulsation occurs in one of three distinct forms. It may be transmitted from the abdominal aorta and is then felt as a forward systolic thrust occurring a little later than the cardiac impulse. It may result from pulsation of the liver and is then visible but rarely palpable and is associated with abundant signs of venous congestion. It may be transmitted from the heart; in that case the movement is almost always a systolic retraction and pulsation is felt against the upper margin of fingers pressed into the epigastrium. Thus the usual cardiac form of epigastric pulsation is valueless in estimating the size of the heart being produced very readily by overaction of the heart and occurring in many perfectly normal people. In cases of hypertrophy of the heart a strong downward systolic thrust from the heart is sometimes felt by fingers that sink into the epigastrium.

Percussion—This method of measuring the heart has much value if used reasonably. It is however a highly subjective method and has an error that is often considerable. It is crucial in measuring to know the error of the method; to have but an inaccurate measure may be regrettable but to have it and not to know it is deplorable. Very fantastic ideas have been and still are held of the accuracy that can be attained by percussion. It is illuminating to practise percussion while blindfolded and repeatedly to percuss in a rib space chosen by some critical observer who chalks the margin of dulness on the chest wall as it is estimated. It does not require much of this experience nor long comparison with orthographic records to know that great accuracy is impossible. The left wall of the heart often moves in by as much as 1 centimetre in systole; it often moves more than this with respiration. Yet we take no heed of the phase of the cardiac or respiratory cycles in which strokes are delivered. A left margin of dulness coinciding with the maximal impulse is usual enough and to be expected since this impulse often forms the outermost point of the heart that is uncovered by lung. The impression of a sharp margin of dulness in other rib spaces is mostly fictitious; this must be so in that there is a gradual recession from the chest wall as the intervening lung tissue gradually increases in depth.

nearly all show auricular fibrillation, half of them present mitral stenosis and most of the remainder some thickening of the valve and widening of the ring. The auricular wall is often of the thinness of paper, and fibrous tissue replaces its muscle. It is because the muscle has perished that the dilatation occurs. The left auricle in expanding bulges to the right behind the right auricle, which normally lies in front of it, into the lower right pleural cavity. The condition is important in that it may be mistaken for pleural or pericardial effusion and be tapped. Dulness is extensive to the right of the sternum, from




FIG 2 —Electrocardiograms from leads I II and III illustrating preponderance of the left ventricle. R is tallest in lead I and shortest in lead III. S is deepest in lead III.




FIG 3 —Electrocardiograms from leads I II and III illustrating preponderance of the right ventricle. R is shortest and S is deepest in lead I. R is tallest and S is least deep in lead III. Time in $\frac{1}{4}$ sec.

the 4th rib or space downwards the border of dulness extends as a convex line outwards and downwards through or beyond the right nipple, the whole of the lower axillary region may be dull. The dulness at the base posteriorly is usually less distinct. Other signs helping at the bedside to differentiate the condition from effusion are cardiac pulsation and clearly audible heart sounds over the anterior area of dulness.

DISTINCTION BETWEEN HYPERTROPHY AND DILATATION IN CHRONIC CASES

In considering whether hypertrophy and dilatation can be separately assessed in chronic heart cases during life we deal with questions of measurement. The only reliable way of measuring the

amount of muscle in the heart is to measure it after death, we weigh the muscle after death and thus is our measure of the hypertrophy that was present in life

In measuring the size of the heart during life we are measuring the size of the organ with its blood content. Our gauge is one relating to the heart's volume. By measuring in several planes we might arrive laboriously at an estimate of volume; this would include the volume of the muscle and the volume of the contents. Volumetric measures after death are fallacious because the heart is very apt to distend in dying and because much blood may run into it after it has ceased to beat.

How is dilatation to be gauged? If a heart of normal volume swells acutely then the measure of increased volume is a simple measure of dilatation. The dilatation that happens during a severe paroxysm of tachycardia can be gauged approximately by repeated observation during the progress of the paroxysm. But no estimate of the amount by which the heart has dilated can approach accuracy if it depends upon observation confined to the end of the attack since the original size of the heart will be unknown. Thus in a case of chronic cardiac enlargement our only hope of obtaining an accurate measure of dilatation during life is to estimate the volume of the heart during life and a little later if opportunity offers after death to subtract from this the volume of the muscle. Hypertrophy can be measured after death; volume might be measured during life. Dilatation could be measured by combining these two but it has not been done. It is to be observed that physical signs, purporting to indicate degrees of dilatation in chronic cases, may not be introduced into the argument for any that we may possess are unreliable in that they remain unchecked against a proper measure. We cannot assess hypertrophy or dilatation where both occur together with any real accuracy either during life or after death.

COMPENSATION

More than a hundred years ago the view was put forward that when a heart has gained, in virtue of its hypertrophy, as much as it has lost by reason of its dilatation, there results a sort of compensation or equilibrium which maintains the cardiac function in its normal state. The view, thus expressed, gradually became more dogmatic, governing the interpretation and even influencing the recognition of cardiac failure. If it were held, in the case of aortic regurgitation that hypertrophy is a process of 'compensation',

enabling the heart to meet the added load, no exception could be taken to such use of this word. But the word stands for a fuller hypothesis. It is that if dilatation appears, the circulation is imperilled, and that the danger continues until a corresponding degree of hypertrophy is added, then the balance is restored, and the heart again functions normally. The later appearance of adverse symptoms is heralded as "broken compensation" and implies that the balance between dilatation and hypertrophy is again upset. It is to be presumed that the balance is upset by a gain of dilatation rather than by a loss of hypertrophy. The hypothesis is open to grave objections, for this reason, and also because it has influenced diagnosis prejudicially, it is necessary that it should be abandoned as a clinical tenet.

When physiology teaches as it now does, that the normal heart in meeting extra work dilates, and thereby increases its capacity for work, it utters a clear warning that the hypothesis is not unassailable, for the essence of the hypothesis is that dilatation is deleterious. Again, if it is agreed that every breach of compensation is remedied by a corresponding hypertrophy, it is clear that each event of this kind must leave the heart larger. Now this is definitely not the case, it is not necessarily the case even after repeated attacks of cardiac failure with congestion. The fact is that it is not dilatation that is injurious but that which is responsible for dilatation. Dilatation of the heart is the result rather than the cause of failing efficiency. When a patient suffering from fibrillation of the auricles loses his venous congestion under digitalis, he does not do so because hypertrophy develops and balances dilatation; neither does he do so because dilatation of the heart disappears. Venous congestion disappears because with the lower rate of beating the heart acts more efficiently, and if the heart shrinks, as it may sometimes in the process, this is not the cause of a restored balance but a secondary effect of longer diastoles and more powerful systoles. The use of the term "compensation" at the bedside conjures up or fosters the belief that, by signs derived from the chest wall, judgment upon a balance between dilatation and hypertrophy can be passed, such a belief is grotesque and, in so far as it is believed, it obstructs diagnosis. It is evident that many use the terms "compensation" and "broken compensation", or its ugly equivalent "decompensation", at the bedside, and mean no more than to infer that venous engorgement is absent or present, but that is not what is to be understood. A plain observation is garnished with an assumption

and served up in an unrecognisable form this too prevalent habit is pernicious for it destroys precise thinking Terms involving unproved hypothesis should be eschewed and in this instance reference should be to the simple presence or absence of venous congestion By this directness cumbersome needless and in fact erroneous assumption is avoided and simplicity and exactitude of expression are secured The diagnosis of venous congestion is within the easy reach of all it is a simple expression of what we see The diagnosis of broken compensation in its proper sense is within the reach of none

DEGREES OF CHRONIC ENLARGEMENT

It has been seen that an estimate of the relative degrees of muscular growth and stretching in chronic enlargement of the heart cannot be made Moreover it is unnecessary in the sense that the goal at which the attempt is aimed is reached more quickly and more certainly by examining the veins It is probable that in most instances of permanent enlargement hypertrophy and dilatation coexist and are more or less proportioned to each other for weight can be guessed from the volume of the heart without gross error during life The chief example of inaccuracy is the instance in which a heart definitely pronounced small during life weighs much more after death than was expected These instances are rare

In patients suffering from or examined to exclude chronic disease of the heart we can undoubtedly estimate the size of the heart in a broad way in most instances by simple bedside signs Great accuracy in measurement as we have seen is unattainable were it attainable it would serve no very practical purpose It is rarely a matter of much consequence to treatment and never in a chronic case whether we diagnose the heart of normal size or very slightly enlarged The idea that a fine gauge is needed is erroneous a coarse gauge suffices If the right border is found in its proper place and the outer margin of the impulse and left percussion border lie $4\frac{1}{2}$ inches or more to the left of the mid line or if these lie a lesser distance to the left and there is unmistakable thrust of bony or cartilaginous structure *definite enlargement* (meaning slight but unmistakable enlargement) of the heart can be diagnosed If in similar circumstances the left border lies farther out and the impulse farther out and perhaps down and there is perhaps heaving of ribs or sternum *moderate enlargement* can be diagnosed When the impulse and percussion border are far out in the axilla and there is forcible movement of much of the chest wall *considerable enlargement*

can be diagnosed. The subdivision is an arbitrary one and subject to minor modification but it serves to place patients, from the standpoint of the heart's size, in a few simple categories which suffice for all practical needs. It is far better to concentrate upon this relatively simple end result than to attempt fine distinctions of more than doubtful value.

PROGNOSIS

Dilatation—It follows from the manner in which pathological dilatation of the heart is produced that it is significant of inefficient beating, as this is produced diversely so the outlook varies for the manner in which dilatation is brought about must be taken into consideration. When dilatation is due to very rapid heart action, prognosis is really concerned chiefly with this rapid action and its cessation for when the heart begins to beat slowly again it will cease to beat inefficiently. Dilatation in itself does not spell disaster unless it is extreme and extreme dilatation is usually prevented by an unstretched or normal pericardium. The circulation can be carried on for long periods as paroxysmal tachycardia illustrates while the heart is much dilated and recovery is prompt and complete at the resumption of normal beating. Similar considerations apply when the heart is dilated during infectious illnesses the dilatation is a measure of the severity of the poisoning appreciable dilatation occurs only when intoxication is reaching a dangerous level. It is natural to regard such a condition as grave but attention should not concentrate on the heart other parts of the organism are simultaneously suffering. The seriousness of the situation lies in the course the infection is taking rather than in the dilatation itself. Dilatation as seen in infectious disease rarely reaches the grade that may occur temporarily in paroxysms of tachycardia except in the terminal and hopeless stages of the infection. These examples sufficiently illustrate prognostic considerations in acute dilatation of the heart.

Enlargement of the heart in cases of chronic heart disease is an important guide to prognosis. In considering this matter it is sufficient to enquire if the subjects of a group presenting enlargement live longer than subjects of a group presenting none. The comparison should be between two series of patients exhibiting similar signs of valvular disease similar exercise tolerance etc., but differing only

In the chronic disease of adults it is remarkable how little the actual physical signs of enlargement change from year to year. Enlargement is not a gradually progressing malady. Enlargement comes in the early stages of the disease and tends to remain fixed in its degree. Hypertrophy in response to a constant new burden is now held to develop fully within a few weeks. The reasons why enlargement affects prognosis are that enlarged hearts are far more prone than are small ones abruptly to display signs of failure, that they are more apt to suffer from thrombotic accidents, coronary or endocardial, the latter leading to embolism, and that they tend more to acquire serious disorders of beating. It has long been held, and I think rightly, that the muscle of a greatly enlarged heart is essentially unsound, but this statement is based, not upon histological or experimental evidence, but upon the experience that large hearts are not dependable.

The place taken by enlargement in the prognosis of chronic heart cases in general is given in the final chapter.

TREATMENT

Treatment in cases of acute dilatation should be governed by the origin of this dilatation. If it arises from a paroxysm of tachycardia, then the paroxysm is to be treated; once this has ended and, as usually happens, the dilatation has disappeared completely within a few heart beats, the fact that the heart has been dilated does not call for special treatment. Patients who feel exhaustion will want to rest for a day, others will soon feel quite well and may be allowed to go about their ordinary affairs. If the dilatation results from an infection, then it is the latter to which the remedies must be applied and which by its course and severity will dictate the period of rest. If it comes from asphyxia, then any obstruction of the respiratory passages that is present must be removed. These are the only common ways in which acute dilatation of the heart arises, and treatment corresponds. There is no acute condition to which we can suitably apply the term 'idiopathic' dilatation, or that calls for special treatment and after care (for Heart Strain see Chapter XVI), but if acute dilatation is so severe as to bring with it great venous engorgement, then, if it cannot be attacked at its source, special remedies for congestion may come under consideration. These have been discussed already under heart failure.

Enlargement of the heart, as a chronic malady, is beyond curative measures. Those who display it must live well within the limits of

their reserves. Patients who exhibit slight but definite enlargement of the heart may have good exercise tolerance, but they should be restricted to the milder forms of exercise. Patients exhibiting moderate enlargement very rarely possess good exercise tolerance, and those with considerable enlargement never do, the activities are to be limited correspondingly.

CHAPTER XIII

DISEASE OF THE AORTIC VALVE

PATHOLOGICAL ANATOMY

DISEASE of the aortic valve is of several types. Rheumatism which affects the valve in young people inflames it and causes numerous verrucous vegetations to appear along the line of apposition of the cusps. Cusps so affected ultimately thicken and shrink and often become adherent to each other at their commissures. Shrinkage produces incompetence, adherency produces stenosis. Secondary calcareous deposits may occur in these same cases and these also cause stenosis. Syphilis affects the aorta in middle life; the aorta and its ring become dilated, the cusps lengthen correspondingly but are no longer wide enough to meet at their margins; which though opaque may be only a little thickened, sometimes the cusps are greatly thickened especially at their margins, shrunken and retracted against the aortic wall. Syphilis yields free regurgitation rather than stenosis. In cases of arteriosclerosis of the aorta this process may spread to the valve cusps along their commissures thickening them and rendering them incompetent. A severe type of ascending degeneration of the cusps (page 244) is not infrequent in elderly subjects; large nodular calcareous deposits are developed in the bases of the cusps and frequently cause stenosis. In infective endocarditis the valve substance is in part destroyed by ulceration; in part it may be distorted by accompanying fibrosis in the subacute or chronic case in which the formation of new vegetations and healing proceed together.

AORTIC REGURGITATION

In aortic regurgitation and especially in free regurgitation the pressure at the end of the pulse cycle (diastolic blood pressure) is low, owing in large part to the reflux of blood from aorta to ventricle and in part as we shall see later to vasodilatation. To main-

tain a full circulation such blood as regurgitates must be thrown out again as well as the normal ventricular content. Thus an increased quantity of blood is ejected rapidly into relatively empty arteries especially when regurgitation is free. The top of the pulse which represents systolic blood pressure is high. The pulse pressure (*the difference between diastolic and systolic*) is great.

SYMPTOMS

There are few symptoms that belong properly to incompetence of the aortic valve. At the most we may recognise those that arise from the unusual momentum of the issuing blood and from the rapid expansion of the arteries. Consciousness of the heart beat (palpitation) is usual and may be severe. Throb is often felt in the neck and head and beating is heard in the ears. There may be a general throb of the body with each heart beat. The momentum of the blood's movement may suffice to shake the bed or couch on which the patient rests and he hears it. Occasionally the patient is aware of a rhythmic noise within the chest. Dizziness is not uncommon and faintness may be felt while the patient is standing.

SIGNS

There are two signs that stand out as of chief value in diagnosis and these are the characteristic steep rising pulse of free regurgitation and the diastolic murmur at the cardiac base. With both these signs present the diagnosis is certain; with either one or other it is probable.

Water hammer¹ Pulse—In free aortic regurgitation the quantity of blood ejected at each beat is in excess of normal and it is ejected by a powerful ventricle into an aorta in which pressure is unusually low; the pressure therefore rises with abnormal rapidity and to an unusual height. Thus the characteristic pulse is large, jerking, abruptly to its summit and representing a greatly increased range of pulse pressure. The upstroke of the pulse is much steeper than that of the normal pulse (Fig. 24) and steeper still if the wrist in which it is felt is held aloft. If the whole wrist is grasped with the fingers lying over the front of it the normal pulse is scarcely felt but the aortic pulse slaps the examining fingers sharply and gives often a distinct and correct impression of an accompanying thrill.

(Fig. 24) All the pulses of the body jerk forcibly, as does the radial, and the movement is clearly to be seen over the line of the carotid and brachial arteries. The carotid artery may move the lobe of the ear and the pulsation be continued actively into the superficial temporal vessel. The abdominal aorta throbs violently. As it travels, the pulse wave rises more and more steeply and to a higher crest, and so it frequently happens that the abrupt jerking (or water hammer) quality is extreme in the dorsum of the foot, for the same reason the systolic pressure comes to be higher, often much higher,




FIG. 24.—Tracings from three selected optical records from the radial pulse. The highest curve is the pulse of a case of free aortic regurgitation (AR) and is characteristic. It is of extreme abruptness in its upstroke which continues in a series of oscillations to the summit. It is a perfect example of a water hammer pulse.

The lowest curve (AS) is from a case of aortic stenosis and is anacrotic in form, rising slowly over the shoulder to a very delayed summit.

The third curve (N) is a normal curve for comparison with the remainder. The vertical lines represent time in $\frac{1}{2}$ of a second.

in the leg than in the arm. Many other signs are recorded that belong to the same category, for example, the "pistol shot" sound, heard when a stethoscope is pressed over an artery, and the audible pulse in the palm; there is no need to cite them all, for they express one and the same phenomenon and are in reality redundant signs.

Because the summit of the pulse rides high, the fall of pressure to the diastolic notch is more rapid than normal, a feature that has led to the use of the name "collapsing pulse." Actually, exaggeration of the rate of fall is far less than is exaggeration of the rate of rise, and cannot be detected with the finger, the term "collapsing" detracts attention from the essential quality of the aortic pulse and should be abandoned.

The pulse in aortic regurgitation deserves close study, not only because it is so important for diagnosis but because it provides the only satisfactory guide to the degree of aortic incompetence. From the full arterial signs described there is a complete transition to the arterial signs that appear in healthy people, in very slight aortic reflux the pulse pressure is normal in extent and the upstroke normal in rapidity.

Diastolic Murmur—The characteristic murmur of aortic regurgitation begins abruptly in early diastole and fades away. It is usually a long, smooth, rather loud, blowing and high pitched murmur, sometimes it is much shorter and far less audible and requires very close attention to detect it. It is heard best over or near the sternum at the level of the 2nd rib or space to right or left of the sternum, or in the 3rd or 4th spaces to the left of the sternum. The stereotyped phrase that the murmur is maximal at the 2nd right cartilage is inaccurate. It is also often heard along the right border of the sternum, or may be traced over the precordium as low as the cardiac impulse. In the region of the 2nd rib it is almost always preceded by a systolic murmur, and this is so whether stenosis is present or not. It is instructive to note, as a comment upon the timing of murmurs (page 133), that there is no murmur that gives rise to more frequent differences of opinion or to more changes of opinion by one and the same observer, than this diastolic murmur when isolated, and especially when the heart is beating a little rapidly. Happily, the systolic murmur rarely does fail. If the stethoscope is placed over the 2nd right cartilage and to and fro blowing murmurs are heard, it is sufficient there is then no need to attempt the timing of these murmurs—the first is necessarily systolic and the second necessarily diastolic. The diastolic murmur is often heard more loudly than the systolic, and at lower levels. When the systolic murmur fails, actual timing of the murmur becomes quite essential to diagnosis, for its character will not betray its position in the cycle.

The diastolic murmur in cases of suspected aortic regurgitation should be sought for diligently along both borders of the sternum, and especially along the left border, where it is particularly apt to appear in cases of slight regurgitation that do not affect the pulse. It should be sought in the erect and lying posture, and particularly with the breath held in deep expiration. Even so in some cases it will prove elusive, vanishing completely from time to time.

The degree of incompetence cannot be gauged either from the

loudness or the length of the murmur. When the pulse indicates free regurgitation the 2nd sound is often but not always inaudible at the aortic cartilage. Ventricular enlargement is the rule in these cases but its extent is too indirect an evidence and is in fact quite unreliable in gauging the state of the aortic valve. It is the pulse that most reliably indicates the degree of regurgitation.

A diastolic murmur is sometimes heard over the main artery of a limb in case of conspicuous aortic incompetence; it is produced beneath the stethoscope and is not heard except when other signs of aortic reflux are quite obvious; consequently it has little value.

DIFFERENTIAL DIAGNOSIS INCLUDING PULMONARY REGURGITATION

The Pulse—The characteristic signs found in the arterial system in aortic incompetence are repeated fully in only one condition, namely, arteriovenous anastomosis. Here the leak is from artery to vein rather than from artery to heart. Every arterial sign of aortic reflux that has been named is repeated in free arteriovenous anastomosis and repeated in its extreme form. The pulse has abruptness though in less degree in some cases of patency of the ductus arteriosus where the leak is from aorta to pulmonary artery it is also abrupt though not extremely so in conditions of simple peripheral vasodilatation. Aortic reflux, arteriovenous anastomosis and patent ductus arteriosus (page 262) are distinguished by the signs that occur directly over the corresponding lesions.

The Murmur—A low pitched blowing early diastolic murmur in the apical region strictly confined to the impulse in a case of auricular fibrillation with low ventricular action is due to mitral stenosis (see page 134).

An early diastolic murmur heard in the 3rd and 4th left space in cases of mitral stenosis should be attributed even when the pulse is unaltered not to pulmonary but to slight aortic incompetence. The murmur described by Steell and attributed to pulmonary regurgitation in mitral stenosis cannot be differentiated from an aortic murmur in the living subject. The pulmonary valve is very rarely deformed except by congenital disease (page 265) or by acute infective endocarditis (page 191). When infection is certain and there are no arterial signs of aortic disease or of systemic embolism a diastolic murmur at the pulmonary cartilage will sometimes render pulmonary vegetations suspect.

The to and fro murmur of aortic valve disease is sometimes difficult to distinguish from to and fro friction sounds (described on

page 174) when these are confined to the base of the heart. Difficulty in diagnosis occurs in cases of subacute rheumatism when a soft to and fro murmur develops in this situation and as yet other signs of pericarditis are obscure. The point resolves itself with time. There may also be difficulty in the same infection when widespread friction sounds have developed and, clearing away, leave a persistent to and fro murmur at the base, which may be interpreted either as pericardial or aortic in origin. The presence of much regurgitation will be recognised or excluded by the pulse. Uncertainty as to the presence of slight aortic valve damage, when the heart is known to have been exposed as a whole to rheumatic inflammation is of little consequence.

RUPTURED AORTIC CUSP


Spontaneous rupture of a healthy aortic cusp does not occur. Most instances of rupture are of cusps that have been affected by syphilis (or subacute bacterial endocarditis); a little aneurysm forms in the centre of a cusp and projects towards the ventricle. One day this aneurysm breaks or a diseased cusp is torn from its edge to its base. The accident is most likely to occur when the subject is making a considerable effort, thus raising aortic pressure. In such a case the subject may feel as though something has broken in the chest; he becomes very breathless, or falls in a state of collapse or actual unconsciousness, and on occasion shows signs of acute dilatation of the heart. Sometimes from the moment of the accident, sometimes a little later, the patient may notice a rhythmic noise within his own chest. On examination the Wassermann reaction is generally found to be positive, and signs of regurgitation, usually free, are discovered. The diastolic murmur may be so loud that it can be heard by the ear placed near to the chest wall; it is often musical in quality (Fig. 25) and is associated with thrill.

CONSECUTIVE PHENOMENA

Rapid Pulse — In describing the signs of aortic regurgitation certain phenomena have been omitted because they are not the direct and immediate consequences of regurgitation, though they often follow from it. A chief consequence of free aortic regurgitation is a low mean blood pressure, the low pressure is the usual cause of the increased pulse rate that is common in this state, though it may also arise from infection.

Vasodilatation — A phenomenon usual in free aortic regurgitation is vasodilatation, exhibited especially in the face, also in the upper

limbs and sometimes more generally. The facial skin is of a fresh colour is a little or sometimes greatly flushed it is warm or hot. The idea that aortic regurgitation causes pallor is erroneous it arose at a time when subacute infective endocarditis was unrecognised for the latter disease affects the aortic valve and often causes intense pallor (pages 186 and 189) so does long continued rheumatism. It is mainly because of associated vasodilatation that capillary pulsation (page 40) is so frequent in free aortic regurgitation this pulsation is usually to be seen occurring spontaneously in the cheeks and forehead and can be elicited readily in the lip and lobe of the ear by gently pressing a slide upon these parts it is also seen though less frequently and conspicuously in the fingers. It is only to be found



sounds are indicated. Time in $\frac{1}{4}$ sec

in parts of the skin that are warm or hot and never in those that are cold thus it is clearly linked with relaxation of arterioles. Doubtless the large pulse pressure helps to render it conspicuous but large pulse pressure is not its cause. Capillary pulsation is seen equally well in cases of free union between the superficial femoral artery and vein here too there are both high pulse pressure and the signs of vasodilatation. If the common femoral vessel is obliterated by pressure and the arteriovenous anastomosis thus cut off from the

Enlargement—A measure of cardiac enlargement is clearly consecutive to aortic regurgitation but has been more conveniently discussed in a previous chapter (page 106)

Anginal Pain occurs in many cases showing aortic regurgitation

it might be just to regard this pain as determined in some cases by the extra work that the valve defect imposes, but its inclusion in the symptom complex of aortic incompetence would not be in the best interest either of simplicity or clarity.

AORTIC STENOSIS

It is a sound rule rarely to diagnose conditions that occur rarely. Aortic stenosis is far less common than aortic incompetence and its diagnosis should be infrequent. It occurs chiefly in elderly subjects, occasionally in the rheumatic hearts of younger subjects.

SYMPTOMS, SIGNS, AND DIAGNOSIS

There are no symptoms of aortic stenosis.

Anacrotic Pulse—The outstanding diagnostic sign of aortic stenosis is a small pulse, rising slowly to a delayed summit, the condition should never be diagnosed without this sign. An experienced finger will detect it so often will inspection of pulsating arteries. Sometimes an earlier summit can be detected on the slow upstroke by palpation, but more often in sphygmographic records (Fig. 24) the pulse is then termed "anacrotic." This change in the pulse's form is the direct consequence of stenosis, the degree of change is related to the degree of stenosis. It is for this reason that the sign is so essential for, if the pulse rises quickly, such stenosis as may be present is not worth diagnosing as such. The slow rising, and the anacrotic, pulse may occur in any other condition in which there is an incomplete and local obstruction anywhere between the aortic valve and the point on an artery at which the pulse is felt. The mechanism is always the same—the primary wave of the pulse is spent in overcoming the obstruction, for this reason systolic pressure is lower than it otherwise would be. No other lesion, however, produces this pulse with anything like the same frequency as does aortic stenosis. In aortic aneurysm, through clot or through constriction of the mouths of branches arising from the sac obstruction may occur in outgoing vessels. The obstruction, however, is usually confined to one, less frequently to two of the three vessels that issue from the aorta, namely innominate, carotid, and subclavian artery so it is not very common to find the pulse universally slow rising in aneurysm, or to find it rising with equal slowness in the two arms. A universal anacrotic pulse occurs as a rare phenomenon in cases of advanced arterial disease with a weakly acting heart. In all the different

conditions named, including aortic stenosis, the systolic blood pressure, like the peak of the pulse, is reduced in level.

Systolic Thrill —The second sign is a systolic thrill at the base of the heart, usually maximal at the 2nd right cartilage or space. This sign is usually to be detected if carefully sought, by placing the palm of the warm hand freshly on the base as the subject is completing a deep expiration to hold respiration in this position. Though it is an expected sign, diagnosis may proceed without it, but a diagnosis cannot be made by means of it alone. It can occur in aortic valve disease when there is little or no stenosis, and occasionally when there is no disease of the valve, but anaemia is present.

Systolic Murmur —Every case of aortic stenosis presents a systolic murmur, heard loudest where the thrill is best felt and carried up to the clavicle or beyond into the neck along the carotid vessel. A harsh murmur is the hint to examine for thrill. The systolic murmur is a guiding sign but has no further value in diagnosis. A systolic murmur at the aortic cartilage is usual in any form of aortic valve disease in dilatation or aneurysm of the ascending aorta, in atheroma of the base of the aorta. It is quite common in anaemia and in any condition in which the action of the heart is augmented. To diagnose stenosis on the basis of such a murmur is unpardonable.

Presence of Aortic Regurgitation —While the slow-rising pulse is an essential sign it is not quite sufficient. The systolic thrill locates the obstruction and thus aids diagnosis, a harsh murmur may be used in the same way. The presence of the diastolic murmur of incompetence can be used similarly to locate the disease, for stenosis rarely occurs without this sign of regurgitation. It is to be noticed that the murmur is cited alone, for an abruptly rising pulse is sufficient evidence that there is no material stenosis.

CHAPTER XIV

DISEASE OF THE AURICULO VENTRICULAR VALVES

PATHOLOGICAL ANATOMY

Mitral Valve —Disease of the mitral valve usually results from an inflammation of the valve occurring during the course of active rheumatic disease (page 195) In the early phases of its inflammation, the two cusps present narrow bands of small sessile and closely set vegetations along their surfaces of apposition Beneath and in the substance of the cusp is a more widespread inflammation The inflammation leaves, in healing, a thickened margin to the cusp, the thin and flaccid edge that is so necessary to perfect closure being destroyed Early thickening and rounding of the valve margin is a recognised cause of mitral regurgitation Repeated or long continued inflammation leads to higher grades of valvular deformity, the margins become more thickened and adhere together at their adjacent borders, at first the finest subdivisions of the chordae tendineae, later coarser subdivisions, become matted together and to the cusp against which they lie The process spreads down the main chordae, and these and the cusps become cemented and encased in new tissue Gradually the valve apparatus is converted into a contracted and rigid structure, and pierced by a buttonhole or funnel shaped orifice that can neither close nor open This is stenosis The rheumatic inflammation that first produces crumpling with regurgitation ultimately produces stenosis, it is one disease presenting a variant picture at different stages of its progress

There is a condition of atherosclerosis of the cusps, and especially of the aortic cusp of the mitral valve, that sometimes proceeds to calcareous deposit sufficient to cause stenosis Calcareous valves in elderly subjects are sometimes of this kind, they may be formed earlier by deposition of lime salts in tissues sclerosed by rheumatism Syphilis does not damage the mitral valve

In enlargement of the left ventricle it is frequent to find the

chamber and auriculo ventricular ring dilated after death and the whole mitral valve enlarged its chordae long, the valve curtains having a wider sweep but their substance thicker and more opaque than normal. Such valves are probably often incompetent we have however no adequate test of mitral incompetence at autopsy.

Infective endocarditis ulcerates the valves and often cuts the chordae thereby rendering the valve incompetent or by heaping up vegetation may cause obstruction at the orifice.

Tricuspid Valve—The only common cause of lesions of the valve is rheumatism and it follows just as does mitral disease from a characteristic inflammation of the valve. The tricuspid however, is less frequently and far less severely involved than the mitral valve. When stenosis is present it is slight the thickened valve edge forming a fixed ring of considerable size severe mitral stenosis usually coexists.

MITRAL STENOSIS

ITS RECOGNITION

Although many symptoms may be complained of by patients suffering from mitral stenosis there are none that can be ascribed properly and usefully to this deformity of the valve. A number of patients with developed stenosis are quite free from symptoms and the discovery of the disease comes as a surprise. Only signs of the disease will be discussed here.

An abrupt and accentuated 1st sound at the impulse an accentuated or reduplicated 2nd sound at the pulmonary cartilage are signs commonly occurring in mitral stenosis but these signs alone or in combination with or without an associated history of rheumatic fever form an insufficient basis for diagnosis. They result oftener from augmentation of the heart beat than from mitral stenosis but they serve usefully as guiding signs to the latter condition suggesting closer scrutiny of the mitral valve than it would otherwise obtain. The presence of signs supposedly indicating hypertrophy of the right ventricle should not be cited as evidence of mitral stenosis this method of procedure is too indirect to be sound. One sign only justifies the diagnosis and that is the appropriate murmur. The corresponding thrill when this is present is of course to be included it is no separate phenomenon the ear hears and the hand feels much the same vibrations when these are both audible and palpable. The murmur of mitral stenosis is the only heart murmur that merits and requires much study.

A word first of all about the thrill. It is best felt with the palm of the hand freshly applied to the chest wall in the region of the impulse, to have value it should be sufficiently sustained to possess a distinct purring quality. It is unwise to accept as thrill anything but an unmistakable sign. The thrill has much the same distribution in the cycle as the murmur though it is not always perceived to last so long, it is more elusive than the murmur and less important in diagnosis.

The murmur is almost always loudest at the impulse or close to it, and is usually confined closely to this region, though sometimes heard over a wider area of the precordium. It will be described first as it is derived from a normally beating heart, and later, the sounds as modified by auricular fibrillation and other abnormal mechanisms will be discussed. The description, which in certain aspects differs from current teaching is based upon a long experience of auscultation and, perhaps uniquely, upon a close comparison of what is to be heard with what is to be recorded by microphonic methods in a large number of selected cases. The graphic method alone places the time relations of these murmurs beyond dispute.

With Normal Rhythm —The characteristic murmur is of low pitch usually rough rumbling or rasping and ends abruptly in a loud sound with which it is continuous. This sound is in fact the first sound of the heart, which being usually and often greatly accentuated is chiefly responsible for the idea that the murmur rises in a crescendo. It is erroneous to believe that the murmur is usually recognised by timing it, although this is steadily taught. Most people though capable of identifying the chief murmurs readily, can not and never will time murmurs reliably. They recognise this murmur of mitral stenosis the instant it is heard, as I do myself in routine work, by its low pitched abruptly ending noise. Much labour is lost and many ultimately fail to know this murmur through persisting in the effort to time, instead of learning to know it as one learns to know a dog's bark.

The murmur is due to the passage of blood through the constricted orifice of the mitral valve in diastole. The phases of diastole in which



may be attained for a shorter or longer part of diastole. What parts of diastole are filled by murmur depends mainly upon the rate of heart beat and upon the degree of stenosis. Given that a certain

amount of blood has to flow through the orifice each minute the velocity of flow will be greater as diastole becomes relatively shorter with faster heart rate in like circumstance the velocity will be greater if the orifice is smaller. In most cases of mitral stenosis, as these come under examination the rate of beating is rapid and the stenosis is moderate or considerable consequently the velocity of flow is high and it is the rule for the murmur to occupy all diastole, beginning with the 2nd sound and ending with the abrupt 1st sound (Figs. 26, 27 and 28). Names become stereotyped and it is still customary to hear this full diastolic murmur called presystolic although this term can rarely be used with accuracy. When the heart is beating at 120 per minute as it often does in our patient clinically it clearly cannot be used accurately for in these patients the interval is no more than $\frac{1}{2}$ second and presystole shall we say, $\frac{1}{4}$ second in duration. The point would not be stressed so much were it not important to remark that there is a manifest tendency, which is traditional for the medical profession to exaggerate the accuracy of its subjective methods of examination. In using any method of measurement it is essential to know the error of it as has been stated earlier in connection with the measure of cardiac enlargement (page 110) and the error in timing can scarcely be less it is usually more than $\frac{1}{2}$ second. Graphic registration and even very attentive auscultation directed especially to detect a gap between 2nd sound and murmur shows that murmurs confined to late diastole are comparatively rare. They are only displayed by hearts that beat slowly and in which the stenosis is not advanced. In rare cases in which the heart is beating slowly the murmur may appear in presystole and continue early diastole leaving a gap in mid diastole clear of sound.

Mitral stenosis is not usually diagnosed in children before the age of ten years. At this stage and for a few years the murmur characterizing mitral stenosis in these young subjects very often occurs in mid diastole (page 199). The reason for its curious position is imperfectly understood but it is well to know that this variation occurs.

Early Diagnosis—In teaching that the appropriate murmur is the only sign justifying a diagnosis of mitral stenosis, I do so subject to the reservation that the murmur is diligently sought, and that a reasonable effort has been made to elicit it. The murmur is heard with much greater frequency when the patient lies than when he stands and oftener when the patient lies upon the left side than upon his back. It is heard with especial frequency when one or other of these two last positions has just been assumed by a patient

who has been sitting or has just taken sufficient exercise to raise the pulse rate and the velocity of blood flow through the orifice as in sitting up and lying down a few times. Close attention should be paid to the sounds at the impulse during the first 5 or 10 beats



ECG tracing showing a regular rhythm with a P wave, QRS complex, and T wave. Below the tracing is a strip of film showing the mechanical movement of the heart, with labels 'P', 'R', 'T' and 'S' indicating specific points in the cycle.

of the heart immediately following the assumption of one or other of these postures after exercise. So often is the murmur audible in these circumstances only and for a few cycles only or so often does the murmur become unmistakable only under these same conditions, that no subject deemed likely to have mitral stenosis should be proclaimed free of it until these special tests have been employed. They

should be used in every case of suspected heart trouble where there is an accentuation or a reduplication of the 1st sound or of the 2nd pulmonary sound or where there is an apical systolic murmur. If they are so used as a routine, experience soon shows that the signs just named are useful guides in suggesting the special tests but once so used can be neglected as actual diagnostic signs of mitral stenosis. Under the routine here described which eventually rejects all signs but the appropriate murmur early mitral stenosis will very rarely be overlooked. What is of equal if not of greater consequence is that the likelihood of diagnosing this disease of the valve where none exists will be lessened greatly. There are of course cases in which owing to the presence of venous congestion, exercise or postural tests are undesirable it is of little consequence to such patients whether a diagnosis of mitral stenosis is made or not.

In Auricular Fibrillation — When the auricles are fibrillating the character of the murmur of mitral stenosis does not usually differ from the common murmur heard while the rhythm is normal. That is to say it is a low pitched rumbling or rasping murmur. In most untreated cases seen for the first time the ventricles are beating rapidly. In such circumstances the murmur occupies full diastole and ends in an abrupt 1st sound thus giving as usual a crescendo effect.

In cases of normal rhythm as we have seen the murmur usually occupies full diastole but with slow heart action may become confined to one part of diastole the favoured part of the cycle is then pre-systole. In auricular fibrillation when the ventricle slows the murmur likewise fails to occupy full diastole but the favoured part of the cycle is now early diastole. That is so because the auricles have ceased to function and to force blood rapidly through the orifice at the end of diastole and in these circumstances the entry of blood from the auricle is at its fastest immediately after the opening of the auriculo ventricular valves the auricles being then engorged with blood that has been dammed up in them during ventricular systole. The velocity soon lessens as diastole proceeds and so the murmur in question tends in general to be diminuendo. In cases of fibrillation where the ventricular action is very slow, the murmur is confined to early diastole in all cycles and for an unexplained reason, often tends to lose its rasping quality becoming soft and resembling more in character the murmur of aortic regurgitation. Usually the ventricle beats less slowly though irregularly and the murmur retains its characteristic roughness it fills the whole diastole of short cycles and fills early and mid diastole in long cycles tailing away and leav-

ing an appreciable pause between the end of the murmur and the subsequent 1st sound. This combination of sounds is extremely frequent and should be known thoroughly, it requires a little thought to understand it, to help its appreciation I have inserted a graphic record of such sounds taken simultaneously with an electrocardiogram (Fig 29)

When auricular fibrillation develops in cases of mitral stenosis with normal rhythm, a late diastolic murmur previously heard sometimes disappears or becomes much less distinct owing to the loss of auricular systole. It is well recognised that, in auricular fibrillation,




FIG 29 — Simultaneous electrocardiographic and heart sound record from the region of the apex

mitral stenosis may be concealed, the heart at times presenting no murmurs. The tests previously described will usually elicit the murmur, or render a doubtful murmur unmistakable, here, as in instances of normal rhythm.

The chief variation of the murmurs in mitral stenosis that have been discussed are summed up in the accompanying table, the commonest example being those in the 3rd and 4th lines.

MURMURS OF MITRAL STENOSIS

Heart	Normal Rhythm	Auricular Fibrillation
1 Rate slow, stenosis early	Presystolic	No murmur
2 Rate very slow, stenosis developed	Early and late diastolic	Early diastolic
3 Rate rapid stenosis developed	Full diastolic	Full diastolic
4 Rate moderate stenosis developed	Full diastolic	Full diastolic in short cycles, early and mid diastolic in long cycles

This table shows at a glance the part played by auricular systole in determining the phase of diastole in which the murmur appears.

The influence of auricular systole can also be studied in cases of mitral stenosis associated with heart block. If the interval between auricular and ventricular contractions widens a shorter or longer gap of silence is noticeable between the murmur and the 1st sound. If 2:1 block develops then two similar murmurs will occur the one in presystole and the other in early diastole and each the result of a corresponding beat of the auricle. Instances of these kinds are not very uncommon instances of mitral stenosis and complete block in which the murmur alters its position from cycle to cycle are rare.

Pituitary Murmur—This murmur may be defined as one that resembles the murmur of mitral stenosis and is heard in cases of free aortic regurgitation in which mitral stenosis is subsequently proved absent. The precise manner in which the murmur is produced is unknown it is not uncommon. Very rarely it may be as rasping as that of mitral stenosis and may so it is said be accompanied by a thrill usually it is of low pitch is rumbling in quality, occupies full diastole but is less audible than the similar murmur of stenosis. In cases of aortic regurgitation where a diastolic rumble is heard at the impulse it is impossible to tell the origin of the murmur from its character or timing alone. Those who aim at complete anatomical accuracy in diagnosis will come nearest to accomplishing their purpose by using associated information. If regurgitation is slight if the heart is small or there is a history of rheumatic fever or chorea mitral stenosis may be diagnosed if the heart is large or the aortic disease is suspected to be syphilitic in origin mitral stenosis should not be diagnosed. Actually the point is not of practical importance the decision taken one way or the other will alter neither prognosis nor treatment.

MITRAL REGURGITATION

SYSTOLIC APICAL MURMURS

Cardiorespiratory Murmur—This is the commonest murmur heard in the region of the apex of the heart. It is usually heard best at or a little beyond the impulse. Often it is audible over a wide area being frequently distinct not only over the precordium but in the axilla and at the angle of the left scapula. It is a short high pitched blowing sound. It is in reality part of a breath sound and is most audible or only audible in inspiration the normal vesicular murmur being

broken into two or more short murmurs by the heart beats. But until thoroughly known its respiratory relationship is generally overlooked, when a short and inconstant murmur of superficial quality is heard, its relationship to respiration, and the effect of suspended breathing upon it, should be observed. The ventricle at each systole creates a little vacuum in its neighbourhood and so increases the rate at which air enters the lung. It is a normal phenomenon when the heart is beating quickly and strongly, as in exercise or excitement in steep uphill walking, when the mouth is open, a sharp movement of the air past the larynx, at each systole of the heart, can often be appreciated by the subject. The murmur should be construed as the natural accompaniment of the overacting heart.

Exocardial Murmurs —Precordialsystolic murmurs that are short, seem superficial and have any scraping quality are usually regarded as produced outside the heart by roughening of pericardium or pleura, the opportunity of examining the suspected membrane does not often occur. When it does occur, it is sometimes, but by no means always, possible to find in them a lesion to which the murmur may be attributed, proof however, is rarely forthcoming. During life the distinction between systolic murmurs that arise within and without the heart, so called "endocardial" and "exocardial" murmurs, is often impossible, unless the latter are represented by the characteristic, widely heard, to and fro friction sounds of recent pericarditis, or are frankly cardiorespiratory.

Inconstant Murmurs —Not infrequently a soft, blowing systolic murmur is heard in the region of the cardiac impulse while the patient is in one posture only, usually the lying, sometimes the standing. Some murmurs of the same character vary much in intensity from day to day, others appear only after exercise. Such murmurs are usually audible over a relatively small area of chest wall, though this is not always so.

Constant Murmurs —A considerable proportion of systolic apical murmurs are relatively constant from cycle to cycle and from day to day, they are in general harsher and longer murmurs, and are heard over a wider area of precordium and axilla than the inconstant murmurs previously described. These are the murmurs that are usually regarded as chiefly signifying mitral regurgitation. Amongst them is the murmur that starts in mid systole, a delay that causes it frequently to be mistaken for one occurring in early diastole.

MITRAL REGURGITATION AND ITS SIGN

It is desirable to emphasise at once that in discussing the diagnosis of mitral regurgitation we are not discussing except indirectly the diagnosis of mitral disease the two problems must be dealt with separately

There are no symptoms to be attributed usefully to mitral regurgitation The only sign that can be named is a systolic murmur It has been said that in mitral regurgitation the 1st sound of the heart may be replaced by murmur graphic records always show the 1st sound The fact is that the ear in listening for the 1st sound is confused by a loud murmur

A real difficulty in attempting accurately to correlate certain systolic murmurs with mitral regurgitation lies in the frequent impossibility of stating from the appearance of the valve after death whether leakage has occurred or not during life From this stand point it may be said that if regurgitation can be declared to have happened in any one group of cases more than another, it is in the group in which the cusps are subsequently found so thickened that the rigid valve opening is incapable either of opening farther or of closing During life these cases of stenosis should present the sign of regurgitation The facts are these In about one third of the cases no murmur can be discovered in the remainder a systolic murmur is heard in the apical region This murmur may be quite local more often it may be followed well out to the axilla It may be soft almost to inaudibility it may be musical or it may be harsh and loud occasionally it may be associated with thrill It is usually constant from cycle to cycle and from time to time but is not always so Thus the sign of regurgitation is to be recognised as very variable from case to case Moreover all these varieties of murmur are heard from time to time in cases in which at autopsy no real fault can be found with the valvular apparatus It is then quite usual to suppose that the muscular ring surrounding the base of the valve is relaxed sufficiently to produce leakage In the past we have been near to believing on *a priori* grounds that a distinct systolic murmur at the impulse must be caused by regurgitation identifying a systolic murmur and then assuming it to arise out of regurgitation In most instances this assumption may be right in some instances it is wrong for there are other ways in which these murmurs are produced There are very definite elements of weakness in our way of diagnosing mitral regurgitation It is never very satis

factory to base a diagnosis upon a single sign and the position is insecure when the sign is not known to be of a high grade of reliability

If we are treating a case of acute infectious disease or a frail elderly person and we are sure in such that a systolic murmur has recently developed at or near the impulse we may be justified in interpreting this as the first indication that the heart is dilating and so it may be important but we are hardly justified in definitely drawing this conclusion unless simultaneously or later direct signs of enlargement can be found The systolic murmur is chiefly a guiding sign and as such may have a distinct value An open window may be such a sign to a police officer on night rounds he may suspect but does not necessarily conclude there is a thief in the house A fixed systolic murmur in cases of chronic heart affection has a similar value in that it is sometimes the first sign to attract attention and lead to close examination of the heart

Confidence in the value of a diagnosis of mitral regurgitation is reduced not only by frequent doubts as to whether regurgitation actually exists or not but by our total inability to state its degree if it is there There may be some grounds though they remain inadequate for claiming that harshness thrill audibility over a wide area or other characteristics mark the murmur of mitral regurgitation no claim has yet been made that particular qualities betray the slight or the large leak Assuredly if the diagnosis of regurgitation *qua* regurgitation matters it matters according to its degree

Briefly the diagnosis of mitral regurgitation has a very limited importance it may be useful as an early indication of muscular failure in acute infectious disease and occasionally in old age Further reasons will be given for concluding that it should not be allowed to modify either prognosis or treatment (page 146)

RECOGNITION OF MITRAL DISEASE

To recognise mitral disease by means of systolic murmurs it is first essential to be accurate in eliminating all murmurs cardio respiratory or not that arise otherwise than from the mitral valve it is secondly essential to differentiate clearly between a mitral murmur that is due to faulty ventricle or to diseased valve To regard any distinct systolic apical murmur as meaning a diseased valve is indefensible The statement that a constant long harsh systolic

murmur heard at or beyond the impulse usually signifies mitral disease and that this interpretation is more reliable when there is a history of rheumatic fever as strong a statement as can be defended

Accurate diagnosis of disease of the mitral valve depends upon the recognition not of mitral regurgitation but of mitral obstruction. It has been stated already that rheumatic fever produces both these defects and that the underlying conditions are but stages of one and the same disease. Thus diagnosis of rheumatic mitral disease is accomplished most accurately (and in speaking of




Fig. 30—Orthodiagram. Man aged 41. Weight 115 lb. Mitral stenosis (developed below the aortic knob (K) appear the projection of P the dilated pulmonary artery of C the enlarged conus and the outline of the ventricle (V); the apex of which is unusually small. The right auricular shadow (RA) is enlarged.




Fig. 31—Orthodiagram. Man aged 40. Mitral stenosis (developed aortic regurgitation (slight) and great enlargement of the heart. The right border is composed of ascending aorta (AA) which is displaced an enlarged right auricle (RA) and the beginning of the enlarged right ventricle (RV). The left border consists of aortic knob (K) dilated pulmonary artery (P) greatly enlarged conus (C) and ventricle (V).

accuracy we must take into account not only correct guesses but the elimination of incorrect guesses) at the present time by starting at stenosis and carrying back the diagnosis of this lesion as far as is possible. If the method of early diagnosis (page 132) is followed many cases that might otherwise pass as mitral regurgitation are transferred into the decisive category of stenosis by using care to unveil these early cases the attempt to interpret many systolic murmurs becomes needless. Of the cases that remain it is questionable if a diagnosis of mitral disease can be made accurately in a sufficient percentage to warrant the attempt upon the basis of present physical signs.

TRICUSPID AFFECTIONS

Tricuspid Stenosis —There is only one sign of this condition that has much value it is a murmur like that described for mitral stenosis, but occurring over the lower part of the sternum. Characteristically in these cases, a murmur of mitral stenosis is heard at the impulse, is lost, and the similar murmur of tricuspid stenosis becomes audible as the stethoscope is moved inwards towards and to the sternum. Absence of a tricuspid murmur does not mean that the valve is undiseased. A presystolic pulsation of the liver recorded graphically was once, but is no longer thought diagnostic of the lesion. I have not known the diagnosis, when made, to affect the management of any case.

Tricuspid Regurgitation —A constant blowing systolic murmur, confined to the region of the lower sternum and ensiform cartilage, is not uncommon in cases of venous congestion and is usually ascribed to tricuspid regurgitation. Much more often than not, in cases of general venous congestion no distinct murmur of the kind can be heard, moreover a similar murmur is audible from time to time, and especially after exercise, in people who are apparently in good health. It is probable that the tricuspid valve readily becomes leaky.

In the old theory of back pressure, the pressure was supposed to rise in the right ventricle before it rose in the veins, and the tricuspid valve was pictured as one of the last lines of defence. Incompetence of this valve was regarded as the signal for engorgement of the veins, and so tricuspid incompetence came to be used almost as an equivalent term for venous engorgement. The hypothesis neglected the facts that, by obstructing the outlet from the right ventricle, this chamber cannot be engorged without raising venous pressure correspondingly, moreover, the rise of pressure in venous

difficulties in the way of diagnosing tricuspid incompetence by auscultation are similar to those already stated for mitral incompetence, the diagnosis is in general merely a statement of the discovery of a murmur, which is then assumed, perhaps in many cases rightly, to indicate this regurgitation. The degree of reflux is not indicated, the influence of the reflux cannot be estimated unless it can be detected in deformity of the waves in the cervical veins, and this is doubtful

murmur heard at or beyond the impulse *usually* signifies mitral disease and that this interpretation is more reliable when there is a history of rheumatic fever, is as strong a statement as can be defended

Accurate diagnosis of disease of the mitral valve depends upon the recognition not of mitral regurgitation but of mitral obstruction. It has been stated already that rheumatic fever produces both these defects and that the underlying conditions are but stages of one and the same disease. Thus diagnosis of rheumatic mitral disease is accomplished most accurately (and in speaking of




FIG 30.—Orthodiagram. Man aged 41 weight 113 lb. Mitral stenosis (developed). Below the aortic knob (K) appear the projection of P the dilated pulmonary artery of C the enlarged conus and the outline of the ventricle (V) the apex of which is unusually round. The right auricular shadow (RA) is enlarged.




FIG 31.—Orthodiagram. Man aged 42 Mitral stenosis (developed) aortic regurgitation (slight) and great enlargement of the heart. The right border is composed of the ascending aorta (AA) which is displaced an enlarged right auricle (RA) and the beginning of the enlarged right ventricle (RV). The left border consists of aortic knob (K) dilated pulmonary artery (P) greatly enlarged conus (C) and ventricle (V).

accuracy we must take into account not only correct guesses but the elimination of incorrect guesses) at the present time by starting at stenosis and carrying back the diagnosis of this lesion as far as is possible. If the method of early diagnosis (page 132) is followed, many cases that might otherwise pass as mitral regurgitation are transferred into the decisive category of stenosis, by using care to unveil these early cases the attempt to interpret many systolic murmurs becomes needless. Of the cases that remain it is questionable if a diagnosis of mitral disease can be made accurately in a sufficient percentage to warrant the attempt upon the basis of present physical signs.

TRICUSPID AFFECTIONS

Tricuspid Stenosis —There is only one sign of this condition that has much value it is a murmur like that described for mitral stenosis, but occurring over the lower part of the sternum Characteristically in these cases, a murmur of mitral stenosis is heard at the impulse, is lost, and the similar murmur of tricuspid stenosis becomes audible as the stethoscope is moved inwards towards and to the sternum Absence of a tricuspid murmur does not mean that the valve is undiseased A presystolic pulsation of the liver recorded graphically was once, but is no longer, thought diagnostic of the lesion I have not known the diagnosis, when made, to affect the management of any case

Tricuspid Regurgitation —A constant blowing systolic murmur, confined to the region of the lower sternum and ensiform cartilage, is not uncommon in cases of venous congestion and is usually ascribed to tricuspid regurgitation Much more often than not in cases of general venous congestion no distinct murmur of the kind can be heard, moreover, a similar murmur is audible from time to time, and especially after exercise, in people who are apparently in good health It is probable that the tricuspid valve readily becomes leaky

In the old theory of back pressure the pressure was supposed to rise in the right ventricle before it rose in the veins and the tricuspid valve was pictured as one of the last lines of defence Incompetence of this valve was regarded as the signal for engorgement of the veins and so tricuspid incompetence came to be used almost as an equivalent term for venous engorgement The hypothesis neglected the facts that by obstructing the outlet from the right ventricle, this chamber cannot be engorged without raising venous pressure correspondingly, moreover, the rise of pressure in venous

difficulties in the way of diagnosing tricuspid incompetence by auscultation are similar to those already stated for mitral incompetence, the diagnosis is in general merely a statement of the discovery of a murmur, which is then assumed, perhaps in many cases rightly, to indicate this regurgitation The degree of reflux is not indicated the influence of the reflux cannot be estimated unless it can be detected in deformity of the waves in the cervical veins, and this is doubtful

Tricuspid incompetence forms an instructive example of an old time clinical philosophy. An intangible valvular defect draped with complex but unproved hypotheses has formed a screen, tending to conceal the essential. The essential is not tricuspid incompetence but general venous congestion and the precise information that is required for practical purposes is obtained, not from the precordium, but from the vein themselves for in this system be the earliest signs and the late signs of congestive failure. The vein provide direct signs and those thoroughly familiar with them use neither the tricuspid murmur nor the theory that goes with it.

CHAPTER XV

SIGNIFICANCE OF VALVE DISEASE

PROGNOSIS

THE undue emphasis placed upon disease of the cardiac valves in diagnosis was the chief reason why the prognosis of heart disease long remained so very unsatisfactory. This over emphasis resulted largely from an exaggerated notion of the extent to which valve defects burden the heart mechanically. The notion is dispelled by considering the extent of these burdens a little more closely. Take, for example, stenosis. Let a hole 4 cms in diameter be cut in a tennis ball, and the ball filled with water and gripped by the hand. It requires little force to expel the water. Repeat the experiment with a ball in which the hole is reduced to 2 cms diameter and the result is little different: the contents are still easily and quickly ejected. Yet this exemplifies the effect of reducing a valve aperture, such as the mitral, to one fourth its original area. A very small rise in driving pressure overcomes the increased resistance offered by a considerably narrowed orifice. It is only when narrowing is very conspicuous that the work of the corresponding chamber is materially increased. The heaviest burden imposed by a valvular defect is probably that imposed by aortic regurgitation, the extent of this is not known exactly, but it is known that the normal heart is very far from finding this burden intolerable. Thus, suppose that the burden imposed by a free aortic reflux actually doubles the work that the left ventricle must do at each beat: when the body is under resting conditions, the ventricle expelling its normal quantity and also what has regurgitated, even so the reserves of the heart are little changed, since the work the heart is capable of doing is very many times greater than that required of it under resting conditions (Fig 32, page 156). If the load it imposes has been fairly estimated, free aortic regurgitation robs the healthy ventricle of only a small fraction of its full reserve. The difference in the

amounts of work that must be done in the presence or absence of free aortic reflux is probably much less than the difference in the amounts that the heart must do when a man is engaged in heavy manual work on the one hand, or in sedentary work on the other.

When the significance of valve disease is regarded from the standpoint of the burden imposed on the heart, the question must become almost purely one of estimating the remaining reserve of the heart. There is no evidence to show that any valve defect imposes a burden upon the healthy heart which, even in the absence of intervening hypertrophy, will result in the heart's failure. If we knew the original capacity of the heart for work and could precisely estimate the burden we might assess the

traction. In the case of aortic regurgitation, it is rarely necessary to estimate the burden imposed by a valve defect, it is certainly undesirable to approach prognosis from a standpoint so indirect, so theoretical and therefore so uncertain.

The only sound basis of prognosis is actual experience, and theoretical considerations should never be allowed to weigh against this. But thought must be clear. The simple fact that there is a heavy mortality amongst patients who suffer from aortic valve disease helps to

as to their prognosis. In a given grade of disease, we are on the point of death or may live a long and active life, reach an advanced age and die of intercurrent malady: how can we proceed satisfactorily to use valve disease in prognosis in the face of such contrasts? It is true that such contrasts are less frequent amongst those suffering from advanced than amongst those suffering from slighter grades of aortic mischief, but this is not altogether relevant, and it is also true that, with the exception of the aortic valve, the grade of lesion cannot be estimated accurately during life. The cases with extremely bad and extremely good prospects are there, and between the two is every transitional type. It should be obvious that if the future is to be foretold in cases of valve disease with any approach to accuracy we must look beyond the valves. Help is obtained in prognosis from the condition of the valves in some cases, in other cases not. If a patient is diagnosed to be suffering from disease of valves and is also recognised to have failure with congestion, the first is so outweighed by the second that it can usually be neglected altogether. A similar statement applies to valve cases in which there

■ great enlargement of the heart and the cardiac reserves are reduced. Failure and enlargement will be the factors determining a reasonable prophecy in these cases.

Just in so far as symptoms and signs of failure, and of cardiac enlargement, are displayed, so the value of the valve lesion recedes into the background.

The presence of a valve defect is chiefly significant in patients who present few or no other signs of heart disease. Amongst patients who have outlived for some years the period of responsible infection, there are some who display aortic regurgitation or mitral stenosis, but in whom bodily vigour is unimpaired and the cardiac reserves are seemingly intact, in whom there is no complaint of symptoms, in whom no definite signs of cardiac enlargement can be found. They are usually, but not exclusively, young subjects and experience shows that in them the prognosis is in general very good, that is to say, these people will usually live many years, and enjoy good health. In this statement lies the basis of correct prognosis in so far as the valve disease is concerned, but it must here be emphasised that in speaking of valve disease I am speaking purely of aortic disease or of mitral stenosis and am not taking into account any of the common accompaniments of these diseases. A few in this uncomplicated group of aortic disease and mitral stenosis and a few only, will develop graver manifestations in a few years. It is most important, however, that this fact should not be allowed to introduce gloom into the prognosis of the group as a whole. It is right and proper that the general attitude to the group considered should be one of almost unqualified optimism provided that the subjects individually realise the wisdom of simple precautionary measures. In this group there is rarely the means of discriminating between the many who will make good and the few who will fail, therefore the attitude to be consistent must be optimistic to all or pessimistic to all. If it is optimistic to all, it may be that an occasional patient will fail without much warning and perhaps dramatically, but the majority of the patients will enjoy what they have a right to enjoy, namely, a reasonable sense of security. If it is to be pessimistic to all, most of them will needlessly experience periods of apprehension or alarm and a constant feeling of insecurity. Surely there can be but one decision in considering these alternatives, for the first only is reasonable, or indeed humane.

If what has been said has been understood, then it is unnecessary to proceed to discuss the case of valve disease that presents minor

degrees of failure or of cardiac enlargement. The general arguments that have been advanced, however, may be given further point by dealing with the example of valve defect that has been the chief focus of recent discussion, namely, mitral regurgitation. This will be discussed before summing up.

Mitral Regurgitation—The difficulties inherent in the diagnosis of mitral regurgitation have been such that the diagnosis is insecure, and its degree cannot be ascertained. When made, the diagnosis hangs exclusively upon one sign, there is no advantage in discussing the significance of mitral regurgitation in any particular circumstances. Therefore, when this murmur is heard by certain

physicians, such as length, harshness and conduction, when considering a murmur that has just appeared (see page 139) in the first fifty who do not, there is no doubt that the prognosis in the first group will be less satisfactory than in the second. But it is beside the point so to discuss the matter, for the second will not form a justifiable control group to the first. It is quite essential, if the significance of the apical systolic murmur is to be ascertained, that the two series of cases should be precisely alike in every respect apart from the murmur. The second group of subjects presents a problem. It never has been

made, and when made, no difference is noticeable. Reliable method of examination is almost always evaded. Thus, if it is stated that mitral regurgitation has little or no prognostic significance, the question is not infrequently asked if it has been associated with ventricular failure. The reply is

The already mentioned of extrasystole. The principle at issue, however, is so vital to prognosis in general, and is so

generally neglected, that this direct emphasis is necessary. An example from a different field, even though it is regarded as extreme, may serve to establish the point that associated circumstances are really irrelevant. A simple healed scar on the tongue clearly has little or no prognostic significance, yet patients who display such scars have not a normal outlook, since some of them are epileptic. The fact does not alter the prognostic significance of the scar itself.

The prognosis in cases of aortic regurgitation has often been stated to be worse if mitral regurgitation is coexistent. This statement is based, not upon experience, but again upon purely theoretical considerations. It is one of the last steps leading to the absurd culminating point, where prognosis is based upon a count of murmurs. In a patient who has aortic disease, to discuss the meaning of a systolic apical murmur is to discuss a triviality and to miss the vital considerations.

To sum up, if it is stated that a diagnosis of aortic valve disease or of mitral stenosis has value in prognosis, and that the diagnosis of mitral regurgitation has little or none, we shall not be far from truth. It has been manifest to me for many years, in watching the work of other medical men, that those who constantly rely upon cardiac murmurs or upon the diagnosis of valve disease are always in difficulty when it comes to prognosis and treatment, and that those who deal most efficiently with cardiac patients lay the chief emphasis upon other phenomena.

This discussion of the significance of valve disease opened with a consideration of the burdens imposed on the heart by valve defects. This led to the statement that prognosis must be based upon experience and not upon the theoretical effects of valve lesions, and it has been concluded that lesions of the aortic valve and the lesion mitral stenosis alone possess material prognostic significance. If we consider why the other

ser-
are

why valve disease is significant is not to be found mainly in the burdens which it imposes upon the muscle, but mainly in associated disturbances that come with the valve disease and from a common source. The heart displaying disease of its valves has usually been the seat of inflammatory process or of degenerative change, disease in the valve is rarely isolated, it is accompanied by disease in other parts of the heart. Thus it is legitimate to regard mitral

stenosis or aortic disease in young adults as furnishing evidence of a rheumatic inflammation of the heart past or present an inflammation affecting all parts of the heart in greater or less degree. An apical systolic murmur even when interpreted as meaning mitral regurgitation has not the same significance. A heart ascertained to present mitral stenosis is one of which the muscle should be suspected before the examination has proceeded farther such patients are more prone than normal people to develop fibrillation of the auricles and failure with congestion. Similarly in aortic disease of syphilitic origin the aorta and parts of the ventricles have been exposed to spirochaetal infection the disease is progressive and there is considerable possibility of damage to the mouths of the cor-

A diagnosis of aortic regurgitation

is added significance by extension

structures. Properly speaking the

the prognosis of the valve disease is not but they are certainly relevant to the question of the significance of valve disease in the more general sense. It will be no great exaggeration to state that the greatest significance attaching to a diagnosis of valve disease in the light it often throws upon the history of the heart affected by it and the help it gives in leading up to a diagnosis of decreased reserve of active infection of disease of the aortic wall or of coronary obstruction.

MANAGEMENT

When active disease is present in the valves the treatment concerns the infection this is dealt with in so far as it is understood under the appropriate headings of infective endocarditis (pages 191 and 193) rheumatic endocarditis (page 201) and syphilitic aortitis (page 225). This really ends what has to be said of the treatment of valve disease since chronic and inactive lesions can hardly be regarded as susceptible to treatment. Surgical attempts to relieve cases of mitral stenosis presenting failure by cutting the valve have so far failed to give benefit. I think they will continue to fail not only because the interference is too drastic but because the attempt is based upon what usually at all events is an erroneous idea namely that the valve is the chief source of trouble.

The treatment of patients displaying valve defects will be found in other parts of this book the treatment is not of the valve defect but of its common accompaniments failure angina etc. The presence of a valve defect should modify treatment rarely. I frequently

when it has been allowed to do so this attitude has been unwise. Thus digitalis was at one time forbidden in cases of aortic regurgitation, under the plausible theoretical plea that it increased regurgitation by lengthening diastole and raising arterial pressure, but the moment the facts were investigated it was found that digitalis in clinical doses produces neither of these effects in the aortic case. The proper use of digitalis is governed by quite different considerations, such as the presence of auricular fibrillation, and these are not different in aortic and other cases. The valve deficiency may be said itself to affect treatment chiefly when the deficiency is uncomplicated. If a patient has aortic regurgitation, aortic stenosis, or mitral stenosis, but presents no sign of cardiac enlargement, and has a good exercise tolerance, it is a wise precaution to forbid any very strenuous act of exercise or long hours of manual work. To confine such cases to purely sedentary occupations and to warn them against all acts of exercise, even to forbid quick walking, cycling, swimming, is not to treat them well. On the contrary, exercise well within the limits of tolerance is to be encouraged, only a limitation of bodily activity is to be insisted upon. The same patients, having recognised cardiac weakness, should be acquainted with and be asked to conform rigidly to simple rules of health (page 283).

The restrictions will naturally be increased to suit the needs of patients in whom the valve defect is complicated by associated conditions.

When a patient presents what is regarded as uncomplicated mitral regurgitation, or, more strictly speaking, exhibits a constant systolic apical murmur, the heart being of normal size and tolerance of exercise perfect and there being no reason to suspect infection, it is my habit to leave bodily activity quite unrestricted and to encourage the full continuance of all pursuits, however vigorous these may be, to pass the patient as a recruit for military service, and to advise ordinary premiums for purposes of insurance. This policy is not settled by any theoretical consideration, but its soundness has been proved by practical experience. It naturally carries with it the proviso that he who adopts it must neither carelessly assess exercise tolerance, nor neglect to examine the heart scrupulously from other points of view. To illustrate this much contended point, there is the case of a young man, seemingly possessing the perfect health and exercise capacity of an athlete. He was sent to see me in 1922 because he had been found to have a blowing systolic murmur near the heart's apex conducted out toward the axilla. Such a murmur was indeed

DISEASES OF THE HEART

CH. VI

present but all other signs were quite natural. He wished to climb Mount Everest and with my consent did so, going as high as any living man has climbed (over 27 000 feet) and returning and remaining in good health. This emphatic example is exceptional only in the unusual severity of the test imposed.

CHAPTER XVI

HEART STRAIN, WORK, AND FAILURE

HEART STRAIN

It has long been thought that the healthy heart frequently suffers strain as an immediate result of strenuous or prolonged bodily exercise. The underlying idea seems to be that the heart may be injured by strenuous effort or by prolonged and severe exercise of the body, and this idea conforms with, and forms part of, the more general and long held conception that heart failure results mainly from the heart being overburdened by work.

THE BURDEN

In discussing heart strain it is not always remembered that the strain of weight lifting or of other strenuous act never falls directly upon the heart, it falls immediately upon muscle, tendon, and bone, the load on the heart is increased purely through the blood pressure. The muscular effort may be abrupt and violent, the load falls on the heart more gradually and less severely as the circulation responds to a call set in motion through normal physiological channels. The chief sources of increased energy expenditure by the heart in exercise come from increased rate of beating, increased output, and raised blood pressure.

ACUTE STRAIN DILATATION

When after some unusually strenuous or violent effort the muscle of a limb at once gives pain and is locally tender, but shows no

fact and overlooks what has just been stated, namely, that the load placed on the muscle of the heart is purely a blood pressure load, and unlike the strain that a heavy weight places directly upon the somatic muscle.

A more commonly held view is that the overloaded heart dilates. The most striking instance of what is to be interpreted as acute dilatation of the ventricle under a suddenly increased load is found in the occasional consequences of rupture of an aortic valve cusp. The rupture happens usually during bodily effort, while blood pressure is raised and the subject may become breathless, cyanotic, and perhaps unconscious. There is no reason to doubt that in these circumstances the heart has become dilated and embarrassed, but the precise circumstances should be noted. It may be assumed that the rupture occurs in early diastole, for that is the time at which the valves will be in greatest tension, at this phase of the cycle the aorta is full of blood and the pressure within it, as a consequence of bodily effort is unusually high. The regurgitant stream of blood in these circumstances must be large and is directed into a rapidly relaxing or relaxed ventricle. The ventricle is submitted to a great reflux abruptly without warning of any sort, and in a moment during which it is powerless to react or adjust itself. The production of an acute overfilling in such circumstances is not difficult to understand especially if we assume, as we are almost certainly justified in doing, that the heart experiencing this accident is showing this response is generally, if not always.

In such cases, here the heart is encountered by the ventricle during its systole, these are quite insufficient to embarrass a healthy ventricle, which, however, will dilate in diastole to accommodate an increasing input of blood at a very low pressure from the veins. The increase of input is met by increase of output of the heart. It is now generally held that the heart is not so easily embarrassed as was formerly supposed. During strenuous effort or exercise, the heart has not been obtained. Since great dilatation of the heart during exercise would certainly have been detected in the many fluoroscopic examinations which have been undertaken we may be content to assume that such dilatation does not occur. It has been shown to increase the efficiency of the organ as a pump. Thus dilatation in exercise

would not be regarded by contemporary physiology as injurious but as distinctly advantageous

Moreover, we know that injurious dilatation, if such is ever threatened, is safeguarded by the parietal pericardium, a comparatively unyielding membrane limiting the capacity of the heart to dilate excessively in health. There are abundant examples in which the healthy, or comparatively healthy, heart becomes greatly distended for shorter or longer periods without undesirable after effects occurring. A simple instance is where the heart is dilated by inspiring strongly with the glottis closed. If this inspiration is maintained, the heart on the X ray screen is seen to dilate very greatly. Yet this experiment can be performed again and again by healthy men with impunity. In paroxysms of tachycardia, as has been described, the patient's heart may become so dilated that the impulse passes well out into the axilla, and the veins and liver may engorge acutely. The attack, in which the heart is beating perhaps at 200 beats per minute, continues thus for many hours or days. Yet, when the paroxysm ends the heart resumes its original size, often after only a few beats and within a short time the patient is as he was before the attack. Such paroxysms, even if repeated, impose no recognisable injury on the heart yet the dilatation is greater, and far more prolonged, than any that can reasonably be suspected from present evidence to occur in effort or prolonged exercise.

The idea that the heart may be strained by a physiological act has received a good deal of stimulus from the incorrect interpretation of certain instances of acute dilatation of the heart, attributed to effort, or occurring during pregnancy. All the cases of this kind that I have been called to see, and in which there were clear signs of dilatation, have been instances of unrecognised paroxysms of tachycardia, the mistake, however, is less frequent than formerly. In the interpretation of these cases two erroneous ideas came into play. The first was that strain can cause pathological dilatation. The second was that dilatation of the heart causes tachycardia, actually dilatation of the heart has little influence upon its rate. In these cases the tachycardia is primary and pathological and it causes the dilatation (pages 72 and 75).

Paroxysms of tachycardia in certain patients are provoked by exercise. In such a patient the first attack complained of may have been determined by some act of exercise. There has been a strong, perhaps a natural, tendency to ascribe the attack altogether to the

act of exercise in this circumstance. But close enquiry shows that this interpretation is unjustified, for the same patients are found previously to have suffered, or subsequently to suffer, attacks that are unprovoked by exercise or by other determinable cause. And close questioning usually reveals that the act of exercise originally blamed, even though it was particularly vigorous, had been undertaken several or many times before without any adverse effect.

It is reasonable to conclude that the burdens imposed by physiological acts upon the normal heart, however heavy these burdens may be, never injure the heart's fibres, never produce injurious dilatation, and never exhaust the heart's reserve.

PROLONGED OVERWORK AND HEART FAILURE

It has been thought that if the healthy heart is worked excessively for prolonged periods, or if it is repeatedly called upon to do strenuous work, that it suffers, becomes diseased, and fails. Heavy manual work has been regarded, with too little thought for its frequent association with exposure, alcoholism, and infection, to be a main cause of cardiac failure, but acceptable evidence has never been advanced for this view. It seems strange, if long-continued heavy work can damage the heart, that in horses, the most heavily strained beasts of burden, chronic heart maladies are almost unknown. The occasional racing horse or greyhound should not have been cited in the past. Because a given hound won many races and ultimately developed heart disease and died, it is unreasonable to believe that the disease was the result of racing. It is known that an increase in the amount of its work causes hypertrophy of muscle, whether somatic or cardiac. Such hypertrophy does not constitute disease, it is physiological. Our evidence is more than a slight physical

real ground for the belief that such hypertrophy is a pathological condition. In fact, the hypertrophy, even in the case of long-distance runners. If racing dogs or race horses or racing men showed a much heavier mortality from heart disease than those that did not race, but were under conditions otherwise similar, this evidence would be suggestive. Actually, it has not been shown that there is an increased mortality from heart disease in any one of these instances. In the case of boat-racing it has been proved that the expectation of life is unchanged, and that the subsequent mortality from heart disease is no greater than amongst the rest of the population. So it will be wise to assume

that the development of disease in the isolated instance is the result of a special circumstance and not of overwork. To cite the isolated case, without attempting to explain the exemption of the great majority, is to depart from the simple rules relating to cause and effect, a cause, to be acceptable as such, must always produce its effect when it acts in given circumstances.

Coarctation of the aorta (page 263) contradicts the idea that failure easily follows from prolonged overwork. In this congenital condition of obliteration or of great stenosis of the aortic isthmus, the systolic pressure may range about 200 or 250 mm Hg under conditions of bodily rest. Here it might be expected that the early appearance of symptoms and signs of breakdown would be the rule. This is not the case, for such subjects are known to maintain these pressures for very many years and to lead active lives, in which frequently these pressures are further and considerably increased, as also is the pulse rate. If the heart can beat against high pressure in this disease for forty or more years without failing, and this seems often to be the case, then even if failure sometimes comes, we should be rash to assume that excessive work is the simple cause. We may recognise that the process underlying fatigue is cumulative, but there is no physiological sanction for an accumulation requiring years, let alone decades, to arrive at a critical level. Since the heart fails to do to-day the work that it has easily done for years, it is necessary to assume that an irreversible or permanent change has happened meanwhile in the muscle, and that the relatively rapid development of failure when it comes is the result of a change in the muscle that occurs, or occurs largely, in the latest period, and not necessarily, or even probably, as a consequence of the primary disease. This change may be a change usual with advancing years, and one that accounts for the similar long delayed failure in essential hypertension, or it may be associated with intercurrent infections, we do not as yet know with any degree of finality. The example remains a conspicuous instance of the uninterrupted overwork with which a muscle, presumably healthy, can cope effectively for very long periods of time.

The case of the burden imposed by a valve defect has been dealt with briefly already (page 143), and it has been stated that patients suffering from such defects often enjoy long and active lives. Such an example as aortic regurgitation may be used conveniently to illustrate the relation between work and heart failure, according to a conception that is of proved and great practical value. I use the

term work here as previously in its ordinary meaning and not in the precise sense of effective work done. In Fig 32 the full or normal capacity of the heart for work is represented by the height of column A which is divided into two parts to represent the work required of the heart to meet the needs of the body under resting conditions (marked rest) and the reserve. The reserve is deliberately shown to be many times that required under conditions of rest.




FIG 32.

In the second column (B) the heart is supposed to be meeting the needs of vigorous bodily exercise. The work of the heart in this exercise is many times greater than that required of it during rest; a reserve is still shown though it is greatly decreased. A hypothetical example of aortic regurgitation, the muscle of the heart being perfectly healthy, is represented in column C and the column is divided into rest factor, aortic factor, and reserve. The aortic factor is arbitrarily represented as equal to the rest factor, the work required of the heart when the body is at rest being no more than doubled in such a case; the reserve is little diminished and the heart may be represented as still just capable of meeting everything that is required of it in vigorous exercise (column D). Column D represents the very rare case of

aortic regurgitation possessed of almost perfect exercise tolerance. In columns E and F the capacity of the heart for work is represented as reduced while there is an abundant reserve under resting conditions; this reserve is inadequate to meet the requirements of vigorous exercise and exercise tolerance is diminished. The capacity of the heart for work is represented as still more deficient in column G, the rest and aortic factors covering a considerable proportion of the whole column and the reserve being greatly diminished. The case is one in which only small calls for extra work can be answered and

exercise tolerance is much reduced. Lastly, in column H, the capacity of the heart is represented as so much reduced that this organ is incapable of meeting more than the requirements at rest and the aortic lesion. The reserve has gone, the patient is breathless at rest, failure with congestion is beginning. A main object of this diagram is to emphasise the idea that failure of the heart is not to be regarded as the simple result of increased work, but chiefly of decreased capacity to work. The difference between the cases represented in columns C and H is not a difference in the aortic burden but a difference in the reserves of the heart.

When heart failure develops it does so, certainly in the great majority of instances, because the capacity of the heart for work has decreased. The extra burden of a valve defect is insufficient to embarrass the heart, it is merely sufficient to reduce the reserves. The burden is not in itself lethal even when long continued. But it hastens the onset of failure when the muscle is becoming weaker. Heart failure often occurs, as previously stated, in cases in which there has been no evidence of an unusual burden. The comparative immunity of the heart from failure in coarctation of the aorta, in which resistance to the output of the left ventricle is continuously raised for many years, the similar immunity in cases of pulmonary stenosis, which throws a heavier load on the right ventricle than does mitral stenosis, are striking examples, the immunity is due to the burden arising out of a congenital, and not out of an inflammatory or degenerative cause—the hypertrophied muscle is healthy and its strength therefore unimpaired.

The manner in which failure is related to the work of the heart is not a mere matter of interesting discussion. There is no branch of medical practice that requires more than that which deals with heart disease a simple and clear philosophy to guide it. No doubt now remains that the treatment of the cases usually called heart strain, and considered in the next chapter, has been gravely mishandled in the past, and that a large number of people who have been thought to suffer from this condition have been reduced, quite unnecessarily, to a state of partial invalidism. It is to the older philosophy, which so greatly over-emphasised the factor of strain or of mechanical defect as a cause of heart failure, that that unfortunate result is to be ascribed.

CHAPTER XVII

EFFORT SYNDROME BREATHLESSNESS OF EFFORT

EFFORT SYNDROME (ATHLETE'S HEART SOLDIER'S HEART)

WHEN a healthy man is taking exercise that is sufficiently strenuous he experiences certain symptoms and certain physical signs develop. The most constant symptom is breathlessness which comes during exercise and continues for a variable time afterwards in gradually lessening degree. He may become aware of his heart beat or of giddiness, he may actually faint. After long or very strenuous efforts the tremulousness and weakness of exhaustion are felt and later the lassitude and aches of fatigue. Heart rate and blood pressure are raised during the period of exercise and the rate and depth of breathing are increased, accessory muscles of respiration being brought into action. The cardiac impulse becomes more forcible and is often diffuse, systolic murmurs often occur over the basal and less frequently over the apical regions of the heart. The term physiological syndrome of effort is applied to this group of symptoms and signs.

The term effort syndrome is used to cover a similar series of symptoms and signs that occur in a large class of patients complaining of ill health. From the standpoint of symptomatology the difference between these patients and healthy people consists mainly in the ease with which the symptoms are brought on by exercise though in the patient additional symptoms may appear. In normal subjects it requires more work than is required in patients to produce the same symptoms. The dividing line is not abrupt. Very possibly several distinct but similar syndromes are being confused for the moment useful subdivision cannot be effected. Certainly the syndrome is preceded by different primary conditions but important as it is we are unaware of its cause.

The syndrome is essentially a malady of growing boys or young men but it is not confined to these ages or to the male sex. Occurring

in budding athletes or in young soldiers, it has attracted particular attention, because it renders these quite unfit to follow their ordinary occupations. It is most important to realise that the syndrome is not peculiarly, or even particularly, a soldier's malady, or an athlete's malady, it is one of the commonest chronic affections of sedentary town dwellers. It tends, however, to remain concealed in the latter and to be much less a matter of complaint than in active people. When the young men of a nation are all called upon for military service, then a considerable proportion of those drawn from sedentary occupations are found to be temporarily or permanently unfitted for active work, and the commonest form of unfitness is this malady, which is revealed by the unaccustomed physical drill to which the recruits are subjected.

It is the syndrome that is more commonly mistaken for cardiac disease than any other. Amongst those who display it there is the very definite group of cases suffering from chronic infections—obvious or concealed, the chief being pulmonary tuberculosis, local pus infections of nose, ear, throat, or other part, infections of bowel or bladder. Another large group is that comprising patients who are convalescing from acute infectious diseases such as pneumonia, pleurisy, infections of the gastrointestinal tract, bronchitis, tonsillitis, or influenza. Transient symptoms are the rule rather than the exception in the first days of convalescence, but they may be prolonged for weeks, months, or even years. In most cases, however, no infection is found and there is none in the recent history. A few cases are suspected of hyperthyroidism.

SYMPTOMS

The onset is usually gradual, or so far back as to be forgotten. It may be more abrupt, in which case it has often followed directly upon an infection. When it has begun abruptly, it is stated that acts of work previously performed can no longer be undertaken without abnormal discomfort. Very rarely, the malady is said to have come after an unusually strenuous or prolonged physical effort, the subject believing himself at the time to be in vigorous health. Breathlessness brought on by exercise is an invariable complaint and is the chief symptom, palpitation is very common. Both these symptoms may be provoked by very small amounts of exercise and may be severe in the more serious cases. Dizziness accompanying change of posture or the end of an effort is very common, attacks of fainting of the vasovagal type (page 98) occur in a number of the subjects.

As in the response of the normal to strenuous or prolonged work, tremulousness, weakness lassitude and fatigue are often added even after minor efforts, precordial discomfort, in the form of aching pain, soreness, and tenderness of skin and muscle, is frequent but rarely severe, and is produced or aggravated by exertion

SIGNS

The subjects are usually thin often undersized, or presenting long, narrow or flat chests. A number show signs of nervous instability as evidenced by the quickness of emotional responses and irritability. Coldness of the hands and feet, and free sweating of these and of the axillae is common

The most significant signs are found in the response of respiration and of the cardiovascular system to exercise. At absolute rest in bed and during sleep the respiratory rate is normal, but a simple hopping test (page 4) deepens the excursion and raises the rate, according to the severity of the condition to between 30 and 70 per minute. In normal subjects of the same age the respiratory response to this test is scarcely noticeable. The heart rate in sleep is normal, but it is almost always found raised in subjects that are simply resting, and if they are excited or restless or are up and about, increased pulse rate usually draws attention. A simple hopping test raises the normal heart rate of young people to 90 or 100 per minute and thus subsides in a minute or less. In the milder cases of effort syndrome the rates jump to 120 or 130 and in severer cases to 150 or even 180 per minute and the subsequent return to the pre-exercise rate takes much longer than normal. The systolic blood pressure, while normal at rest is in general higher than is expected in these subjects when up and about and is raised unusually by emotion and by exercise, and falls slowly again as does the pulse.

The main cardiac impulse which is often forcible and quick, lies in its accustomed place, but the impulse as a whole is diffuse, appearing widely in the 3rd 4th and 5th and even in the 2nd and 6th spaces. The cardiorespiratory murmur is very frequent, as are brachial systolic murmurs, systolic murmurs at the impulse appear also but less usually. These abnormalities of the cardiac signs are particularly prominent when the heart is beating rapidly after slight exertion or under mental excitement. The signs are often so striking that it is difficult to persuade those who have no special experience of the cases, or experience of X ray examination in them, that the hearts are undilated.

SIGNIFICANCE OF THE SYNDROME

The meaning of the malady that has been described is none too definite, it is perhaps better that the picture should have been painted without a central motive to fix attention, for if the condition is regarded as a specific form of disease, the right attitude towards the patient is lost. This should usually be one of deferred judgment. Each case requires to be considered on its own merit, a number will at once find their place under well defined diagnostic headings, such as incipient tuberculosis, some will by development enter their well defined categories later while under close observation. Most of the cases, however, will remain under the term "effort syndrome", a term that deliberately signifies a group of symptoms rather than a specific disease and which, by refraining to imply the cause, helps to maintain an attitude of enquiry towards the malady.

The symptoms are essentially provoked by effort, and are provoked very easily in severe cases. It is for this reason, and because certain cardiovascular manifestations are prominent, that the syndrome came to be regarded in young men and soldiers as a disorder of the heart, resulting primarily from the strain of strenuous athletics or the long continued efforts of active military service. The idea of heart strain found support in the diffuse impulse and the occasional systolic murmur, both being regarded as indicating dilatation of the heart. The diffuse impulse is unreliable, and the systolic murmurs in these cases are exocardial more often than not, both diffuse impulse and murmur result simply from overaction of the heart, the former being rendered conspicuous if, as is common, the chest is narrow or flat, and thinly covered.

Another factor that has contributed to the idea that the malady originates in effort is the not infrequent history that breathlessness had first been experienced with unusual exercise. It seems often to be overlooked that the effort has been made many times before without unusual symptoms. If the respiratory reserve is declining, or has declined, the first symptom will naturally appear with the most vigorous effort that is customary, that is the occasion on which the reserve is put to the test, and is found to be deficient. The chief part that effort plays is clear, it calls upon a waning reserve, bringing a previously concealed defect to light. Though many are thought to be of this kind, I have seen no case amongst civilians or soldiers in whom I could regard the evidence for strain as a primary cause of breathlessness as at all convincing.

During the early days of the Great War patients of this type were regarded, according to views then current, as primarily cardiac and were rested, or, when exercised, were exercised with much caution. Experience soon proved any fears groundless, many thousands were ultimately given graduated exercise rising to the full point of toleration, and in many to strenuous exercise. In not a single instance was there suspicion of adverse after effects of this treatment, on the contrary, the exercises came to be used widely as remedial measures.

PROGNOSIS

The prognosis of cases of effort syndrome in which no underlying infection is found is good, though progress may be slow. Within five years, and living under ordinary home conditions, a third of these patients are quite fit or are improved. Most of the remainder remain in an unchanged state. The sickness and mortality rates amongst the group are normal for their surroundings, with the single exception of tuberculosis, which is a little more prevalent than it should be. Heart disease does not appear more frequently than is usual. The younger the patients the greater is the percentage of recovery within a given period.

MANAGEMENT

Cases presenting the effort syndrome should be closely searched for foci of infection when first coming under observation and later, from time to time. Local infection will demand appropriate treatment, but in most cases none will be found. The patients should be persuaded that they have no heart disease, and that the sensations they have attributed to the heart have in reality another source, or are due to temporary and trivial disturbance of the organ consequent upon general lack of tone and fitness. Very many have been perturbed by a cardiac diagnosis and are anxious about the future. Simple reassurance of the patient and his friends often brings about improvement. To diagnose heart strain in these patients and, on this or other ground, to forbid exercise or to enjoin rest, is to cripple these patients and to bring many of them to a state of partial or complete invalidism. The most successful method of treatment is by simple exercises in the open air, for this encourages a more vigorous habit of mind and body. The exercise, whatever form it takes, is to be governed by the patient's responses. In the more severe cases it should be light walking exercise or very simple drilling, in less severe cases it should be moderate, and so forth, the

patient should not exercise or be exercised to the point of distress, but to a point that quickens and deepens respiration appreciably. There are few cases that will not benefit from two periods of exercise, each of half an hour or more, each day. The longer the cases are in the open air the quicker, in general, they improve, and so the hours of work amongst students and those employed in sedentary occupations require to be curtailed, and often may cease with profit for a period of weeks or months. Exercise is pressed as recovery permits. Breathing exercises are useful in expanding the chest and increasing the vital capacity of the lungs, especially in young subjects. The patients should be well fed. Smoking interferes with recovery and should be forbidden or kept to the strict limit of a small smoke after each chief meal. Treatment by drugs, other than occasional tonics or laxatives when required, has little value.

DIFFERENTIATION OF BREATHLESSNESS

The symptom breathlessness on effort is common to heart failure and to the syndrome now described. The recognition of early heart failure, displayed solely as breathlessness, and its differentiation from the effort syndrome depends upon associated signs or symptoms. Thus, in early heart failure breathlessness is usually accompanied by definite signs of cardiac enlargement or valve disease, while in the effort syndrome it is associated with conspicuous overaction of the heart. For the most part the differentiation is a simple matter therefore. But there are instances of real difficulty. The effort syndrome is a very common one, and there is little reason to doubt that it occurs not infrequently in patients who actually display cardiac disease. This is suggested not only on *a priori* grounds, but because breathlessness in some young cardiac cases is out of all proportion to the signs and shows no tendency to progress to congestion. They must be distinguished, so far as possible, since on general principles the treatment of failure consists of rest, and in the effort syndrome exercise is encouraged. The contrast in treatment, however, is not so sharp as this, for limited exercise is beneficial in cases of heart disease in which breathlessness is slight, and exercise should not be allowed to distress patients who are regarded as free from structural disease. In practice, therefore, it means only that the heart case is allowed a good deal less exercise than the other. The general rule should be to regard breathlessness, the cause of which is not obviously attributable to renal disease, bronchitis, infection,

or neurosis, as a manifestation of the effort syndrome, when it occurs in young people who present no signs of structural mischief in the heart, or insufficient signs to be consistent with the symptoms, and to regard breathlessness as signifying early cardiac failure, when it happens in young subjects with well defined signs of heart disease, and in middle aged and elderly subjects, whether clear signs of heart disease are present or not, and to arrange treatment accordingly

Some physical signs that help the identification of cardiac breathlessness and which sometimes stand alone, are described in Chapter XXV

Nervous breathlessness rarely gives rise to difficulty in diagnosis. Apart from the characteristic behaviour of most of those who display it the breathlessness is not specially associated with effort, but rather with emotion or nervousness. It usually increases in degree when attention is drawn to it and may or may not be increased by effort, and notable inconsistencies are presented, thus, the patient will usually hold the breath for ten or more seconds at any time to a sharp request and the breathing will not usually be deeper or faster, but often the reverse, when it is resumed. Nervous breathlessness disappears during sleep.

CHAPTER XVIII

PULMONARY CONGESTION, OEDEMA, AND INFARCTION

CHRONIC PASSIVE CONGESTION AND EARLY OEDEMA OF LUNGS

PASSIVE congestion of the lung is due to increased pressure in its vessels. This can be caused by anything that obstructs the outflow of blood from the lungs. Thus a chronic state of pulmonary congestion results when the pulmonary veins are compressed or thrombosed, or in such narrowing of the mitral orifice as raises the average pressure in the left auricle appreciably. It is the rule to find evidence of pulmonary congestion when the heart as a whole is failing, its association with systemic venous congestion has been noted already. In passive congestion there is not only engorgement of small vessels of the lungs but increased transudation from them, even making the tissues a little, and in more profound forms manifestly, oedematous.

Reduction of alveolar space by engorgement, and by an increase of fluid within alveoli and bronchioles, puts many alveoli out of effective action, the blood passing through the corresponding vessels on into the left heart and systemic arteries without changing its venous state.

When of very long duration, congestion widens the pulmonary arteries, and these often become atheromatous, the tissue of the lungs is engorged and pigmented, the bronchi are a little inflamed, and the connective tissue generally is diffusely increased, a condition termed 'brown induration' of the lung.

SYMPTOMS

Pulmonary congestion causes breathlessness by decreasing the power of the lungs to aerate the blood passing through them. In cardiac failure pulmonary congestion contributes in this way to breathlessness that is occurring at rest. It causes a hard cough with a little sticky and sometimes more watery and slightly blood stained expectoration, according to the amount of fluid transuding

SIGNS

High pressure in the pulmonary artery is often accompanied by a ringing 2nd sound at the pulmonary cartilage. Such accentuation, however, may not be used as certain evidence of such increased pressure (see page 229). Dilatation of the pulmonary artery may be identified by X ray (Fig 30 page 140). Congestion causes or contributes to cyanosis, which in advanced cases is recognisable clinically as of central type (page 36). The breath sounds at the bases of the lungs are weakened, crepitations are heard there first, and over the lungs scattered rhonchi become audible. These adventitious signs increase in extent, and the sputum becomes more abundant and more watery when oedema plays a larger part.

ACUTE CONGESTION AND OEDEMA OF LUNGS

ACUTE CONGESTION WITH EARLY OEDEMA (CARDIAC ASTHMA)

The term cardiac asthma is used to indicate a severe form of breathlessness occurring in the form of paroxysms that are mainly nocturnal. The attacks are observed chiefly in patients between the ages of fifty and seventy and are very rare before the age of forty. Men suffer much more frequently than women. Three fourths of the patients are sufferers from essential hypertension, many of these and many of the remainder, have advanced arteriosclerosis, with or without sufficient involvement of the coronary arteries to yield angina pectoris. Some cases present syphilitic aortitis. A few others show definite evidence of renal insufficiency. There is no distinction between what is called 'cardiac asthma' and what is called 'renal asthma'. The case may show aortic valve disease or mitral stenosis less commonly, in the great majority there is no evidence of valve disease but enlargement of the heart is invariable, and gallop rhythm, alternation of the pulse, or a gross anomaly of the electrocardiogram often bears witness to involvement of cardiac muscle (see Chapter XXV). Cheyne Stokes breathing is a frequent accompaniment.

The Attack —An elderly man, having few symptoms or experiencing a little breathlessness on effort in the daytime, retires to bed and, lying recumbent, sleeps. He wakes in the early hours of morning feeling oppressed and breathless, sits up in bed and pants, becomes restless as breathlessness increases, is distressed

and sits on the edge of the bed or moves to the window, after a few minutes or longer he becomes wheezy and soon the attack subsides

The severer attack wakes the patient and brings him at once to a sitting position with a sense of intense suffocation. The breathing is increased in rate and becomes more and more forcible, he clutches surrounding objects and brings all the accessory muscles into play. But the chest, increasing in size, moves less and less effectively. Distress is terrible. Little air can be taken into the lungs and expiration is prolonged and powerful. Cyanosis comes and quickly deepens, while at first the veins remain unengorged. Pallor may be added. Sweat breaks out and is profuse. The man is brought within a very short time into the throes of a fierce struggle for breath. The chest is filled with sibilant rhonchi and râles. The patient becomes semi-conscious. The attack begins to subside. In this, and sometimes in the milder attack, cough develops and the patient brings up a little frothy and usually blood stained sputum. He is left exhausted. Such a severe attack may last a half hour, an hour or more. It threatens life ending sometimes but not often in fatal oedema of the lungs. In the early stages the pulse is quick (up to 120) and strong, and the systolic blood pressure is found to have risen, it is the rule for pressures to be high in the attack. In the later stages pressure may fall and the pulse become almost imperceptible.

Short attacks may be repeated one or more times in a night. Though usually confined to the night similar attacks also occur occasionally when patients lie down by day.

The attacks are almost certainly due to a rise of pressure in the pulmonary vascular system, producing turgescence and early oedema of the lungs. The manner in which the attack is brought about is unknown.

Diagnosis—The attacks of breathlessness here described are mainly nocturnal. They must be distinguished from attacks brought about differently. Patients suffering from failure with congestion and orthopnoea slip down in bed off the pillows in their restless sleep, after a time they wake breathless and distressed. This form of breathlessness can easily be remedied if recognised. Patients suffering from Cheyne Stokes breathing are sometimes awakened repeatedly at night by particularly long and violent hypernoeic periods.

ACUTE PULMONARY OEDEMA

In acute oedema the lungs are not only congested but at their bases or throughout are heavy and have lost their spongy crepitant character. They are firm, pale and pit on pressure. When the lung is cut a thin blood stained fluid runs from the surface and when it is squeezed this fluid runs freely and is frothy. The same fluid is found in the respiratory passages. It is believed to occur as a consequence of primary failure of the left ventricle. A similar condition of the lungs can be produced experimentally and acutely, by interfering mechanically with the action of the left ventricle, the right continuing to beat efficiently.

In elderly subjects suffering from arteriosclerosis usually accompanied by hypertension in syphilitic aortitis and in occasional cases of mitral stenosis acute and severe pulmonary oedema can occur in single or recurrent attacks. It happens occasionally as an immediate sequel to an attack of angina pectoris. At the beginning of the attack systemic blood pressure has usually been found raised. A very instructive association is the occurrence of pulmonary oedema in the attacks of those rare cases of paroxysmal hypertension which are thought to result from release of natural suprenal secretion (page 60).

The onset may resemble exactly that of the attacks previously described occurring usually at night and waking the patient from sleep, or it may come in the hours of daylight and wakefulness. The separation of these attacks from those previously described is in fact arbitrary and necessitated only by relative dryness of the latter and their very frequent repetition. In frank pulmonary oedema while breathlessness is usually urgent from the start cough is more prominent and frothy blood stained fluid is freely or profusely expectorated. The march of events is often speedy. Cyanosis deepens rapidly breathing becomes progressively more urgent and less effective. The attack may last at its height for hours. In fatal attacks the pulse becomes steadily weaker until imperceptible, the veins swell and the breathing becomes more gasping less frequent, and then weaker until it ends.

In a fulminating form the oedema occurs so rapidly and is so intense that within a minute or two of crying out in fear, the patient is drowned by the copious blood stained fluid that pours into the respiratory passages and overflows frothing from the mouth and nose.

In coronary thrombosis a mild form of oedema of the lungs unaccompanied by intense breathlessness or much cyanosis is common. The sputum is scanty and frothy, or consists of a little watery blood stained fluid that continues to be ejected for a few hours or several days.

Symptoms usually slight of pulmonary oedema may also come acutely in attacks of paroxysmal tachycardia that bring about failure with congestion. Here the mechanism is clearly that of passive congestion of the lungs, though coming quickly. Acute oedema may supervene upon chronic passive congestion.

PROGNOSIS

From the milder attacks of nocturnal breathlessness with or without early oedema recovery is invariable. Recovery is the rule in the more severe attacks of manifest oedema of the lungs, but these are more threatening to life and a fatal termination is not very uncommon.

Of patients suffering from repeated nocturnal attacks of breathlessness of the type described, or from less frequent attacks of pulmonary oedema, many are dead within six months and the majority within two years of the first seizure. A few survive longer and occasional cases suffer one or more attacks and surviving experience no more. In most of the patients the attacks are not the only ominous manifestations (see page 215).

TREATMENT

Patients suffering from nocturnal attacks should sleep on high pillows or upon a cardiac bed with well raised back, whether they are orthopnoeic or not. Opiates help to fend off the attacks. A milk and vegetable dietary has been advocated.

In the severe attacks, whether signs of oedema are clear or not

relieve mild attacks occasionally, but in severe attacks are much less effective than is morphia. Atropine ($\frac{1}{15}$ grain) is often given with morphia when oedema is obvious, but its value is not established. Strophanthin and pituitary extract have found their advocates. Adrenaline should on no account be given. In patients in whom the systemic veins are swelling appreciably, bleeding should be undertaken without delay.

ACUTE PULMONARY OEDEMA

In acute oedema the lungs are not only congested but at their bases or throughout are heavy and have lost their spongy crepitant character. They are firm, pale, and pit on pressure. When the lung is cut, a thin blood stained fluid runs from the surface and when it is squeezed this fluid runs freely and is frothy. The same fluid is found in the respiratory passages. It is believed to occur as a consequence of primary failure of the left ventricle. A similar condition of the lungs can be produced experimentally and acutely by interfering mechanically with the action of the left ventricle, the right continuing to beat efficiently.

In elderly subjects suffering from arteriosclerosis usually accompanied by hypertension in syphilitic aortitis and in occasional cases of mitral stenosis acute and severe pulmonary oedema can occur in single or recurrent attacks. It happens occasionally as an immediate sequel to an attack of angina pectoris. At the beginning of the attack systemic blood pressure has usually been found raised. A very instructive association is the occurrence of pulmonary oedema in the attacks of those rare cases of paroxysmal hypertension which are thought to result from release of natural supra-renal secretion (page 60).

The onset may resemble exactly that of the attacks previously described, occurring usually at night and waking the patient from sleep, or it may come in the hours of daylight and wakefulness. The separation of these attacks from those previously described is in fact arbitrary and necessitated only by relative dryness of the latter and their very frequent repetition. In frank pulmonary oedema while breathlessness is usually urgent from the start, cough is more prominent and frothy blood stained fluid is freely or profusely expectorated. The march of events is often speedy. Cyanosis deepens rapidly, breathing becomes progressively more urgent and less effective. The attack may last at its height for hours. In fatal attacks the pulse becomes steadily weaker until imperceptible, the veins swell and the breathing becomes more gasping, less frequent and then weaker until it ends.

In a fulminating form the oedema occurs so rapidly and is so intense that within a minute or two of crying out in fear the patient is drowned by the copious blood stained fluid that pours into the respiratory passages and overflows frothing from the mouth and nose.

As a rule few signs, other than an occasional rhonchus can be found. In cases where the area infarcted is larger and suitably placed, signs of the consolidation may be detected.

In extensive infarction from numerous or very large emboli, breathlessness may be severe, and signs of failure appear and progress, or oedema of the lungs may supervene and kill the patient. The accident naturally has more serious effects in patients already presenting signs of failure. In most patients recovery is speedy and uneventful.

Massive infarction of an apex, which is occasional, may be confused with pulmonary tuberculosis.

TREATMENT

This consists usually in maintaining the patient at complete rest, with or without morphia, and, in cases of auricular fibrillation, in bringing the heart under the influence of digitalis, without inducing vomiting. In the exceptional severe case, the treatment is that of cardiac failure.

HAEMOPTYSIS

A spitting of blood is not an infrequent complaint of the subjects of heart disease. In these it is first wise to exclude haemorrhage from infection of the gums or from the nose. Haemoptysis is marked by cough and haematemesis by vomiting, but the history is not always reliable in this respect, when blood comes in any quantity through the trachea. Smaller quantities are coughed subsequently. Brisk, fatal or non fatal, haemoptysis may happen occasionally when an aneurysm ulcerates trachea, bronchus or lung. Small quantities of pure blood bright or dark may be brought up from an aneurysm oozing in the same situations. Infarction yields small dark masses of almost pure blood having a foundation of mucus. In acute bronchitis or tracheitis expectorated mucus is streaked with blood. Congestion of the lungs stains the sputum intimately and may yield almost pure blood. In oedema the blood stains a serous fluid.

CHAPTER VII

PERICARDITIS

ACUTE PERICARDITIS AND PERICARDIAL EFFUSION

PATHOLOGY

PERICARDITIS arises in a variety of ways. The commonest is for it to display itself as a chief manifestation of an infective process. There are very many examples but the most frequent are rheumatism and pneumonia of these the former and commonest is a subacute and the latter an acute process. Tuberculosis causes chronic pericarditis. Another way is for pericarditis to appear as a terminal event in a chronic disease that affects nutrition of which Bright's disease is the most notable example. Pericarditis is not infrequently caused by a direct spread of infection from neighbouring suppuration—for example from empyema or ulcerated oesophagus—or by pus tracking through the diaphragm from appendix stomach or gall bladder. It may follow direct wounds of the pericardium. It occurs quite locally when the pericardium is involved as a sequel to coronary thrombosis.

In acute inflammation the pericardial membrane loses lustre, becomes injected, swollen and sometimes gelatinous. On its surface lymph collects in flakes and loosely attached tags. The lymph may be in small amount or a large quantity collects covering the surface with a thick irregular rough or shaggy sheet of yellow deposit. Such is the pericarditis of rheumatism and of Bright's disease. Some excess of fluid is nearly always present. There may be much fluid this is serofibrinous in rheumatism, purulent in pneumonia and in pericarditis by extension, it is serofibrinous hæmorrhagic or caseous in tuberculous disease. Very large collections of fluid are usually chronic, often tuberculous, more often of unknown origin.

The parietal pericardium is regarded as being a relatively inextensible membrane. This is true in the case of tension that is of brief

duration, and the pericardium thus serves to guard the heart from injurious over distension, such as would occur, for example, in asphyxia. But the pericardium yields readily to chronic distension, whether this is brought about by an enlarged heart or by effusion. The heart lies between the bodies of the vertebra and the sternum, and there are no intervening structures that are compressible, distension of the pericardial sac occurs therefore first to left and right, but especially to left, and from the left it distends backwards, pressing upon and causing collapse of the basal part of the left lung.

SYMPTOMS

Pericarditis is usually quite painless. Pain, when present, is probably due usually to involvement of extrapericardial tissues, such as the diaphragm. It may be a continuous severe precordial ache, or a stabbing pain reaching the abdomen or the left shoulder, and felt especially on inspiration. Breathlessness resulting from obstruction of venous return or from pressure on a respiratory organ by large effusions is sometimes to be counted a true symptom with other occasional pressure symptoms, but it is hardly possible in most cases to separate such from associated symptoms of cardiac failure. The remaining symptoms result from the causal infection or accompanying disease, and these vary greatly in kind and in severity.

SIGNS

When pericarditis comes as an acute infection, the dominant signs may be those of intoxication. Suppurative pericarditis gives high fever and pulse rate, breathlessness, often great restlessness, and sometimes delirium. As a part of a subacute rheumatism the toxæmia is less intense, though fever is the rule, restlessness is common, and increased pulse rate invariable.

Cardiac Impulse—In effusion the heart usually maintains its contact with the chest wall, but the heart's impulse often changes nevertheless. It becomes diffuse and wavy. It may be displaced upward to the 4th space or outward. Occasionally it disappears, and the heart sounds become more distant. Such changes are not always the simple result of pericarditis, there may be simultaneous dilatation of the heart. In children the precordial intercostal spaces may flatten.

Distension of Veins—The heart is very sensitive to intrapericardial pressure which, like intrapleural pressure, is normally below atmospheric pressure. A small rise of pressure in the pericardial sac at once interferes with the filling of the heart and causes a fall of

arterial and a rise of venous pressure. Any considerable increase in intrapericardial tension greatly impedes the venous return and the veins become distended. Distension of the veins is frequent in pericarditis with effusion but can rarely be shown to arise in the manner described that is so because pericarditis is so often accompanied by changes in the heart that lead to failure with congestion.

Cardiac Dulness—In considerable effusion cardiac dulness extends up into the 2nd and then into the 1st interspace. This new area of dulness may spread as much as 2 inches to the left of the sternal margin and to the right may overlap the sternum. It is the most important sign of fluid that we possess; it is best displayed by patients in the recumbent posture, and may lessen or even disappear in the erect posture.

Cardiac dulness also increases laterally, especially at the level of the 4th and 5th spaces and both to left and right, the latter is the more significant. To the left, dulness may appear well beyond the outer margin of the impulse. The contour of the heart's dulness in large effusion is that of a blunt cone, right and left margin running obliquely but unusually straight, or it is pear shaped.

Friction Sounds are produced by the rubbing together of visceral and parietal membranes that have lost their polish or are actually rough. The sound often simulates a blowing murmur, but is usually shorter, seems more superficial, and ends more abruptly than the latter, usually, too, the sound has a characteristic scraping quality. There may be a single systolic sound, and this usually begins a little later than systole, but it is the rule to hear a diastolic element as well, to and fro sounds of equal intensity and duration. The to and fro sounds do not change in relative intensity from place to place as do the conducted to and fro murmurs of aortic disease. Very occasionally a third friction sound is added, to make a sort of canter rhythm of friction. Creaking or leathery sounds are unusual. Pericardial friction limits itself strictly to the precordium, may occur over the whole of it, but is more often confined to the central parts and base, it may be quite local. It is seldom palpable.

The friction rub is the only sign we have of dry pericarditis. It may also result from chronic roughening of the pericardium, and so the advent of friction sounds, or change in their character or extent from day to day is more important, in fact alone conclusive of active mischief. Although friction sounds often change in character, or disappear as fluid collects, the presence of a rub does not exclude effusion, or even large effusion, for the heart is often maintained in

contact with the anterior parietal pericardium in these circumstances

Friction sounds having a cardiac rhythm may also be heard when there is inflammation of the pleura overlapping the heart, such friction has a respiratory rhythm as well and is called "pleuro-pericardial friction" It is heard along the right or left border of cardiac dulness

Basal Signs —In large effusions, the lung becomes compressed at the left base. An area of dulness appears, extending from the spine as far as the mid scapular line or a little beyond it, and upwards as far as the angle of the scapula. This rather rectangular form is characteristic, as are the signs which accompany it. The breath sounds at the base are weak or absent, but just below the angle of the scapula loud tubular breath sounds and aegophony or bronchophony are heard, often accompanied by crepitations. When these signs are early they will sometimes disappear if the patient leans far forward. In recurrent effusion the signs may vanish and return several times. In rheumatic children they are said to accompany dilatation of the heart without effusion.

X ray Signs —The shadow of the heart is widened laterally, the basal part of the shadow is increased in breadth and density, and its upper left margin is much more oblique than normal. In the recumbent position, these signs become less conspicuous in the upright position. Pulsation of the margin of the silhouette is reduced.

Diagnosis of Effusion —To differentiate between acute dilatation and pericardial effusion in rheumatic children is often impossible, especially since the two conditions often coexist. The most important guide to effusion is upward extension of dulness. In adults small pericardial effusions are rarely diagnosed. When large, the characteristic area of dulness and the signs at the base are usually sufficiently distinctive. Friction sounds or the loss of cardiac impulse may draw attention to and X ray examination will help to identify, effusion. The extension of cardiac dulness well beyond the impulse is a valuable sign in favour of effusion. When dulness to the right of the sternum is extensive, this may result from a greatly enlarged left auricle (page 112), in that case heart sounds will generally be loud over it.

COURSE AND TREATMENT

Acute suppurative pericarditis, whether it occurs as a primary or secondary manifestation of infection, usually runs a rapid and fatal

constrict or block the mouths of entering veins, it forms a "constrictive pericarditis"

The inflammatory process may involve surrounding structures, and in healing bind the heart firmly to the chest wall and to the diaphragm. The lateral parts of the pericardium, with the adjoining layers of the pleura, become welded into a sheet of dense tissue, which then forms a firm partition between heart and lung, stretching from the root of the latter to the diaphragm. The same process invades the roots of the lungs, the base of the heart and its outgoing and incoming vessels, and the posterior mediastinum, further anchoring the heart, and sometimes constricting the main vessels. This condition is called "mediastinopericarditis"

SYMPTOMS AND SIGNS OF AN OBLITERATED SAC

The smooth, glistening pericardial membrane, lubricated by fluid, normally allows the heart to move within the sac with little friction. Yet this smooth sac is not indispensable, as is shown by its occasional congenital absence, out of which no trouble seems to come. Obliteration of the sac, as after rheumatic infection, does not increase the work or weight of the heart appreciably. It gives rise to no symptoms. The intrapericardial adhesions may be responsible occasionally for a friction rub or cardiac crepitus, but they cannot be diagnosed, unless on the basis of a remembered acute inflammation of the sac.

SYMPTOMS AND SIGNS OF MEDIASTINOPERICARDITIS AND OF CONSTRICTIVE PERICARDITIS

The symptoms of great thickening of the pericardium and of extrapericardial adhesions are mainly those usually associated with venous congestion. The signs that may occur are many.

Systolic Retractions—During systole the heart becomes smaller and tends to create a vacuum in its vicinity. In front this pull is exerted, especially by the right ventricle, in the region of the 4th and 5th left rib spaces and the epigastrium. A heart of normal size, beating quietly and covered naturally by lung, pulls sufficiently to cause a little recession in the epigastrium only, and that not always. An enlarged and powerfully beating heart often causes systolic recession of at least the rib spaces too. In mediastinopericarditis these recessions may be much more emphatic, and not only rib spaces but the ribs themselves and the sternum may be retracted. The

structures most affected in this way are the 5th and 6th rib cartilages and the xiphisternum, but retraction may extend into the axilla and involve lower ribs than those named. Through the diaphragm, systolic retraction of the lowest parts of the bony thorax may occur, seen at its left margin when the chest is viewed from the front, or at the left base when the chest is viewed laterally.

It is not quite correct to regard the heart as tugging the precordial ribs inwards through firm adhesions between it and the chest wall. When the heart tends to draw away from the chest wall, the latter must inevitably follow it, whether bound by fibrous tissue or not, provided that the lungs cannot move in to fill the gap. Ordinarily, decrease in the size of the heart in systole is balanced by an equivalent expansion of the air-containing lungs, which bound it laterally, this systolic expansion of the lungs calls for the expenditure of little cardiac energy. When inflexible fibrous tissue in pericardium and pleura separates the heart from the lungs, the heart expends energy in pulling in all directions upon relatively unyielding structures. Thus, fibrous tissue covering the heart laterally impedes its beat more than do any adhesions anteriorly, for the former alone guards a yielding structure, and these lateral adhesions come to be more important in bringing about precordial retraction than do adhesions binding heart and chest wall directly. Some energy may also be expended uselessly in direct pulls by the heart through fibrous bands and diaphragm upon more distant structures.

Retraction of the more rigid parts of the chest wall will naturally happen more readily when these are young and pliant, and will be greater the more vigorous the movement of the ventricle. Thus, although systolic retractions, and especially emphatic retractions, are valuable, they are inconclusive signs, there is no known form of retraction that is not to be observed on occasion apart from adhesions, moreover, advanced mediastinopericarditis with strong adhesions between heart and precordial chest wall may be present without any retractions occurring.

Immobile Impulse—Normally the impulse and left margin of cardiac dullness move 2 to 4 cms ($1\frac{1}{2}$ inches) to the left when a young subject rotates from the supine to the full left lateral posture, and equally to the right when the full right lateral posture is assumed. Adherency of the heart to the chest wall, or its enclosure within firm partitions of fibrous tissue, naturally renders it immobile. However, the positive use of this sign and of corresponding

signs obtained by X ray screening, is lessened in value by the fact that an enlarged heart is rarely very mobile

Distension of Veins—When pericarditis and mediastinitis are of rheumatic origin, the heart has always suffered simultaneously, and thus venous congestion may be the result of muscle failure. But congestion of the veins may also occur in these conditions because the return of venous blood to the heart is impeded. This may happen through constriction of one or other of the great veins or by the heart being held more or less encased by unyielding bands of fibrous, or occasionally calcified, tissue. Such cases are uncommon, but venous congestion so produced has its peculiarities. These are due perhaps to the output of the heart being less affected or to smaller disturbance of the pulmonary circulation when the pulmonary veins escape constriction, than happens in failure. Be that as it may, patients of this class may be less breathless than would be expected if congestion of the same degree were due to cardiac failure. Engorgement of the liver too is unusually prominent owing, so it has been thought, to direct interference with the hepatic veins. Great enlargement, cirrhosis of liver, and extreme perihepatitis are some times displayed, and are probably to be associated with long continuance of congestion. Dropsy of the legs is less, and ascites is more, prominent than in comparable cardiac failure, and dropsy of head and neck can occur, whereas in cardiac failure it does not. When the superior vena cava is obstructed, pulsation in it lacks freedom.

The feature that has special value in diagnosing constrictive pericarditis is conspicuous venous engorgement in a case that presents few or no signs of cardiac enlargement.

Inspiratory Swelling of Veins—The cervical veins normally collapse in inspiration because blood is sucked into the heart when thoracic pressure falls. In some cases of mediastinopericarditis an obstruction to the entry of blood occurs during inspiration, presumably because fibrous bands tighten about the heart or about the great veins; thus it happens, paradoxically, that the veins swell in inspiration.

Inspiratory Decline or Failure of Pulse—In mediastinopericarditis, inspiration of ordinary depth may interfere with the filling of the heart, as has been seen, or may in certain cases kink the aorta. Either of these interferences will cause the pulse to become smaller and pressure in the arteries to fall. In the same circumstances a deeper inspiration may sometimes obliterate the pulse. It is important to note that all the pulses in the body are affected

structures most affected in this way are the 5th and 6th rib cartilages and the xiphisternum, but retraction may extend into the axilla and involve lower ribs than those named. Through the diaphragm, systolic retraction of the lowest parts of the bony thorax may occur, seen at its left margin when the chest is viewed from the front, or at the left base when the chest is viewed laterally.

It is not quite correct to regard the heart as tugging the precordial ribs inwards through firm adhesions between it and the chest wall. When the heart tends to draw away from the chest wall, the latter must inevitably follow it, whether bound by fibrous tissue or not, provided that the lungs cannot move in to fill the gap. Ordinarily decrease in the size of the heart in systole is balanced by an equivalent expansion of the air containing lungs, which bound it laterally. This systolic expansion of the lungs calls for the expenditure of little cardiac energy. When inflexible fibrous tissue in pericardium and pleura separates the heart from the lungs, the heart expends energy in pulling in all directions upon relatively unyielding structures. Thus, fibrous tissue covering the heart laterally impedes its beat more than do any adhesions anteriorly, for the former alone guards a yielding structure, and these lateral adhesions come to be more important in bringing about precordial retraction than do adhesions binding heart and chest wall directly. Some energy may also be expended uselessly in direct pulls by the heart through fibrous bands and diaphragm upon more distant structures.

Retraction of the more rigid parts of the chest wall will naturally happen more readily when these are young and pliant, and will be greater the more vigorous the movement of the ventricle. Thus, although systolic retractions, and especially emphatic retractions, are valuable, they are inconclusive signs, there is no known form of retraction that is not to be observed on occasion apart from adhesions, moreover, advanced mediastinopericarditis with strong adhesions between heart and precordial chest wall may be present without any retractions occurring.

Immobile Impulse—Normally the impulse and left margin of cardiac dulness move 2 to 4 cms ($1\frac{1}{2}$ inches) to the left when a young subject rotates from the supine to the full left lateral posture, and equally to the right when the full right lateral posture is assumed. Adherency of the heart to the chest wall, or its enclosure within firm partitions of fibrous tissue, naturally renders it immobile. However, the positive use of this sign and of corresponding

X ray Signs—When the pericardium is greatly thickened and fibrosed, the borders of the heart are straighter than normal and the points of junction of vessels and different cavities tend to be indistinct. The rounded apex of the heart is not seen free of the diaphragm. Irregular shadows project from the silhouette of the heart or basal vessels, interfering with its normal outline and with the sharpness of its margins. The diaphragmatic movements are often greatly limited, especially in the region of the heart's apex, which moves little or not at all. Sometimes definite distortions of the diaphragm are observed, especially directly to the right of the heart, during inspiration, where adhesions may sometimes be seen. Immobility of the heart with change of posture can be identified. A well-exposed film may exhibit a calcified sheath upon the heart.

ASSOCIATED PHENOMENA

Signs of enlargement of the heart are usual, though not invariable, in mediastinopericarditis. Adhesions that impede the heart's action help to produce this enlargement. Because many of the patients have suffered from rheumatism, any other rheumatic manifestation, such as mitral or aortic disease, enlargement of the heart, auricular fibrillation, or failure arising from a defective muscle, may be associated (see Chapter XXII).

DIAGNOSIS

The diagnosis of mediastinopericarditis or of constrictive pericarditis depends largely upon their being remembered as a possibility in certain types of patient, for this will lead to a close and special examination. The possibility should be in mind in all cases of prolonged venous congestion of rheumatic origin, especially in young people. It should not be forgotten when, in these circumstances, the heart seems little enlarged (constrictive type), or when, the heart being enlarged, there are no definite signs of accompanying disease of the valves.

The signs upon which chief stress should be laid are retractions of the chest wall, combined with those indicating persistent interference with venous return, which may or may not be enhanced by the breathing being held in the inspiratory position.

SIGNIFICANCE AND PROGNOSIS

Simple adherency of the pericardium has probably no effect on the weight of the heart or upon the life of the subject. The condition may be found in people who have lived long and very active.

lives whose hearts are not enlarged and who have died of some quite separate malady. As it is undiagnosable it has no clinical significance.

It is rarely possible to separate the consequences of the more serious extrapericardial adhesions from those of accompanying lesions of valves or defects of muscle. The course run is therefore very variable. Thus the subject may die in childhood from involvement of the heart or from rheumatic reinfection or may survive and the course may be many months or several years during which time venous congestion is shown. In most cases however pericardial adhesions do not cause venous stasis by themselves but merely form one of several burdens the heart has to bear. Thus the adhesions are but a feature of the case and then have little influence either upon prognosis or treatment.

SPECIAL TREATMENT

When the heart is so shut in by fibrous bands that it cannot expand the lungs in systole much of its energy is expended in pulling resistant structures like this fibrous tissue and the chest wall towards it. It can be relieved of this work by making one side of the resistant chamber in which it lies relatively flaccid. It is to this that removal of rib and cartilage owes its success as a remedy. Not only the tug on the chest wall but on surrounding fibrous tissue is relieved. The operation may be performed under local anaesthesia. The three ribs mainly tugged upon—and these are usually the 3rd, 4th and 5th—are removed for a length of about 12 cms over the precordium. The ribs must be separated from the heart with the utmost care so that tearing the muscle of the heart and pneumothorax may be avoided. The operation is a very successful one in well or fortunately chosen cases and in skilful hands. Any suspicion of active rheumatism contraindicates operation. The chief indication for operation is definite intugging of the ribs and the amount of work that the heart is expending on these may be estimated by noting how much force is required to depress symmetrical ribs similarly. On occasion strong bands of pericardium have been divided successfully. The direct mortality of the first and less difficult operation is stated to be about 5 per cent. To balance against this mortality is conspicuous improvement in a considerable percentage of the cases and occasionally restoration of a complete invalid to active life. The main obstacle to surgical treatment is however the diagnostic one.

CHAPTER XX

BACTERIAL ENDOCARDITIS

SUBACUTE BACTERIAL ENDOCARDITIS

THE subacute variety of bacterial endocarditis is here emphasised because it is much the commonest form, and because it often complicates old standing heart disease. The malady is most prevalent between the ages of twenty and forty and is commoner amongst males than females.

PATHOLOGY

There is much to suggest that subacute bacterial endocarditis is due to an infection of relatively low virulence, which attacks valves that are already abnormal. The bacterium responsible for the disease is almost always a nonhaemolytic streptococcus and one that grows poorly in culture. Very occasionally it is the bacillus "influenzae". The vegetations are almost confined to the left side of the heart, the side on which chronic valvular disease is common. The chief exceptions to this rule are cases of congenital malformation in which the right heart is involved, as in patency of the interventricular septum, pulmonary stenosis, or patent ductus arteriosus. The disease ordinarily attacks the mitral or aortic valve. In most cases one or other of these, and usually the aortic valve, is the main seat of the disease, both may be heavily affected and, if the valves are searched for minute vegetations, it will be realised that the cases are few in which either valve escapes completely. In a very considerable number of the cases involving the aortic valves, these valves are congenitally bicuspid, this deformity occurs in little less than 1 per cent of all autopsies, and about a fifth of all such cases of congenital malformation acquire infective endocarditis, it is therefore a notable factor in determining the disease. In other cases it is clear that vegetations have formed on old standing rheumatic disease of mitral or aortic valves, a history of rheumatic fever or chorea is present in

a high percentage of the patients Syphilitic disease of the aortic base is also found, though less frequently

Clinical observations show that cardiac patients who acquire this disease do so for the most part while in good or fairly good general health cases of congestive failure are relatively immune It is the rule to find numerous vegetations and not a single mass the vegetations often involve all the cusps of either aortic or mitral valve or both The vegetations are mostly small, warty, or shortly pedunculated some are much larger, club shaped, strap like, or form large irregular masses they affect the lines of apposition of the cusps first of all but in heavier involvement plaster the whole surface of the cusp and appear in smaller or larger confluent masses upon the walls of auricle or ventricle upon the lowest parts of the aortic wall, and upon the chordae tendineae some of which are frequently cut through The parts of affected cusps that are free of actual vegetations are nearly always thickened, often much fibrosed and deformed and sometimes show healed perforations Often vegetations are found that are in an advanced stage of fibrous, strap like vegetations are always tough at their bases calcareous deposits are not infrequent In such cases, and they are those that have run long courses, the appearances indicate that a process of new infection by contact or spread, and a process of repair, continue to challenge each other in a fight for predominance, which is carried on over long periods of time Sometimes freshly broken vegetations are found where emboli have become detached

CHIEF CLINICAL MANIFESTATIONS

Onset —The malady, regarded as an infection, generally begins in early adult life and its onset is usually insidious, so insidious that sometimes it may be many months before the subject realises his health to be seriously affected The usual early symptoms are lassitude and weakness, combined perhaps with occasional feelings of chilliness and malaise, perhaps with aches about the joints or muscles, these symptoms continue and the patient passes gradually to a state of debility and anaemia Anorexia with or without nausea, and later vomiting, is an early symptom in some cases The first symptoms may be those of an acute febrile illness with chills and sweating The first symptoms may arise from embolism of one of a number of important vessels, giving a varied and sometimes puzzling symptomatology

The phenomena of the established condition are conveniently described in three groups. Firstly, there are the local signs in the heart corresponding to old standing mischief and to new damage of the valves consequent upon their recent inflammation. Secondly, the disease may be regarded as equivalent to a condition of chronic septicaemia or intoxication, most of the manifestations are properly placed in this category. Thirdly, there are the very varied effects of embolism.

Local Signs—Since the disease is one which attacks valves that are often much deformed already, and as the disease itself damages the valves further, perfectly clear signs of valvular mischief are almost always to be obtained. The cases are generally those of aortic regurgitation or of mitral stenosis, the remainder are interpretable as mitral regurgitation or are instances of congenital heart disease. When there are no signs pointing emphatically to disease of the valves, signs of definite or moderate enlargement of the heart will usually be found. It will be understood that the degree in which these local signs are attributable to the disease we are considering can be estimated clinically only if the state of the heart before its infection is precisely known.

That the valves are actually the seat of active mischief is sometimes betrayed during the course of the malady by signs of new local damage, as when a murmur indicative of aortic regurgitation or of mitral stenosis changes its character conspicuously, or actually develops. These changes, however, are not witnessed very frequently and therefore have no great clinical importance.

Failure of the heart with the development of venous congestion, is more appropriately placed in the next category (page 187).

Symptoms and signs of infection are numerous and most of them are common to this and similar infections. They include weakness, gradual loss of flesh and many manifestations that require special mention.

Fever is the rule though it is often inconspicuous. In the early stages of the disease and particularly when the onset is insidious or the course long drawn, the temperature may be raised only 1° to 3° F, and there may be periods of days or weeks during which it is normal. It is rather the exception than the rule for temperature to be raised in isolated readings taken in the daytime from patients who are up and about. The temperature chart of bed cases is very variable, almost any type of curve may be shown but low grades of fever are the rule, high remittent or intermitt-

continuous fever is rare before the terminal phases of the malady are reached

Pallor is not invariable even when infection has long established itself. A few patients have the full and fresh colour of rude health. But pallor is the rule and is often as extreme as in pernicious anaemia. The tint is variable as is that of the bloodless normal skin. ■ special reason for variation is found in chronic cases, which acquire a distinct but diffuse cutaneous pigmentation. A yellow pallor is usual in blondes, pale brown or muddy tints in those pigmented spontaneously or by the disease. Pallor ■ due largely to lowness of red blood cell count which is commonly near to three million or less; this is often aggravated by a low colour index, the anaemia being in some of chlorotic type.

Leucocytosis is the rule, the count of white cells being usually between 10,000 and 15,000, occasionally much higher, occasionally normal or actually lower than normal. The differential counts vary, but polymorphonuclear cell increase is much commoner than lymphocytosis.

Splenic enlargement is frequent in almost all cases and, the spleen being firm, is palpable in most cases; this is a very important sign. The tip of the spleen is just palpable or comes three or four fingers' breadth below the costal margin; exceptionally it is much bigger and fills a large part of the left upper abdomen. This enlargement is to be ascribed to infection as in the case of typhoid fever.

Clubbed fingers ■ another manifestation due in all probability to long lasting toxæmia, as it is in phthisis and bronchiectasis. Swelling is confined to the last phalanges of the fingers; the swelling may be anteroposterior and accompanied by increased convexity of the nails, or it may be lateral. It is not often so advanced as to produce really bulbous fingers as in congenital heart disease, a noteworthy contrast is that these clubbed fingers in subacute endocarditis rarely exhibit cyanosis. Clubbing of the toes occurs, but is more difficult to recognise.

Petechiae, scattered in small numbers and appearing often in crops, are usual; they are especially frequent above the clavicles and around the axillae, on the upper arms, chest, and abdomen. They are scarlet, red brown, or brown, according to their age, and they last a few days. They are to be distinguished from the small bright crimson angiomas that are so common in the skin, and which are either distinctly elevated or, if smaller, blanch a little or much on pressure. When petechiae are seen in the conjunctival or buccal

mucous membrane or under the nail, where they are often linear, there can be no confusion with the bites of parasites. Petechiae should not be regarded as embolic the minute vessels of the skin in this state permit an escape of blood more easily than normal, the petechiae can be produced by congesting the arm. Though occurring so often in endocarditis their diagnostic importance is much decreased by their frequency in many anaemias and intoxications.

Purpura occurs occasionally and may be widespread, but is usually found as a symmetrical eruption on the legs, it occurs in the later stages of the malady as do rare haemorrhages from mucous membranes. Other cutaneous eruptions, including urticaria and different forms of erythema are occasional.

eff
ch
and pleurisy occur rarely

Renal involvement is common. Abnormalities of the urine, as the presence of a trace of albumen hyaline and granular casts or pus cells in small numbers are frequent. Occurring alone these changes are insufficient evidence of nephritis. Per

frequent and sometimes affects these structures in large numbers. In the later stages clear evidences of parenchymatous nephritis are not infrequent the urine is loaded with albumen, renal colic occurs and uraemic manifestations may appear.

Cardiac failure with congestion is rarely present in the early stages of this disease which attacks those who are in relatively good health but it is so frequent in the later stages as to form a chief cause of death it is always terminal. Failure with congestion in these cases is the end result neither of the original burden imposed upon the heart nor of its increase through an aggravation of the valve defects. The total valve defect is often slight, and the failure develops while the patient is at rest in bed and is an obvious accompaniment of infection. The disease affords one of the clearest illustrations of failure being determined by an infective process (page 23) but as there are no recognisable lesions in the heart muscle that can be called to account for this failure the weakening of the muscle is to be attributed to poisoning the weakening

Embotic Accidents—The emboli are derived from the heart, a large vegetation, or a considerable portion of it, becomes detached, or it breaks into a number of pieces, and vegetation or its fragments are swept on with the blood current. The vegetations being almost confined to the left heart, embolism is almost confined to the systemic system. The size of the embolus determines the size of vessel that can be entered and plugged. The direction of arterial branching may be disregarded and the travelling embolus considered as merely part of the current of blood. No artery is immune from embolism and chance chiefly determines which is affected but blocking of certain vessels has conspicuous effects, thus attracting unusual attention and giving a false idea of the relative frequency of embolism in these.

The chief effect is interference with local blood supply in instances in which a proper collateral circulation fails to become quickly established. Embolism of the main vessel of a limb may cause gangrene of the middle cerebral artery, hemiplegia, of a coronary artery sudden death of a retinal artery partial blindness. Embolism of a splenic vessel infarcts that organ, causing local pain and tenderness vomiting and audible friction in the splenic region. Infarction of the kidney brings albumen and blood into the urine, pain in the back and sickness and rarely when blood is abundant and clots form causes symptoms resembling renal colic.

The emboli of infective endocarditis often give rise to symptoms and signs unseen in the case of uninfected clots. The events are well illustrated in superficial vessels. An artery like the radial or dorsalis pedis is blocked by an embolus. This may happen so unobtrusively as to escape any attention. It may be noticed on careful search that the pulse has disappeared. In other instances local pain and tenderness follow the embolism this may culminate in local reddening of the skin lasting a few days, or with swelling and induration of the tissues around the artery for a longer time. In a few cases chronic arterial and periarterial inflammation continues, the vessel wall gives, and an aneurysm forms locally. Abscesses very rarely occur. These different disturbances illustrate the varying power of the tissues to deal with the infection. Essentially similar changes occur also in deeply seated vessels, for example, aneurysms form on branches of arteries supplying the viscera.

Infected emboli in smaller arteries, like the nasal, may give conspicuous inflammatory redness and swelling locally. Digital nodes are small red or purple spots a centimetre or less in diameter, always

tender when fresh sometimes raised and often just palpable and occurring in the pads of fingers and toes They last a few days and are important diagnostically Similar but more extensive lesions also occur in the skin of the thenar and hypothenar eminences These cutaneous or subcutaneous lesions are thought to arise from infective emboli of small arteries or arterioles Similar emboli may probably account for occasional pain and tenderness appearing in muscles and near joints

DIAGNOSIS

The diagnosis of subacute bacterial endocarditis has been obscured in the past by failure to recognise the insidious nature of the complaint and by the expectation of finding high temperature or at least some fever Misinterpretation of three of the cardinal physical signs has impeded and continues to impede its recognition these signs are clubbed fingers enlarged spleen and pallor In adults presenting definite signs of chronic heart disease clubbing of the fingers has been regarded wrongly as a simple association of this heart disease In about four fifths of these cases clubbed fingers are due to infective endocarditis though not diagnostic they constitute one of our most valuable signs of the disease Occasionally they are the result of congenital heart disease the fingers then being cyanosed and their tips more bulbous very occasionally the clubbing is itself congenital and the patient will then usually be sure that the fingers have always been misshapen swelling at the base of the nail due to irritation from breaks in the skin of the lunula should not be confused with clubbed fingers An enlarged spleen has long been regarded as a consequence of cardiac failure Now the spleen unlike the liver has a muscular framework and it is unusual for it to increase much if at all in size in congestive failure In chronic heart disease an enlarged spleen generally means infective endocarditis though its significance is less in countries where malaria is prevalent Pallor has been looked upon as a common result of aortic regurgitation Aortic reflux does not produce pallor when the two are found together an adequate cause for pallor must be discovered when pallor is conspicuous it is usually due either to infective endocarditis or to rheumatic infection according to whether the patient is an adult or a child

In identifying infective endocarditis the first requisite is clear evidence of cardiac disease Without this the diagnosis will contain an element of doubt that can scarcely be resolved before signs of embolism occur and streptococci are found in the blood

Given this evidence of heart disease, the presence of continued or periodic fever always arouses grave suspicion of the disease. The addition of splenic enlargement increases the suspicion, but a diagnosis cannot yet be made, when heart disease is complicated by

only signs of heart disease, pernicious and splenic anaemia must sometimes be taken into account, the presence of clubbed fingers, signs of embolism, or a positive blood culture will settle the diagnosis in each instance. Irrespective of other signs, if the patient is afebrile and remains so for many weeks, the diagnosis becomes highly improbable and is shortly ruled out.

In a case of acquired heart disease, clubbed fingers at once points to the possibility of infection, splenic enlargement, conspicuous pallor or red blood cells in the urine, will each render a diagnosis highly probable, even when at a single examination the temperature is unraised.

In a heart case embolism combined with fever, splenic enlargement or positive blood culture suffices, or it is enough if the emboli are known to be infected, or if digital nodes occur.

A blood culture showing nonhaemolytic streptococci is insufficient for a diagnosis even when heart disease is present, but if fever, enlarged spleen, clubbed fingers, or embolism is added the diagnosis becomes certain. Failure to cultivate the organism from the blood is of little value, unless the failure is repeated during long continued fever. A certain diagnosis can often be made from clinical signs when failure to grow the organism has been frequent.

COURSE AND PROGNOSIS

The course of a disease presenting such a variety of symptoms and signs naturally varies. It is a progressive disease, ending fatally with so few exceptions that little or no hope of the patient surviving it is to be entertained. Instances of reported recovery have not always been substantiated, but a few seem well authenticated. The average course from onset of symptoms till death averages six months, the termination may come earlier, or it may be prolonged, not unusually till eighteen months and occasionally till two, and rarely three, years. Inconspicuous fever and frequent and long apyrexial intervals are usual in the cases running long courses. Death occurs from progressive weakness and exhaustion or from

cardiac failure in about equal numbers of the cases, and these together form about two thirds of the whole. Serious embolic accidents, nephritis, and intercurrent maladies, including pneumonia, account for most of the remainder.

TREATMENT

No treatment that has been tried cures. Very many remedies have been advocated, but none has as yet stood the test of time. Intravenous antiseptics such as salvarsan and sodium cacodylate, eusol, dyes such as flavine and gentian violet, vaccines and serum therapy, deep X rays, radium emanations, diathermy over the heart, heavy dosage with vitamin *a*, intramuscular injections of nucleic acid, ultra violet light treatment, blood transfusion, these are a few amongst the many failures.

The patients should be treated in their homes so far as possible. They should be in bed when febrile, they can move about when afebrile, strolling quietly or sitting or lying out of doors in sunlight and in warm weather, according to general fitness. Their food should be sustaining, light as a rule, heavier when tolerated. It is important for them to be occupied, the visits of cheerful but quiet friends are very helpful. Symptoms are met as they arise, but the treatment is chiefly in the hands of friends and nurses. It injures these patients to ply them with remedies.

ACUTE BACTERIAL ENDOCARDITIS

PATHOLOGY

This form of bacterial endocarditis can occur at any age. It often originates in a serious local infection in some other part of the body, as in pneumonia, osteomyelitis, gonorrhoea, or puerperal sepsis.

The bacteria responsible are usually the pneumococcus, staphylococcus aureus, gonococcus or haemolytic streptococcus, other organisms occur occasionally. The valves of the right heart are much oftener affected than in the subacute infection, the aortic valve is attacked most frequently of any, and after this the mitral. One valve is usually the seat of disease and sometimes one cusp. The valve attacked may be the seat of previous mischief, but usually this is not the case, and the uninfected cusps, or parts of the cusp, have their normal thickness and translucence. The vegetations are soft, usually large, not infrequently forming a cauliflower like mass. Spread to the septa is occasional only. Ulceration is frequent and cuts deeply,

often perforating the valve. The tissue reaction is acutely inflammatory. Signs of repair are few both in the vegetation and around its base. This is so because resistance is slight and the disease runs a short course.

CHIEF CLINICAL MANIFESTATIONS

Onset—Developing in pneumonia or as a complication of other grave infection the onset is inconspicuous. In cases that are primarily cardiac the onset is usually with high fever and its accompaniments—headache, vomiting, malaise, chills and sweating.

Local Signs—Signs of valve disease may be present from the start but more often develop. A systolic murmur appearing at the cardiac impulse or to and fro murmurs at the base, the latter may be to right or left of sternum and due to involvement of pulmonary or aortic valve. Frequently there are no signs of valve disease. Change in the character of the murmurs is noted occasionally. Cardiac enlargement unless existing previously is not the rule. Inconspicuousness or lack of cardiac signs is responsible for the chief difficulties of diagnosis.

Symptoms and Signs of Infection—These may be enumerated as for the subacute form though much more briefly. The contrasts between the two diseases are chiefly due to greater severity of infection and the shorter course run in the acute variety.

Fever is constant, is continuously high, remittent or intermittent. Temperatures of 103° or more (up to 105° or 106°) are reached almost daily. Rigors are common and profuse sweats the rule. The gravity of the malady is shown by the prostration of the patient by his apathy or by delirium or coma. Pallor is usually inconspicuous. Profound anaemia is rare. Clubbing of the fingers scarcely occurs. Time is usually too short. The polymorphonuclear leucocytosis is considerable, the white cell count being from 20,000 to 30,000. The spleen presents signs previously noted but being soft is not so often felt as in the subacute disease. Petechiae and purpura are common. The joints are less often and the pericardium and pleura more often involved than in the subacute disease. Congestive failure is unusual. Albumen and red blood cells are often found in the urine and acute but not chronic nephritis may develop.

Embolic Accidents—These are very frequent and the detached fragments are mostly large. By obstructing the blood flow through important arteries they cause corresponding symptoms already described in detail. Because the valves in the right heart are often affected pulmonary embolism is not unusual. In superficial arteries

an inflammatory reaction is the rule and embolic abscesses are common. Digital nodes are not a feature of this disease.

DIAGNOSIS

The disease is more difficult to recognise than is subacute bacterial endocarditis. That is so because signs in the heart are found with far less frequency and the diagnosis then comes to depend almost entirely on evidence of embolism. Without cardiac signs or emboli the disease cannot be differentiated clinically from simple septicaemia. In pneumonia in which the original temperature curve is continued or rises the diagnosis will waver between endocarditis, tuberculosis and empyema. Continuous temperature lasting for several weeks accompanied by increasing weakness, delirium and splenic enlargement produces a picture like typhoid fever though in this the pulse rate is lower. The Widal reaction differentiates. Miliary tuberculosis is a possibility that must also be kept in mind. Local and deep seated collections of pus such as are found in central empyema and perinephritic abscess must be remembered. Blood cultures are always positive the organisms growing readily.

COURSE, PROGNOSIS AND TREATMENT

In the severest infections and in cases in which the patient is already the subject of grave disease death may occur within a few days. The usual course is a few weeks, sometimes two and rarely three months. Death is generally the result of poisoning the patient weakening progressively. Embolism of brain or heart occasionally kills. Recovery is almost unknown. The patients are mostly helpless and require skilful and constant nursing—to administer food in liquid form through lips often unresponsive to relieve high fever by tepid sponging, to keep clothing dry amidst drenching sweats, to maintain cleanliness and yet not to hasten the exhaustion which comes so quickly despite all care.

CHAPTER XXI

RHEUMATIC CARDITIS

INTRODUCTORY

To understand rheumatic heart disease as it is seen in its developed forms, it is necessary to study its evolution. The conception, still prevalent that what we call by the name "rheumatic fever" is a disease of the joints and that this disease is frequently complicated by involvement of the heart cannot be justified. The disease is a widespread affliction of the body, usually, it is true, but by no means always, producing arthritic symptoms. To regard the joints as the primary or central point of the disease is to go beyond the facts. To convey the idea, by the use of the traditional and almost always unsafe word 'complication', that something is added which is not essentially a part of the disease, is to assume more than we know, and probably more than is true.

The modern conception of the disease is of an organism, possibly, if not probably streptococcal in kind, that invades the blood stream, damages the small arterioles of the body and sets up a condition of inflammation, acute or subacute, that calls attention to itself especially in and around the joints, in the heart as a whole, and in the subcutaneous tissues and tendons. The clinical picture is very varied, but the most notable manifestations are the intoxication, arthritic

nodule, and chorea may each be in chief evidence, or appear as the only obvious clinical accompaniment of the infection, any two of these may be paired (except arthritis and chorea), and often three are present simultaneously or in other instances within short times of each other in a given case.

The disease is specially one of childhood, but age modifies it, chorea, nodules, and carditis becoming much less frequent shortly

before or at the time puberty is reached, while disturbances of the joints increase in severity and acuteness to become the dominant feature of the disease from the first decade of adult life. The name rheumatic fever is apt to conjure up the picture of the acute and relatively transient illness of the adult, but, while the heart may and often does suffer damage in such attacks of the disease in the third decade of life, rheumatic heart disease begins almost always in childhood. In childhood the disease strongly tends to be relapsing and therefore to run a chronic course. The full active course is often run, and life often brought to an end, before the child comes of age. Rheumatic heart disease as it is seen in the adult consists in very large part of the damage arising out of the inflammations of these early years. Mitral stenosis is the result of such damage, this and other injuries of the kind are sometimes found in patients in whom there is no history of illness to account for them, but if chronic rheumatic heart disease is to be seen in the actual making, to be followed through the stages of its progress, it is to the growing child that we must go.

RHEUMATISM OF CHILDHOOD

The disease is one of cold and damp countries of the northern hemisphere, and is found particularly amongst the overcrowded and poorer parts of the population of towns. It appears especially in the winter months. The age period begins at about five years and rises to a peak at ten and subsequently and gradually declines. It affects girls more often than boys.

PATHOLOGICAL ANATOMY

In fatal cases the pericardium is nearly always affected in some degree. Its surface is congested and dull, and there are small deposits of fibrin upon it. Often the pericardium is universally inflamed and swollen, and covered by a thick layer of fibrinous material. Pericardial fluid may be in excess and is turbid, but in children is rarely large in amount. In cases that have run a long course, universal soft adhesions are common. In a few mediastinopericarditis occurs (page 176).

Whenever the valves are involved, and this probably happens always in a case of carditis, the mitral and aortic valves are usually both affected in greater or less degree, the mitral valve is probably involved invariably and usually heavily, the aortic valve is involved

a little less frequently and usually less heavily. The tricuspid and pulmonary valves come next in this order the former being involved in as many as a third of the cases if we count slight disease and the latter infrequently.

The lesions on the surface of the valve consist of the almost characteristic small sessile vegetations set in a band along the margins of apposition and involving the whole circumference of the valve when this is heavily invaded and occasionally affecting other portions of the valves such as the chordae tendineae. Beneath the surface is the more notable active inflammation of the valve substance. These lesions represent the active stage of the disease as this attacks in the first instance healthy valves and in recurrent attacks valves already scarred by similar past injuries. In healing they leave behind thickened shrunken tissues narrowed orifices incompetent valves as previously described in detail (pages 120 and 129).

In most cases cloudy swelling of the heart muscle is present. In cases running a longer course this is less in evidence and the subendocardial nodule is the characteristic lesion. These nodules present a fibrinous or necrotic centre and contain large bipolar or multinuclear darkly staining cells young fibroblasts lymphocytes and a few leucocytes. They are frequently found around small blood vessels and occur in the interstitial tissue of the auricles and ventricles especially around the valve rings in the subendocardial tissue in the pericardium and aortic adventitia. They may be few in number or the heart may be crowded with them. They also occur in many other parts of the body especially in the periarticular tissues. They are thought to be peculiar to rheumatic inflammation and are often used to identify it. In healing they leave behind at the most a minute fibrous scar. The subcutaneous nodules or granulomata though much larger are composed of similar tissue elements and evidently represent a similar form of subacute or chronic inflammatory reaction.

SYMPTOMS AND SIGNS

Onset—Although the disease may begin abruptly with fever an insidious onset is commoner. The child is brought with the complaint that it feels poorly has lost appetite is easily tired is pale and has been losing flesh there may be pains in the limbs or chest.

Pulse and Fever—In this disease the pulse rate is a more valuable guide to the presence of infection than is the temperature chart.

Fever is often absent from both short and long attacks but the pulse rate is always quickened too the temperature is usually low and often falls as joints or heart become involved, it almost always subsides in cardiacitis, it falls away by lysis. Temperature variation. A phase of fever lasting ten to thirty days or may be repeated one or more times with intervals during which there is no fever, but during which the pulse is raised. Cases in which short and slight phases of fever are a period of six or nine or even twelve or eighteen days are not uncommon. Hyperpyrexia is rare. In general, the less fever is in evidence. There may or may not be pallor, often rapid in development is usual and is associated with actual anaemia.

Joints—Involvement of joints is the rule, often and manifests itself by pains only. Usually the wrists and tenderness of such joints as the wrist or the joints are involved they are often involved in the early stages.

Carditis and its Manifestations—Even in the early stages of the disease it is involved in a high percentage of cases it is involved in prolonged or recurrent infections of long duration. Involvement a rise of pulse rate is conspicuous and there is breathlessness. The cardiac dulness and the impulse moves out the first sound becomes less distinct. A murmur is heard in the apical region. The aortic regurgitation may appear to the left of the sternum. Lasting a few days is not uncommon over the first few days.

In severe attacks the onset of pericarditis is vomiting by a sharp rise of temperature, and in severe attacks cordial pain and tenderness are frequent. As the severe attack breathlessness increases the veins are engorged, and other signs of failure with cyanosis.

failure is not due to valvular defects for it is usually undeveloped neither can present day histological changes in the muscle that adequately account for it comes and it goes, hand in hand with other signs of intoxication, and is to be attributed to the

The rhythm of the heart may be disturbed by the occasional occurrence of partial heart block or extrasystoles

Nodules are seen clinically in the subcutaneous tissues in about a tenth of the cases, and especially in those running a very prolonged and severe course. They occur chiefly over bony points such as the elbow knuckles knee ankle spines of vertebrae, and cranium, but also on the tendons and tendon sheaths of hands and feet. There may be scores of them but more often they are few in number and must be searched for closely. Large nodules, several centimetres in diameter are easy to detect but those of a few millimetres which are much commoner are apt to escape attention. They are best displayed by putting the skin on the stretch, when the nodules lift and blanch the skin. The skin moves easily over them, they are painless and firm or almost hard to the touch, and are attached to the deeper tissues though usually slightly movable. The larger masses are often aggregates of smaller ones. Nodules appear quickly, and last a few days or weeks.

Sweat rashes are common of the pronounced skin eruptions, erythema multiforme is the most frequent, purpura is occasional.

Chorea—The nervous syndrome with its mental disturbances, muscular weakness and inco-ordinate movements will not be described here. Occurring alone it runs a course of six weeks or longer, which is almost always without fever. The heart is much less frequently involved in this type than when there are arthritic changes. Carditis occurs however and, though usually mild, is sometimes severe.

COURSE AND PROGNOSIS

Death may occur in acute and very severe cases, but these are rare. A first attack is rarely fatal. When cardiac dilatation has been slight and an apical systolic murmur has developed, these signs disappear more often than not as a first attack subsides. But the disease is one that has a strong tendency to recur or to present relapses and when the original manifestations of cardiac involvement are more serious, or when the signs reappear in a recrudescence of activity, evidence of residual damage is more frequent and permanent. It is not clear that the heart becomes quite free from active disease during periods of symptomatic quiescence, and that there is not still a smouldering inflammation, but it does seem beyond doubt that the main mischief is usually done by bursts of more acute inflammation that are recognisable at the time. Thus, while recovery of normal cardiac signs is frequent after a single febrile illness, it is

unusual after a second and is very rare after a third, the statement being true if the attacks are separated by long or comparatively short periods of time. Each attack may leave the heart noticeably changed for the worse. The amount of cardiac involvement is related to the length of the active clinical course, and the cases, and they are many, that run from one relapse to another for periods as long as six months or even a year or more often emerge from the illness, if they survive it, in a permanently crippled state. Even so there is no immunity, further infection slight or grave being the rule.

The development of mitral stenosis is gradual, the signs becoming clear in general after a number of distinct illnesses or after very long drawn periods of recrudescence. Amongst the early signs are the harsh systolic murmur, the appearance of a sound in mid diastole, giving a form of canter rhythm. The extra sound is probably an accentuation of the normal 3rd heart sound, often occurring at this phase of the cycle and especially to be heard in children. Later a diastolic murmur appears at the same phase of diastole, or runs through diastole as it does characteristically in mitral stenosis of the adult. Another slow development is firm adherency of the heart to the chest wall and other surrounding structures in cases where pericardial inflammation has not confined itself to obliterating the sac but has spread beyond it.

The description of rheumatic heart disease as it develops to become chronic and as it is here described in outline is incomplete in two particulars. On the one hand, it is clear that the heart may also be affected permanently by rheumatic fever occurring in adult life, and manifesting itself as a more acute attack in which arthritis is prominent, though in this form the heart usually escapes. On the other hand, the past rheumatic illness cannot always be identified. In cases of mitral stenosis in adults a history of infection is obtained in about half the cases only: this is due in part to reliance upon a history of an illness with arthritic symptoms. It is due also to the development of serious heart disease as an insidious rheumatic process, unaccompanied by such symptoms as would be regarded by patient or patient's friends as constituting definite illness. The more closely children who have displayed single and definite attacks of rheumatism are watched, the more evident is the difficulty of stating that there is no longer active mischief. Without presenting fever the child from time to time is unfit and has perhaps pains in the limbs, the pulse rate is at times unduly rapid, the routine

The rhythm of the heart may be disturbed by the occasional occurrence of partial heart block or extrasystoles

Nodules are seen clinically in the subcutaneous tissues in about a tenth of the cases and especially in those running a very prolonged and severe course. They occur chiefly over bony points such as the elbow knuckles knee ankle spines of vertebrae, and cranium, but also on the tendons and tendon sheaths of hands and feet. There may be scores of them but more often they are few in number and must be searched for closely. Large nodules several centimetres in diameter are easy to detect but those of a few millimetres which are much commoner are apt to escape attention. They are best displayed by putting the skin on the stretch when the nodules lift and blanch the skin. The skin moves easily over them, they are painless and firm or almost hard to the touch, and are attached to the deeper tissues though usually slightly movable. The larger masses are often aggregates of smaller ones. Nodules appear quickly, and last a few days or weeks.

Sweat rashes are common of the pronounced skin eruptions, erythema multiforme is the most frequent purpura is occasional.

Chorea—The nervous syndrome with its mental disturbances muscular weakness and inco-ordinate movements, will not be described here. Occurring alone it runs a course of six weeks or longer, which is almost always without fever. The heart is much less frequently involved in this type than when there are arthritic changes. Carditis occurs however and though usually mild, is sometimes severe.

COURSE AND PROGNOSIS

Death may occur in acute and very severe cases but these are rare. A first attack is rarely fatal. When cardiac dilatation has been slight and an apical systolic murmur has developed these signs disappear more often than not as a first attack subsides. But the disease is one that has a strong tendency to recur or to present relapses and when the original manifestations of cardiac involvement are more serious, or when the signs reappear in a recrudescence of activity evidence of residual damage is more frequent and permanent. It is not clear that the heart becomes quite free from active disease during periods of symptomatic quiescence and that there is not still a smouldering inflammation but it does seem beyond doubt that the main mischief is usually done by bursts of more acute inflammation that are recognisable at the time. Thus, while recovery of normal cardiac signs is frequent after a single febrile illness, it is

unusual after a second and is very rare after a third the statement being true if the attacks are separated by long or comparatively short periods of time Each attack may leave the heart noticeably changed for the worse The amount of cardiac involvement is related to the length of the active clinical course and the cases and they are many that run from one relapse to another for periods as long as six months or even a year or more often emerge from the illness if they survive it in a permanently crippled state Even so there is no immunity further infection slight or grave being the rule

The development of mitral stenosis is gradual the signs becoming clear in general after a number of distinct illnesses or after very long drawn periods of recrudescence Amongst the early signs are the harsh systolic murmur the appearance of a sound in mid diastole giving a form of canter rhythm The extra sound is probably an accentuation of the normal 3rd heart sound often occurring at this phase of the cycle and especially to be heard in children Later a diastolic murmur appears at the same phase of diastole or runs through diastole as it does characteristically in mitral stenosis of the adult Another slow development is firm adherency of the heart to the chest wall and other surrounding structures in cases where pericardial inflammation has not confined itself to obliterating the sac but has spread beyond it

The description of rheumatic heart disease as it develops to be

and manifesting itself as a more acute attack in which arthritis is prominent though in this form the heart usually escapes On the

(1 2 3 4 5 6 7 8 9 10 11 12)

history of an illness with arthritic symptoms It is due also to the development of serious heart disease as an insidious rheumatic process unaccompanied by such symptoms as would be regarded by patient or patient's friends as constituting definite illness The more closely children who have displayed single and definite attacks of rheumatism are watched the more evident is the difficulty of stating that the illness is a simple rheumatism It is often a fever that presents itself as a rhapsody of symptoms in the routine

examination brings to light a few days of slight temperature, a friction rub a temporary prolongation of the A V conduction interval, a dropped beat or other sign that but for close and frequent scrutiny would escape notice. Similar incidents are to be noted in young patients with mitral disease who give no history of past illness. Another indication of stealthy invasion is the not infrequent discovery of unsuspected active lesions of rheumatic type in the hearts of patients dying of cardiac failure.

The mortality during the active stages of the disease is heavy, usually estimated at about 5 per cent in the first year, and 20 per cent in the first ten years after the onset. The rate varies much, however, and is extremely high amongst slum children who fail to obtain early care and treatment. It is variously reckoned that of those developing rheumatism in childhood 60 to 80 per cent ultimately manifest clear signs of heart disease.

The prognosis for the individual attack of active mischief depends upon the general strength of the child, the appearance of debility and anaemia being unfavourable. It depends especially on the signs in the heart and particularly upon the degree of failure present, and it depends upon the course the infection is taking and that is often impossible to estimate. A high brief fever that has been falling is more favourable than a lower fever long continuing. A first attack is more favourable than a second, a second than a third, as already indicated.

The distant future of the individual case depends upon when the injury to the heart by active rheumatism will end, and the extent of mischief that will then prevail. To prophesy at all accurately would require not only precise knowledge of the damage already present, but foresight of the time and severity of reinfection. The prognosis from this standpoint will be greatly simplified when we are able to prevent the recurrence of active mischief. It is manifest from experience that patients who are properly guarded are in a much more favourable position than those who are not. It is also in general clear that as each year passes and the child is growing towards adolescence and maturity, the most dangerous period is passing. The outlook is most unfavourable in the child continuing in a crowded and unhygienic home, and maintaining in these circumstances signs of ill health and presenting frequent recurrences of low grade infection. The outlook is unfavourable in cases presenting subcutaneous nodules.

TREATMENT

The patient in whom active disease exists should be removed from the unhygienic home to an institution with proper nursing facilities. Rest is essential until active mischief is no longer suspected, namely, until temperature and pulse rate are continuously normal and have been so for several or, in the case of long illness, many weeks. An object of rest is to avoid strains of joints or increased work of heart that may help to light up active trouble. Tender joints should be swathed in cotton wool. Patients who are febrile and sweating should be dressed in flannel or woollen clothing that is easily removed, and food should be soft and much of it taken in the form of fluid. Fever is treated by means of sodium salicylate or the acetyl compound (aspirin), both fever and pain are abated by these remedies, though they appear not to influence the course of the disease otherwise. Pyramidon, being less toxic than salicylic acid derivatives, is also used, but being less effective in lowering fever is more suited to the treatment of chorea. The pain of pericarditis and restlessness are treated with Dover's powder. The use of digitalis for failure with congestion, in this or any other acute febrile illness, is not usually to be recommended.

Patients in whom the disease is running a long course should be moved whenever possible to a hot sunny climate. The benefits of this procedure are rapidly displayed and signs of infection often completely disappear. Experience is beginning to show, too, that recurrences are rare so long as the child is kept in tropical or sub-tropical climates. In temperate climates these conditions should be reproduced as far as possible by keeping warm those patients who are fit to be treated on couch or on chair, and by exposing them to air and sunlight.

Convalescence should proceed slowly and be well consolidated, in cases of carditis the end of active infection is most difficult to gauge, and months must pass before recovery can be regarded as complete. Return to exercise must be very gradual. Tonsils that are obviously diseased, or frequently become infected, should be removed, but routine tonsillectomy has not been shown to reduce the frequency of reinfection.

CHAPTER XXII

CHRONIC RHEUMATIC HEART DISEASE

In northern climates rheumatism accounts for more chronic heart disease than any other single agent, and probably for not less than 40 per cent of all cases. This percentage varies greatly for different age periods. In children and in adolescents, once we exclude congenital malformations it is the only important cause. When a cause can be assigned it is rheumatism in the great majority of patients in the third and fourth decade of life. After the age of forty syphilis begins to take its place and soon becomes a prominent factor. After fifty is passed arteriosclerosis assumes an ever increasing importance.

A description of the established lesions or chronic forms of disease arising out of rheumatic infections, has been anticipated by the account of their development, and by the preliminary descriptions of structural lesions and functional disturbances of the earlier chapters of this book. In this section, therefore, it will be unnecessary to do more than outline the chief manifestations of the chronic rheumatic heart.

The patients may display valve defects, signs of cardiac enlargement, symptoms and signs of failure, irregular heart action, etc. It will be profitable to note the frequency with which many of these phenomena appear, and to examine, where it is useful to do so, their interrelationships, with a view to obtaining a simple basis for subsequent grouping.

CHIEF MANIFESTATIONS

Valve Disease—The frequency with which the valves are affected in the active stage of the disease, and the measure in which they become deformed in healing, have been described (pages 120, 129, and 193). As might be expected from their development, cases of rheumatic heart disease almost always display clear clinical evidence of valve disease, that is to say, they show signs of mitral stenosis, aortic re-

gurgitation, or aortic stenosis. Mitral stenosis is twice as common as aortic regurgitation and twenty times as common as aortic stenosis, aortic and mitral disease are not infrequently combined. Adult cases in which there are definite evidences of disease, such as enlargement of the heart, or fibrillation of the auricles, or congestion, without one of the named valve defects, are comparatively rare. This statement is made subject to the understanding that every care has been taken to recognise early cases of mitral stenosis (page 132). The comparatively rare cases form a group in which systolic apical murmurs may or may not be heard. Rheumatic patients in whom there are no signs of cardiac enlargement and no evidence other than a systolic apical murmur are more numerous, these have been referred to at an earlier stage (page 146). Here it may be well to comment again upon the significance of the systolic murmur in its present relation to rheumatic disease. First it may be said that such a systolic murmur appearing during the period of a rheumatic infection may form a guide to cardiac involvement (page 197), but experience shows that it has far less value as a sign left behind when active mischief has disappeared. A heart that is left with this sign only is one which at the most has been lightly involved, and there is actually no evidence to show that, taken together with the history of rheumatism, it has appreciably more significance than has rheumatism by itself.

Pericardial Adhesions—Many of these cases show after death simple obliteration of the pericardial sac, a condition of little consequence. Very occasionally the heart's action is embarrassed by the adhesions of mediastinopericarditis (page 176).

Enlargement—In an earlier chapter hypertrophy of the heart and of the right and left ventricles have been discussed in relation to the chief valve defects (page 106). In rheumatic heart disease signs of conspicuous enlargement are encountered chiefly in cases of aortic regurgitation, and usually but not always this is free regurgitation, comparable signs are found also in a few patients suffering from mitral stenosis, though in general the hearts in these present far less evidence of enlargement. Occasionally cases are seen in which conspicuous enlargement occurs in the absence of any sign of valve lesion or pericardial adhesion. The signs of cardiac enlargement vary very greatly from case to case, as judged clinically hearts of almost any size may be found in association with any of the valve defects. Enlargement in its various degrees runs more parallel with various degrees of failure than with the estimated and varying burdens of work.

Breathlessness and Failure—Breathlessness on effort is the dominant symptom of rheumatic heart cases. There are patients in whom exercise tolerance is good who nevertheless present aortic regurgitation or mitral stenosis and may or may not have signs of some enlargement of the heart; their number is limited. Chronic rheumatic heart cases show all grades of defective exercise tolerance; it is the symptom that ultimately becomes linked with signs of failure as the reserves are exhausted.

There is a general but imperfect relation between exercise tolerance and the size of the heart. Good exercise tolerance is almost confined to patients with little or no cardiac enlargement. Those with fair exercise tolerance mostly present definite or moderate rarely considerable enlargement. Patients with considerable enlargement are usually breathless with little or no effort; for the most part they show venous congestion or are near to doing so. Patients with congestion usually exhibit unequivocal signs of enlargement. The relation between failure and size, however, is not a strict one; thus it is broken by patients who while having small hearts are easily rendered breathless. The most notable of these are instances of mitral stenosis; breathlessness is more conspicuous in association with mitral stenosis than with aortic regurgitation. There are instances too of deep venous engorgement in patients who show few signs of enlargement. Relatively small hearts fail though not so frequently as do large ones. A small heart and venous congestion that is deep and long continued is an association suggesting encasement of the heart by fibrous bands or calcified pericardium (page 177).

Congestion is the usual termination in cases of rheumatic heart disease. Failure is due in part to the unusual burdens of work that the heart has to bear such as valve defects of various kinds and sometimes strong adhesions for these deplete reserves. But the chief factor nevertheless must be inherent in the muscle itself for failure can occur without the increased burden. And yet muscle that has failed cannot at present be recognised histologically or biochemically. There is increase of the interstitial fibrous tissue a little atrophy here and there a little fatty change; for the most part the fibres seem sound and the muscle presents no more change than is often found in hearts that have not failed. The change in the quality of the muscle is more subtle and frequently it is brought about by some recent infection from which the patient has suffered. Not only a rheumatic infection but infections of other types endanger or ter

minate the lives of those who are near to congestion or actually display it

Anginal Pain is experienced by some, these, if young, are nearly always cases of free aortic regurgitation with much cardiac enlargement. Cases of auricular fibrillation or of mitral stenosis rarely suffer from angina, but a few of the latter and a few cases of rheumatic heart disease without valvular trouble experience angina after passing the middle years of life

Precordial Ache, soreness, or other discomfort is a frequent minor complaint of rheumatic heart cases. Its origin is obscure

Palpitation is most frequently caused by overaction of the heart in aortic cases, and by irregular and rapid heart action in mitral stenosis

Auricular Fibrillation is one of the most serious consequences of rheumatic infection and plays an important part in the course of chronic rheumatic heart disease. That the relation to rheumatic disease is a very close one is shown by the fact that about two thirds of all cases of fibrillation belong to this group. Amongst mitral stenosis cases, and also amongst rheumatic cases that present no definite evidence of valve lesion, seen in adult outpatient departments, about a fifth exhibit auricular fibrillation. For a reason not as yet understood, auricular fibrillation is very unusual with aortic regurgitation occurring in less than one case in fifty of rheumatic origin; this statement is also true of cases in which aortic disease and mitral stenosis are combined

Apart from the past rheumatic history the reason for the frequent association of mitral stenosis and fibrillation is unknown. It would be incorrect to regard it as caused by mechanical strain in the auricle, for it can occur when the auricular muscle is hypertrophied or when it is largely atrophied. It cannot be ascribed with any confidence to such lesions as occur in the auricle, the muscle of which usually shows a good deal of fibrotic, atrophic, and fatty change, but not constantly. There is no known mechanical strain or muscular lesion in the cases of fibrillation which is not known to exist in cases of normal rhythm also. The ultimate cause of fibrillation is still unknown

Auricular fibrillation is prominent during a relatively late phase of rheumatic heart disease. It may occur at any time after infection, from the acute stage onwards, but its incidence begins to be heavy at about twenty five years and, as the disorder of mechanism is essentially persistent and fresh cases of it develop, the proportion of cases showing fibrillation mounts steadily with age, at least half

the cases of mitral stenosis reaching forty years are affected by it. Owing to its very frequent association with mitral stenosis, and to its baneful effects, the onset of fibrillation marks a chief event in the lives of a large proportion of these patients. There may be a preliminary period of paroxysmal fibrillation, but more usually fibrillation comes and, having come, stays to distinguish the last phases of the disease.

The effect of fibrillation upon the circulation has been considered (page 82). In rheumatic heart disease, fibrillation is usually associated with a high rate of ventricular response, owing probably to relative health of the junctional tissues. When it comes it therefore makes a heavy call upon the heart's reserves, and, as these are generally reduced and often much reduced, already, it almost always occasions much breathlessness, and often and quickly brings congestion. Failure with congestion, associated with auricular fibrillation, is the usual terminal picture, and the end in most cases would come quickly were it not for digitalis.

Paroxysmal Tachycardia and Auricular Flutter—The first of these is common and the latter is much rarer. They both favour cases of mitral stenosis with cardiac enlargement and at their onset have much the same effects on the circulation as has fibrillation of the auricles.

Embotic and Thrombotic Accidents—Embolism from the heart giving rise to infarction of the lung has been described fully (page 170). Simple emboli arise in similar circumstances from the left heart and find their way into the systemic system. As in the case of pulmonary embolism, the case is usually one of mitral stenosis and auricular fibrillation; the clot is formed in and detached from the left auricular appendix. The accident may happen when normal rhythm is restored by quinidine (page 91). Emboli are also derived from the apex of the left ventricle. These uninfected emboli may lodge like infected ones (page 188) in any artery, but the most important is the middle cerebral.

Very occasionally in mitral stenosis a large thrombus becomes detached within the left auricle. It lies loose as a ball in this chamber and may cause sudden death by plugging the mitral orifice.

Hemiplegia, with or without aphasia, is not infrequent as a sudden accident in mitral stenosis with fibrillation and in other forms of rheumatic heart disease. In these it is usually embolic. Hemiplegia occurs, of course, in other cardiovascular cases, thus it is not infrequent in arteriosclerosis and hypertension, but in these it more

often results from thrombosis or haemorrhage. Whether it arises from embolism or thrombosis, partial or considerable recovery is the rule, occasionally complete recovery happens within a few hours or days. Treatment of the paralysed limbs by massage and proper posture should begin after the first few days, if the patient's condition permits, and should be continued for a few weeks after improvement has ceased to be perceptible.

Thrombosis of veins occurs chiefly in cases of failure with congestion, the most frequent sites noted clinically being the veins of an arm or leg. The obstruction causes cyanosis and oedema, the latter often conspicuous, in the corresponding limb. The affected limb should be protected and kept at absolute rest for several weeks.

Acute Pulmonary Oedema, an occasional accident in mitral stenosis, and a less frequent one in aortic disease, has been described already (page 168)

OUTLINE OF TYPES, TREATMENT, AND PROGNOSIS

As has been seen, cases of chronic rheumatic heart disease present signs of valve disease of different forms, of cardiac enlargement in

nd symp
not group
of ways,
alike In

attempts to classify, one way would be to use valve defects as the basis, this is the time worn plan, it has never been very serviceable either from a clinical or pathological point of view. A classification by valve lesion is too anatomical, too little expresses functional ability or deficiency. To break up the cases into classes according to the size of the heart or to arrange them according to the amount of functional reserve, each of these would present certain advantages, and yet would insufficiently display the types. A practical subdivision uses neither an anatomical nor a simple functional basis, it adopts a natural classification based upon clinical experience. The groups here specified are not entirely comprehensive, neither are they sharply separable, but they enable the display of all the common forms that chronic rheumatic heart disease assumes, and they subdivide the patients into a few simple and useful categories from the standpoint of prognosis and treatment.

(a) *Valvular Disease the Chief Finding*—A not inconsiderable number of patients present signs of valvular disease, usually a

slight aortic regurgitation or a mitral stenosis but little or no evidence of enlargement of the heart and an exercise tolerance that is good or fair. They are cases that have come under observation for minor symptoms or in which physical signs of valvular disease are discovered unexpectedly. The outlook for these patients is in general good; most of them will live for very many years and live busy and even active lives taking exercise such as walking, riding, swimming, cycling, golf or engaging in manual work such as carpentering, fitting, plumbing, decorating. Many men of this class fought in the trenches in the Great War. The women do house work or in factories such work as packing or machine work; many of them marry and bear children without harm (page 268). The patients require little management. Little should be said to them about their hearts; they need occasional advice about their work and habits.

(b) *Auricular Fibrillation the Chief Finding*—The cases of this group come complaining when untreated of palpitation or breathlessness with brisk walking exercise; present few or no signs of cardiac enlargement and are almost always but not always found to display in addition to auricular fibrillation signs of mitral stenosis. This is the class of case in which the use of quinidine can be considered. Otherwise the ventricular rate is controlled by digitalis. Although it is remarkable how much many such patients do with impunity, the outlook in them is not so good and life should not be so strenuous as in the preceding class.

(c) *The Central and Chief Group*—These patients complain chiefly of breathlessness when they walk on the flat; tolerance of exercise being only fair or decidedly poor. The heart is found to be definitely or moderately enlarged and signs of aortic disease or mitral stenosis or both are discovered. In those without aortic regurgitation auricular fibrillation is often present.

This is the largest class of rheumatic heart case in adults. The prognosis is not good; there is a mortality of one third of the cases within ten years of diagnosis. Death when it comes is from cardiac failure in the majority. This is especially the case for mitral stenosis; infective endocarditis claims many of the aortic cases. Many of the group survive longer, however, and surviving these years show for the most part little change. In this group the death rate is equally great for mitral stenosis, for aortic regurgitation and for a combination of the two. Factors on which chief stress is to be laid are the degree of cardiac enlargement and the cardiac reserves. The patients lead or should lead very restricted lives being unfit for

more than light or sedentary work, or for any active exercise, the prospect of living is lengthened by living carefully. It is unwise of women in this class to bear children. In those who display fibrillation of the auricles, digitalis is to be used appropriately. All require close supervision, their symptoms should be a matter for frequent interrogation and test. The veins of the neck and the liver should be watched most carefully. As years go by, a number develop signs of failure, and usually do so rather abruptly. Most of the examples of infarction of the lung and of hemiplegia that are encountered come from this group.

(d) *Cardiac Enlargement the Chief Finding* —There are instances of considerable enlargement of the heart, without actual congestion, occurring chiefly in conjunction with free aortic regurgitation, but occasionally found apart from it. Almost without exception the cases have poor exercise tolerance, being breathless upon the slightest exertion. Amongst them also are cases of severe angina pectoris at rest, which have been described earlier (page 53). The patients rightly spend much or all of their time in bed, the prospect of life is but a few years, and is very short if the subjects are not well guarded.

(e) *Venous Congestion the Chief Feature* —When present, venous congestion dominates the picture controlling the outlook and governing treatment. Valve lesion, if present is but a subsidiary consideration. Congestion may be combined with aortic disease or with mitral stenosis. The latter is the more frequent combination, there may be no definite valve lesion to diagnose. It matters little. Congestion is associated with various degrees of enlargement, usually moderate or considerable. Because it reacts well to digitalis, the case of congestion with fibrillation is much more favourable than the case displaying normal rhythm. Chronic congestion that cannot be relieved is manifestly and immediately threatening. Cases of congestion even when responding to treatment, customarily relapse, and the prospect here is not in general long, though it varies by some years. Those patients who are relieved slowly, and those who, when relieved, present moderate or considerable enlargement of the heart, are least likely to survive.

This subdivision of chronic heart disease into a few appropriate categories sufficiently indicates the chief lines of treatment to be followed, as these are described in other parts of this book, and it begins to outline a plan of prognosis that proves satisfactory in practice, and that is dealt with more broadly in the final chapter.

In rheumatic heart cases that have become established the heaviest death rate is between the fifteenth and thirtieth year and it is heaviest in the early years of this period because these are the years of active disease. By the fortieth year less than half the patients survive. This statistical statement has only a limited value however in assessing the prospects of individual patients a number of whom live on to old age. Individual prospects are to be gauged on the lines laid down in the preceding classification.

INFECTION

In so far as rheumatic infection is concerned enough has been said in the preceding chapter to make it clear that it brings an element of great uncertainty into prognosis. In dealing with young patients still in their teens or entering their twenties the possibility of present rheumatic infection must still be considered but as the patient is older so search for and exclusion of signs of this infection becomes less important. The occurrence of subacute bacterial endocarditis cannot be foreseen. A proportion of these chronic rheumatic heart cases acquire this infection and when it is diagnosed prognosis changes correspondingly but as its coming is unpredictable and is infrequent it should not enter into the consideration of the prognosis of the uninfected individual.

DEGENERATIVE CHANGES

Patients who survive well into and beyond the forties enter the phase of life when degenerative changes appear. Arterial disease begins to take a prominent place perhaps the coronary arteries become sclerosed and bring in an anginal syndrome perhaps the arterial pressure becomes raised. These new troubles not infrequent in late rheumatic heart disease bring fresh complexity to the case.

CHAPTER XXIII

SYPHILIS OF HEART AND AORTA

AMONGST white people of northern climates syphilis is responsible for chronic disease of the heart and great vessels in approximately 10 to 20 per cent of all such cases. It is the cause of aortic regurgitation in about 30 to 40 per cent of all cases, its incidence, however, being closely associated with age thus between the ages thirty one and thirty five about 10 per cent, between the ages of forty one to forty five about 50 per cent, and between the ages of fifty one to fifty five about 70 per cent of these cases are syphilitic. Nine tenths of the patients are men.

Syphilitic aortic disease like *tabes dorsalis* is a very late manifestation of the original infection, it is the most frequent lethal consequence of this infection. The reason for the long latent period is unknown, but it averages about twenty years infection occurs usually in the twenties, and the peak in the incidence of syphilitic aortitis is near the forty fifth year. It may follow congenital infection, and then the latent period is in general shorter.

PATHOLOGICAL ANATOMY

In attacking the blood vessels syphilis concentrates upon the aorta at its origin, the ascending and the transverse aorta is usually involved in decreasing degree as it is traced onwards, to the mouths of its brachiocephalic branches. Less commonly the process invades the descending aorta, occasionally the abdominal aorta. The mouths of vessels issuing from a diseased aorta are always involved, but the branches of the aorta, in their length, escape, the chief though infrequent exception being the innominate artery. The disease often affects the sinuses of Valsalva heavily, here the mouths of the coronary arteries are implicated, the disease does not invade the vessels themselves, but ends abruptly at their origins.

The disease consists of a chronic inflammatory process, marked

by lymphocytic infiltration and fibroblastic reaction and by intimal thickening of local vasa vasorum. It probably starts in the adventitia and invades the wide media breaking and separating its elastic laminae. Intimal thickening occurs and the process of repair continues to lay down fibrous tissue here and also in and around the weakened aortic wall while the inflammation continues. The internal surface of the aorta is characteristically deformed by numerous large and small often depressed fibrous scars of white or pink colour. Atheromatous changes are mixed with these in a degree increasing with the age of the subject. The mouths of vessels issuing from the diseased aortic wall are much fibrosed and as shrinkage in these is less opposed by dilatation they are often greatly constricted a matter of supreme consequence in the instance of the coronary vessels and of diagnostic import in the case of larger branches.

As the affected wall of the aorta becomes deficient in resisting elastic laminae it stretches under the pressure of its contained blood. It may dilate more or less uniformly until its diameter is at least twice that of a normal aorta at a time not sharply distinguished in the process or by terminology dilatation becomes fusiform aneurysm. Expansion may proceed less regularly. A dilatation in one direction may become a definite bulge this may widen to form a small round projection or saccular aneurysm or one bulge may unite with others to form a much larger and irregular aneurysm. Aneurysms vary much in shape size position and in their direction of pointing. All structures that lie in the path of expansion are compressed distorted displaced or destroyed. Bone of rib or vertebra and other resistant structures are eroded. Saccular aneurysms are often lined and sometimes filled by laminated clot. This may be dense and solid or pierced by blood channels.

The disease attacks the aortic valve through its commissures which become inflamed the cusps become thickened and retracted. Simultaneously in greater or less degree the structures supporting the valve are weakened the valve ring dilates and the cusps shrunken as they are held away from the orifice which they were set to guard.

Hypertrophy of the heart is the rule in syphilitic aortitis it is conspicuous in cases in which aortic regurgitation has been free and then affects the left ventricle in general more than the right. Gummata are occasional at an earlier period of the infection and sometimes invade the auriculoventricular bundle (page 94). But in

the later stages active inflammatory lesions are small and rare. An increase of interstitial fibrous tissue is found, but changes in the muscle fibre are not pronounced.

In dealing with the clinical manifestations of syphilis of the heart and aorta, it will be convenient to consider the aorta first and the whole condition subsequently.

DILATATION AND ANEURYSM OF AORTA

SYMPTOMS

The complaints of patients suffering from dilatation or aneurysm of the thoracic aorta are extremely varied. By far the most frequent are pain and breathlessness. But as the aorta is syphilitic, the coronary mouths often narrowed and cardiac failure is present in greater or less degree these symptoms cannot be ascribed usually either to dilatation or to aneurysm.

Pain across the upper part of the chest may be frankly anginal, often it is less characteristic and then while sometimes to be ascribed to erosion of the sternum, its cause is more frequently obscure. Whether or not pain can be produced by the stretching or inflammation of nerves lying in the wall of an enlarged aorta is unknown.

Breathlessness is usually the result of cardiac failure but it may be contributed to, or caused by compression of trachea or bronchus or by compression of the lung and in other ways as well.

Most of the important symptoms of aneurysm are definitely produced by pressure on other structures. Thus pain comes from irritation of nerve roots or from the erosion of bone; difficulty in swallowing from pressure on the gullet. Haemoptysis comes from erosion and ulceration of the trachea, bronchus or lung, an actual leak from the aneurysmal sac itself being not unusually demonstrable. Cough may result from the same lesions or from infection of air passages or lung. These and other pressure symptoms will be noted again a little later.

LOCAL SIGNS

Pulsations —Several forms of pulsation occur in cases of aneurysm and form valuable signs.

Pulsation in the 2nd right intercostal space near the sternum, is the most frequent early sign of aneurysm of the ascending aorta. The sign occurs, however, much more commonly when the right lung retracts from the base of the heart, or when the heart is over

acting, especially in aortic cases with or without aortic dilatation. But in these instances pulsation is more visible than palpable, and it is a brief and not a sustained systolic movement as is that of aneurysm. When due to aneurysm the area that pulsates is always dull to percussion, but rarely so in simple dilatation of the aorta.

Pulsation is also caused by aneurysm in the 2nd or 3rd left intercostal space near the sternum, though less frequently. Local pulsation in this region much more usually results, however, from retraction of the left lung, dilatation of the pulmonary artery, or upward displacement of heart when the contents of the abdomen are greatly increased. Pulsation in the same spaces is common as part of a diffuse pulsation over the heart (page 109). In aneurysm alone there is a sustained systolic pulsation, dulness, however, may also accompany a dilated pulmonary artery.

Movement of the manubrium sterni, if systolic and sustained, is an almost diagnostic sign of aneurysm and is not infrequent.

Local pulsation from aneurysm may appear on occasion in almost any other part of the chest, front or back, and may affect rib or space, but it is infrequent in the back and on the front of the chest below the 4th rib.

Episternal pulsation, systolic in time, is frequent both in aneurysm and dilatation of the aorta. When the arch of the aorta enlarges upwards appreciably the manubrium tends to be displaced forwards, sometimes in aneurysms it is obvious that the trachea and sternum have become unusually separated, whether this is so or not, a finger pressed down behind the sternum may feel the top of a dilated or aneurysmal aorta.

Pulsating Tumour of the chest wall is a late sign of aneurysm. An aneurysm may break through the chest wall, eroding sternum, ribs, or cartilages in the process, at any of the places named in the last paragraphs. The site of election is the base of the heart, and especially to the right of the sternum. Such tumours bulge at first a little, then become hemispherical or conical as they grow. They are tender, and the skin covering them becomes inflamed. They are recognised for what they are almost always without any difficulty, since their form and site are characteristic, and because they pulsate. The pulsation is systolic and sustained, it is expansile—that is to say, it takes place at right angles to each part of the tumour's surface. Expansile pulsation is not to be expected if the tumour is only just raised above the level of the surrounding skin. It becomes obvious or conspicuous

when the tumour projects obviously or conspicuously. Thus the expansile quality of pulsation in these tumours is not very helpful because their nature is already evident before the sign is obtained. Very uncommon tumours of the chest wall may pulsate slightly and in an expansile fashion but the circumstances in which these occur are so different from those of aneurysm that they rarely should be confused. Pulsating empyema is easy to recognise occurring usually near the left nipple its pulsation is feeble and the fluid contained in the hernial projection of pleura returns to the main pleural sac under the light pressure of the hand.

Tracheal Tug —The aorta arches over the root of the left lung and when expanding unusually forces this root down and tugs on the trachea. The tug is felt if the tips of the two index fingers are placed under the lower margin of the cricoid cartilage and the cartilage gently lifted headwards. The tug when vigorous is diagnostic of aneurysm but tug that is distinct is also encountered in some cases of simple dilatation of free aortic regurgitation and occasionally in mediastinal new growth. Tug is downward and must not be confused with the forward impulse given by freely beating carotid arteries.

Dulness —Dilatation of the aorta usually produces dulness behind the sternum at the level of the 2nd and often up to the 1st interspace. The sign may be only just detectable or distinct. The borders of dulness may extend a half inch to the left and a little more to the right of the sternal borders. An area of dulness exceeding $2\frac{1}{2}$ inches (6 cms) in diameter very rarely results from simple dilatation and as the area is larger so aneurysm becomes more certain. Aneurysmal dulness as it increases extends mainly towards the right and downwards involving the 3rd and 4th as well as the 2nd and sometimes 1st spaces it also extends to the left though not usually so far. As in the case of pulsation dulness may appear over other parts of the chest wall.

Heart Sounds —Because enlargement of the thoracic aorta is generally accompanied by aortic regurgitation to and fro murmurs are usually audible at the base of the heart. When aortic regurgitation is absent or is slight a systolic murmur at the base (with or without thrill) is usual. An accentuation of the 2nd aortic sound is frequent the accentuation is peculiar in having a slight musical quality the noise is reproduced when a slack strip of linen is thrown into sudden tension. Though not diagnostic it is a useful guiding sign to aortic dilatation and is frequently associated with a clearly

palpable 2nd sound an equally suggestive sign termed diastolic shock.

The heart sounds are often remarkably well conducted to those parts of the chest wall towards which large aneurysms are projecting.

X ray Shadow—Much of the contour of the aorta can be determined by X ray exploration. In normal adults the right border of the cardiac silhouette is composed of two parts—namely, the rounded outline of the right auricle below and the almost straight shadow of the superior vena cava extending from this to the clavicle. This straight outline is often invaded by the slight convexity of the ascending aorta (see Fig. 21 page 108). The transverse aorta as it passes backwards through the mediastinum appears as a semi-circular projection or knob forming the top of the left border of the silhouette. The first part of the shadow below this knob may correspond to the beginning of the descending aorta or to the pulmonary artery, and is usually little curved. The left border is completed by the shadow of the ventricle. It is usually possible to obtain a clear idea of the contours of the ascending and transverse aorta by screening. The contours become easier to decipher in their whole length as the aorta becomes enlarged and information as to the size and shape of its several parts is increased by rotating the patient into a right or left oblique position. In orthodiagrams many measurements have been devised with the idea of expressing the size of the aorta in numerical terms but most of these are subject to error or misinterpretation and they do not add appreciably to a simple judgment from the orthodiagram of the form and area of the vessel.

When the aorta is dilated the density of its shadow is increased its curves become fuller but they retain their smoothness the ascending aorta takes a more prominent part in the right outline the knob is outlined by a fuller curve the knob may be raised until it meets the clavicle it extends farther to the left. Often the descending aorta is visible and descends for some distance before it is lost in the left border of the heart's silhouette. Thus the shadow of the basal vessels is widened at all levels (Figs. 33 and 34).

Aneurysms of the aorta present very diverse X ray pictures. Early aneurysms are usually recognised by closely following the visible contours of the aorta in appropriate planes so as to detect any disturbances of its normally continuous and rounded curves the direction of the line changes more or less abruptly where an aneurysm joins the aorta. In oblique radiograms or screening the

contours are examined for parallelism, departure from the cylindrical form is significant. When a definite tumour is present, its shadow if it is aneurysmal, must blend with and present a similar density to that of the remainder of the aortic silhouette. Aneurysms (Figs 35 to 38) present sharply defined margins for the most part smooth contours, and no irregularities of density. Sharply defined circular outlines are particularly suggestive, as they can only be confused with cysts. The presence or absence of pulsation at the border of the shadow is rarely helpful because aneurysms often fail to display it, because the normal aorta may project beyond a solid tumour, and because tumours often have pulsation transmitted to them. Expansile pulsation in a round tumour is of course decisive, but this is not often seen. X ray signs in trachea and oesophagus are described later.

SYMPTOMS AND SIGNS OF PRESSURE

Owing to the anatomical relations of the aorta to the mediastinum aneurysm of this vessel frequently causes symptoms and signs by exerting pressure upon structures surrounding it. Pressure manifestations are usually the result of aneurysm of the transverse aorta, and the chief are as follows —

The oesophagus is one of the most frequent structures to be affected many patients complain of difficulty in swallowing solid food. Very few are unable to swallow and few regurgitate a point of importance in differentiating from other forms of stricture. If a mouthful of barium paste is swallowed while the patient is being screened an actual obstruction to the oesophagus is readily demonstrable. The obstruction is usually level with the top of the arch of the aorta or just below it. A noticeable lateral bend or displacement of the oesophagus to the right may be noted when the patient faces the screen, but this is usually more distinct when the subject turns half left with the right shoulder forward. Minor grades of interference, such as bending or kinking of the oesophagus though insufficient to cause complaint, are frequent in cases of early aneurysm and of simple dilatation of the arch.

Compression of the lung by aneurysm is observed most frequently at the right apex, and results in dulness, deficient air entry, and crepitations. Occasionally the lung is eroded giving rise to haemoptysis.

Pressure on the trachea may be responsible for breathlessness or stridor and when the aneurysm ulcerates the trachea, for cough

with blood streaked sputum, or heavier haemorrhage. The most frequent sign in the trachea apart from tug is its slight but distinct displacement to the left at the root of the neck, this may be detected at the bedside and is well seen in many of the skiagrams. Deformity of the trachea is responsible also for metallic (brassy) cough.

Pressure on the left bronchus causes at first deficient air entry at the left base characteristically it causes absent breath sounds over lung that is resonant to percussion, later, bronchiectasis and local pulmonary infections may follow.

Compression of the left recurrent laryngeal nerve causes an




FIG 33—L rhodiogram. Man aged 4 weight 150 lb. Aortic regurgitation (free) syphilitic aortitis: a general dilatation and elongation of the aorta. The ascending transverse and descending parts of the aorta are all clearly visible.




FIG 34—Orthodiagram. Man aged 60 weight 150 lb. Aortic regurgitation (slight) syphilitic aortitis: systolic blood pressure 240 mm. Great widening of the ascending aorta and of its arch. The ventricle is enlarged and lies horizontally.

abductor, and later complete, paralysis of the corresponding vocal cord changing the voice, making the voice hoarse and the breathing noisy.

When the left sympathetic nerve is caught and paralysed the left eyeball sinks, the palpebral fissure and the pupil are narrow, signs that are rendered relatively more emphatic by instilling a few drops of 1 per cent cocaine into each conjunctival sac, for this dilates the pupil and renders the eyeball more prominent on the unaffected side. Sympathetic paralysis causes increased warmth of the face on the same side, the colour of the affected skin may be unchanged, or it may be redder or paler than that of the other side according to the state of vasomotor tone in the latter. The most frequent of sympathetic signs is inequality of the pupils.

Obstruction of the superior cava causes swelling, most notably in the face, and high colour in, and often cyanosis of, the head and neck, the veins of the neck are engorged but do not pulsate. If the




FIG 35—Orthodiagram. Man aged 56 weight 151 lb. Aortic regurgitation (free) syphilitic aortitis fusiform aneurysm of ascending aorta the descending aorta also visible




FIG 36—Orthodiagram. Man aged 45 weight 139 lb. Aortic regurgitation (free) syphilitic aortitis a conical aneurysm projects to the right from the ascending aorta and the transverse aorta bulges as a sac to the left in the chest




FIG 37—Orthodiagram. Man aged 68 weight 167 lb. A sacular aneurysm (S) of the ascending aorta bulges to the right. The arch of the aorta is greatly dilated or actually aneurysmal




FIG 38—Orthodiagram. Man aged 80. The ascending aorta and arch are dilated and an aneurysm bulges to the left from the first part of the descending aorta.

patient survives long enough, anastomotic veins appear over the clavicle and in the region of the shoulder. The blood flow in them can be traced downwards over the front of the chest or axillae. Ultimately the anastomoses may form very dilated tortuous leashes.

Pressure on a subclavian vein engorges the corresponding arm, and on a subclavian artery weakens the pulse beyond. Occasionally

lower roots of the brachial plexus are damaged, causing neuralgia in the arm

Descending Aorta—Aneurysm of the descending aorta is apt to develop very insidiously and to give neither symptom nor sign till it has grown to a large size. Obstruction of the oesophagus may occur as an early isolated event. Compression of lung is frequent. The aneurysm erodes the bodies of the vertebrae, may involve the dorsal nerve roots, and in more advanced cases may even come to press directly upon the spinal cord. Erosion may be painless, but a steady aching pain localised in the middle line of the back is the commonest symptom. It is sometimes accompanied by local tenderness. The patient may complain when the spine is jarred and use great caution in rising from a lying to a sitting position. Pain may be more severe and radiate along the dorsal nerves. Pressure on the cord produces the familiar picture of paraplegia. Dulness to percussion and even pulsation may appear locally, but are rare signs. A skiagram will demonstrate the state of the vertebrae and reveal the tumour.

DISTANT ARTERIAL SIGNS

Most valuable evidence of the involvement of main arteries by disease is often to be obtained from their branches. Inequality in the strength of the radial pulses is frequent in aneurysm of the thoracic aorta. When found it is not to be used until traced to its source. First the brachial and then the subclavian, pulses are felt to ascertain if these participate in the inequality. Inequality of the pulses at the wrist is quite frequently due to difference in the size of the two radial arteries, or to one vessel turning outside the radius unusually high in the forearm. If, as is usual, the right subclavian is weaker than the left, then it is important to ascertain if the right carotid is the less distinct, for in that case the innominate artery is obstructed. It is unusual to be able to state that more than one of the three main vessels issuing from the transverse aorta is involved, though all three may be.

When decreased the pulse is anacrotic (page 127) or flat topped in form. The up stroke of the pulse is never delayed, but, if the pulse is anacrotic on one side, then its summit is represented by the second wave of the pulse, and this may give the impression of delay. There may be but a flicker in the arteries of one arm, or no pulse may be felt. Weakness of the brachial pulse is associated with a corresponding fall of the systolic pressure reading on that side. It is

erroneous to believe that diminished pulsation in the arteries necessarily means a lessened circulation in the limb in aneurysm gross inequality of the radial pulses may be found yet the hands may be of equal warmth and measurement may show the blood flow in them to be equal in amount

The cause of a decreased pulse in aneurysm is obstruction at the mouth of the corresponding vessel as it leaves the sac This is usually the result of fibrosis and constriction of the mouth as part of the disease in the aortic wall similar constrictions occur in syphilitic aortitis without aneurysm though in these they are less common because the disease is less advanced I have known the pulses in all the three vessels to be lost in a simple syphilitic aortitis A less usual cause ■ outgoing blocking of the mouth by clot lying in the aneurysmal sac it is rare for an aneurysmal sac to press upon a subclavian artery sufficiently to distort it That weakening of the pulse is materially contributed to by the wastage of its force in expanding the sac is a prevalent but erroneous notion If it were so the femoral pulses would often show more weakness than they do

CARDIOAORTIC SYPHILIS

From considering in detail the diverse effects of syphilitic aortitis attention is turned to the disease as it affects heart and aorta together

In making its delayed attack syphilis concentrates upon the aorta Of symptoms directly referable to inflammation of the aorta none are known a dull ache across the sternum is an early and frequent symptom sometimes ascribed to this cause The disease usually develops far before the patient complains and the complaint when it comes arises out of one of four chief events The disease (a) weakens the aortic wall so that this dilates or becomes aneurysmal (b) invades the aortic cusps and their supporting structures rendering the valve incompetent (c) attacks the mouths of the coronary vessels and causes angina pectoris (d) produces directly or indirectly a decline of ventricular efficiency Of these four chief results the first with its pressure symptoms has been considered at length The remainder have been considered at length though not with particular reference to syphilis in previous chapters The chief manifestations vary in their incidence or are combined in different ways But when one is displayed others are not often long delayed

CHIEF MANIFESTATIONS

Dilatation of the Aorta distinctly recognizable is found in at least half the cases *aneurysm* occurs in an additional group amounting to 15 or 30 per cent of the whole

Aortic Regurgitation is present in the great majority of the cases, whether *aneurysm* is present or not When present it is almost always free and extreme signs of regurgitation are frequent Slight regurgitation is much less common than in rheumatic heart disease stenosis of the valve is a rarity The reason for this difference is that in syphilis the aortic ring is dilated and the cusps lose their support Rupture of an aortic cusp is usually the result of syphilis and gives rise often to a particular group of signs and symptoms already described (page 125)

Angina Pectoris of unmistakable type occurs in a number of the cases The anginal subjects nearly always present aortic regurgitation Usually appearing as an angina of effort it often reaches ultimately an unusual pitch of intensity attacks of great severity occurring once or several times a day the attacks are frequently accompanied by a conspicuous rise of pulse rate and blood pressure (page 53)

Cardiac Failure —In cardioaortic syphilis breathlessness on effort is an almost invariable symptom and in a large proportion of the cases signs of congestion appear The two chief known reasons for this failure are the increased work that the heart has to do and insufficiency of its blood supply The extra work is due to the aortic reflux and here allowance must be made not only for the advanced grades of disease in the cusps but for the increase in the size of the orifice that is to be guarded moreover these patients are middle aged or elderly and high or relatively high blood pressures are common amongst them Insufficient blood supply results in part from constriction of the coronary mouths and in part also from the heaviness of the ventricles that are to be supplied through them Enlargement of the heart is conspicuous or extreme in very many of these patients it is particularly so in those with aortic regurgitation Enlargement however, and sometimes great enlargement occurs when there is no incompetence of the aortic valve In general as in the case of rheumatic heart disease the degree of breathlessness corresponds with the amount of enlargement Little can be said of structural change in the ventricular muscle There is a certain amount of replacement fibrosis it is true,

but the cause of failure cannot be recognised histologically here, any more than in rheumatic heart disease. Experience shows that the muscle of an enlarged heart is rarely dependable. Alternation of the pulse (page 248) is not infrequent. In elderly subjects atheromatous changes occur in the coronary vessels and induce secondary changes in the muscle, but coronary thrombosis is not a common accident in these patients. Neither does fibrillation of the auricles often contribute to failure, failure with normal rhythm is the rule.

As in other instances in which failure with congestion comes while the rhythm is normal, it is almost always progressive and terminates life within six or twelve months. As in similar but non syphilitic cases, it is frequently accompanied by Cheyne Stokes' breathing and by distressing paroxysms of breathlessness, experienced especially at night, and more closely described on page 160.

TYPES

The chief types that may be recognised conveniently are — a main group in which there is aortic regurgitation, with or without dilatation of the aorta, but no angina; a smaller group in which anginal pain dominates the clinical picture, and a third group of cases presenting aneurysm. Each of these types may or may not present signs of venous congestion. A fourth group is small because the cases are usually difficult to diagnose. It is one in which there is syphilitic aortitis, with or without dilatation of the aorta and enlargement of the heart, but no aortic reflux. It has practical importance because in it are represented many of the early cases. It has theoretical interest because the patients sometimes develop cardiac enlargement, even great enlargement, and congestion, while the aortic valves still remain competent.

DIAGNOSIS

In diagnosing syphilitic aortitis, it is necessary to obtain evidence of syphilitic infection, and to identify disease of the aorta. The very presence of aortic regurgitation in a man of forty five years at once suggests syphilitic aortitis, and its presence in older men makes its origin from syphilis increasingly probable. Given a history of the infection, or scars recognisable as those of gummata of skin or palate, and the diagnosis is in little doubt. A positive Wassermann reaction or association with *tabes dorsalis* is more valuable still. A negative Wassermann reaction, on the other hand, does not exclude syphilis as the cause.

A middle aged or elderly man who presents signs of aortic re

gurgitation should always be searched closely for the clinical signs of aortic dilatation and of aneurysm. And this examination should include whenever possible a thorough examination of the chest under X rays. For if definite evidence of aortic dilatation is found the diagnosis of syphilitic aortitis becomes highly probable and if signs of aneurysm are found it becomes almost certain. Dilatation of the aorta however may occasionally produce retrosternal dulness to percussion in cases of senile atheroma especially when accompanied by high arterial tension though this is usually distinguishable by the tortuosity of the aorta under the X rays. Dilatation and occasionally aneurysm of the aorta occur in connection of the aorta this condition is relatively rare and not difficult to distinguish (page 263).

The diagnosis of syphilitic aortitis when aortic regurgitation is absent is more difficult. It is well to consider syphilis as the possible origin of enlargement of the heart in all patients over forty years of age in which the cause of enlargement is obscure. A positive Wassermann in these however does not suffice since its association may be fortuitous. To make the diagnosis certain enlargement of the aorta must be found as well.

The diagnosis of aneurysm in its early stages depends mainly upon the closeness with which patients prone to display this form of aortic disease are searched and especially upon judiciously selecting cases for X ray examination. Most cases should be diagnosed before the patient complains of some symptom more or less distinctive of aneurysm and before there are manifest signs over the front of the chest. The diagnosis should hardly ever be made in patients under thirty years. When a sign or symptom of pressure upon a mediastinal organ appears it will be necessary to differentiate between aneurysm and new growth. A positive Wassermann reaction will assist in this case signs of aortic regurgitation will assist even more because they will proclaim the aorta unhealthy but in the absence of such assistance the diagnosis of aneurysm will turn chiefly upon an X ray examination upon finding that an artery leading from the aorta is obstructed or upon the general grouping of symptoms and signs.

COURSE AND PROGNOSIS

Syphilitic aortitis is a malady of later years when the natural expectation of life is not very lengthy. It is a serious disease and usually comes under observation when far advanced. Consequently the outlook in years is relatively short. The average duration of life from the diagnosis is about six years in patients under treatment though

quite a number live ten years or more. A *compensated* case may suddenly in another large group of the cases *develop*, and this then progresses to end life *usually* a few die of acute oedema of the lungs. In a *small* infective endocarditis supervenes. A few die of *cerebral* haemorrhage, or as a consequence of *important* structures. Intercurrent *disease*, account for many more deaths.

In the individual case serious anginal *influence* the outlook unfavourably. The *anginal* severe and rapidly progressive owing to further *mouths* of the coronary vessels. It is in these *aneurysm* that sudden death is so frequent. In *death* results from rupture of the sac externally, *bronchus*, oesophagus, pleura or pericardium. The *in* cases of aneurysm is about four years from the *an aneurysm* has bled or by compression has *lungs*, the course will be measured in months. But *not* invariably unfavourable, some aneurysms *the* condition remains unchanged for eight to *disease* discovered.

Apart from these special considerations, *turns* mainly upon the degree of failure, as *and* signs, and upon the size of the heart, much *early* signs of venous congestion or great *being* ominous.

The prognosis should take into account *the* reaction of the patient to rest, the *patients* under full treatment and not at work *that* the prognosis is much less favourable in *do* manual work especially heavy work, which *and* strains the unhealthy aorta.

TREATMENT

The chief remedial measure in the *in* all its forms is rest. Manual work, unless of *forbidden*, even in the mildest cases, and *enlargement* of the heart, angina, or aneurysm *or* all of their time lying or sitting. Diet *page* 286) Strenuous acts (page 285) are to *and* strictly healthy habits (page 283) *measures*.

(15 or 20 grains thrice daily) has proved to be a useful remedy in relieving pain combined with mercury in the form of grey powders or unguents, it is used as a but is not proved to stay the remedies care should be taken

Neosalvarsan and similar preparations have been employed extensively from the same point of view and under full courses the Wassermann reaction frequently becomes negative Ten to fifteen intravenous injections of neosalvarsan of 5 grains to test and later 10 grains (0.3 and 0.6 gramme) are given at the rate of one or two doses a week, and the course repeated twice each year The uncomplicated aortic cases tolerate these injections well According to most available evidence this remedy prolongs life, though this is not universally agreed Cases having angina should not be given this treatment, but early aneurysmal cases may

In treating thoracic aneurysm a chief point is continuously to maintain a low average blood pressure and this is most safely accomplished by rest, by a small but adequate dietary, and by the promotion of full hours of sleep Vasodilator substances are not recommended It is not unusual to notice a decline in the size of a pointing aneurysm as blood pressure falls with rest The use of very low calorific diets and of other remedies such as the introduction of foreign bodies into the aneurysmal sac, having for their object the promotion of clotting in the sac, is not advised It is quite beyond our power to control the amount of clotting in the sac, and the remedies have each their special dangers, it is clear that clot may form in quantity and yet fail to strengthen the aneurysm against rupture, and it is clear that clots in forming may obstruct important outgoing vessels and even the aorta itself Remedies of this class do more harm than good When an aneurysm is weeping into trachea into oesophagus or externally, as it sometimes does before rupturing, the patient must be kept at absolute rest under morphia

Angina and failure with congestion, when present, are treated on lines already laid down

The time to treat syphilis is during the weeks following infection Once the disease has declared itself in the aorta the damage is usually irreparable¹

¹ In the United States of America the death rate amongst cases of syphilitic aortitis appears to be heavier than in Great Britain, upon which the prognosis here given is based.

(15 or 20 grains thrice daily) has proved to be a useful remedy in relieving pain combined with mercury in the form of grey powders or inunctions it is used as a specific remedy and has been thought, but is not proved to stay the progress of the disease. In using these remedies care should be taken to avoid iodism or mercurialism.

Neosalvarsan and similar preparations have been employed extensively from the same point of view, and under full courses the Wassermann reaction frequently becomes negative. Ten to fifteen intravenous injections of neosalvarsan of 5 grains to test and later 10 grains (0.3 and 0.6 gramme) are given at the rate of one or two doses a week and the course repeated twice each year. The uncomplicated aortic cases tolerate these injections well. According to most available evidence this remedy prolongs life, though this is not universally agreed. Cases having angina should not be given this treatment but early aneurysmal cases may.

In treating thoracic aneurysm a chief point is continuously to maintain a low average blood pressure, and this is most safely accomplished by rest by a small but adequate dietary, and by the promotion of full hours of sleep. Vasodilator substances are not recommended. It is not unusual to notice a decline in the size of a pointing aneurysm as blood pressure falls with rest. The use of very low caloric diets and of other remedies, such as the introduction of foreign bodies into the aneurysmal sac, having for their object the promotion of clotting in the sac, is not advised. It is quite beyond our power to control the amount of clotting in the sac, and the remedies have each their special dangers, it is clear that clot may form in quantity and yet fail to strengthen the aneurysm against rupture, and it is clear that clots in forming may obstruct important outgoing vessels and even the aorta itself. Remedies of this class do more harm than good. When an aneurysm is weeping into trachea, into oesophagus, or externally, as it sometimes does before rupturing, the patient must be kept at absolute rest under morphia.

Angina and failure with congestion, when present, are treated on lines already laid down.

The time to treat syphilis is during the weeks following infection. Once the disease has declared itself in the aorta the damage is usually irreparable.¹

¹ In the United States of America the death rate amongst cases of syphilitic aortitis appears to be heavier than in Great Britain upon which the prognosis here given is based.

Cardiac symptoms are the commonest. Palpitation is frequent and may be due to forcible action of a regularly beating heart but is more usually ascribable to extrasystoles and in occasional cases to auricular fibrillation. The most frequent symptom belongs to cardiac failure namely breathlessness on effort which is mild in the early cases and increases in severity until congestion comes. A little swelling of the feet is not infrequent as in other elderly people and is not necessarily due to cardiac failure (page 20). In advanced cases nocturnal attacks of cardiac asthma (page 166) may be frequent and very distressing. Anginal pain occurring on exercise and ultimately at rest is the chief complaint of a number.

RECOGNITION OF HIGH BLOOD PRESSURE

Accentuation of the 2nd Sound —It is widely believed that variation in the sound at the 2nd right cartilage means abnormality in the systemic and at the left cartilage in the pulmonary vascular system and accentuation of the 2nd sound has long been used and more rigidly than is warranted as evidence of raised pressure in the corresponding artery. It is quite true that an accentuated second sound at the 2nd right cartilage is often heard in cases of high blood pressure but such accentuation is not infrequent when blood pressure is unraised and conversely high blood pressure is not infrequent with the sound at or below its usual intensity. It is to be remembered that it is not systolic pressure that is concerned in closing the valve and rendering it tense but aortic pressure at the very end of systole and that vibration of the valve is governed equally if not actually to a greater extent by the rate at which the pressure falls away in the ventricles at the end of systole. Thus while an accentuated 2nd sound may usefully draw attention to raised blood pressure it is illegitimate to use it as valid evidence of high arterial tension. Similarly the conclusion that pulmonary arterial pressure is raised cannot be based on an accentuated 2nd sound at the left cartilage.

Pulse Tension —The fingers should be trained to recognise a high tension pulse for a pressure meter can hardly be used as a routine in general practice. The experienced will usually suspect high pressure where it occurs as soon as the radial pulse is felt. To make more certain the index finger of the left hand should feel the pulse while the index of the right presses proximally upon the artery to obliterate it. The estimate of tension should never be attempted with the fingers of one hand only. The amount of pressure that must

be exerted to obliterate forms the gauge. A recurrent pulse will only occasionally make it essential to stop the artery low down as well.

Sphygmomanometry —An accurate gauge of pressure at the top of the pulse beat (systolic pressure) in the brachial artery can be obtained by means of special instruments that are familiar. There are many forms and most are sound. A few essentials must be named. The gauge of pressure if not a mercurial column should be checked against the latter from time to time. The outer surface of the cuff must be inextensible (leather or silk) the rubber bag for use on arm only must have a width not less than 12 cms and must itself be long enough to encircle the limb completely, otherwise serious over estimates of pressure become inevitable. For very obese arms and for the leg larger cuffs should be used. In taking systolic pressure the cuff pressure at which the beats begin to reappear at the wrist is in general just as good a guide as is the return of the sounds over the artery below the cuff. There are small errors and occasional fallacies with each method. As pressure in the cuff gradually declines the sounds below the cuff become louder and later diminish in intensity these changes may happen once or twice during the decline. The pressure at which the sounds begin to fall away abruptly and finally is the pressure usually taken as the index of pressure prevailing at the beginning of the pulse beat (diastolic pressure). Diverse opinions are held about the correct index of diastolic pressure and of the value of these readings when obtained. The gauges are largely theoretical. It is probable that normal readings are sufficiently close but the accuracy of many abnormal readings and especially low readings is less certain. The usual range of normal diastolic pressure readings is about 70 to 85 mms. A pressure of 100 mms is generally regarded as definitely abnormal. Mean or average pressure which would express what the heart has to maintain is not measurable in man but it lies nearer to diastolic than to systolic pressure. Other things being equal when the pulse slows the systolic pressure mounts and diastolic pressure falls.

In using systolic pressure fine distinctions of height are useless for the pressure varies much and frequently with pulse rate with emotion exercise and many other causes normally a difference of 10 to 15 mms may be found in the pressure in the two arms. A single reading has little value unless low or unless exceptionally high for previous exercise or present excitement will often lift the pressure to a high point in susceptible people (to 160 or 200 mms Hg).

In adults repeated readings, taken in circumstances of physical and mental rest, show normal pressures of 110 to 130 mms usually. These are definitely normal values, but some would extend these limits in older subjects. Values that are above 160 mms Hg definitely surpass normal limits. Such readings are rightly regarded as less negligible in a young than in an elderly subject, because they are less usual. The average blood pressure rises with age, this is not to say that rising blood pressure is normal, for many vigorous and long lived men retain low blood pressure, it is good that an otherwise healthy but aged man should show a pressure below the average for his years. Resting pressures of 180 mms and over are universally regarded as well beyond normality at any age, though pressures often pass 200 and readings up to 300 or even 350 mms are found.

In using systolic pressure to gauge whether the condition essential hypertension is present or not, it is to be remembered that, when the pulse pressure (or difference between systolic and diastolic pressures) is large as in aortic regurgitation, or when the ventricular action is very slow as in complete heart block, the systolic pressure rises 10 to 20 mms or more higher than it otherwise would.

Often in cases of irregular heart action, and especially in auricular fibrillation, the beats pass the cuff at widely different pressures. The cuff pressure that allows the occasional large beats to pass, while most are just obstructed, will form a useful, though it is not an accurate, gauge.

RECOGNITION OF ARTERIOLAR SPASM

If there are any physical signs of arteriolar spasm that can be recognised clinically these are to be found in the retinal vessels, which are the only vessels of suitable size that can be directly inspected. These arteries or arterioles are often narrowed remarkably.

ACCOMPANYING MANIFESTATIONS

Many of the patients are heavily built or obese, high coloured, and becoming turgid in face when they stoop. But there is no rule. High pressure is also found in small people, and many lose flesh and become thin and anaemic in the later stages of the malady.

Vascular Manifestations—In addition to the signs of high blood pressure already detailed are those of general arterial disease (page 241). Some disease of the arteries is invariable, often it is advanced.

The rupture of small vessels is not uncommon, petechiae are

be exerted to obliterate forms the gauge. A recurrent pulse will only occasionally make it essential to stop the artery low down as well.

Sphygmomanometry —An accurate gauge of pressure at the top of the pulse beat (systolic pressure) in the brachial artery can be obtained by means of special instruments that are familiar. There are many forms and most are sound. A few essentials must be named. The gauge of pressure, if not a mercurial column should be checked against the latter from time to time. The outer surface of the cuff must be inextensible (leather or silk) the rubber bag for use on arm only, must have a width not less than 12 cms and must itself be long enough to encircle the limb completely otherwise serious over estimates of pressure become inevitable. For very obese arms, and for the leg, larger cuffs should be used. In taking systolic pressure, the cuff pressure at which the beats begin to reappear at the wrist is in general just as good a guide as is the return of the sounds over the artery below the cuff. There are small errors and occasional fallacies with each method. A

the sounds begin to fall away abruptly and finally the pressure usually taken as the index of pressure prevailing at the beginning of the pulse beat (diastolic pressure). Diverse opinions are held about the correct index of diastolic pressure and of the value of these readings when obtained. The gauges are largely theoretical. It is probable that normal readings are sufficiently close, but the accuracy of many abnormal readings, and especially low readings is less certain. The usual range of normal diastolic pressure readings is about 70 to 85 mm. A pressure of 100 mm is generally regarded as definitely abnormal. Mean, or average pressure, which would express what the heart has to maintain, is not measurable in man, but it lies nearer to diastolic than to systolic pressure. Other things being equal, when the pulse slows, the systolic pressure mounts and diastolic pressure falls.

In using systolic pressure fine distinctions of height are useless for the pressure varies much and frequently with pulse rate with emotion, exercise and many other causes, normally a difference of 10 to 15 mm may be found in the pressure in the two arms. A single reading has little value unless low, or unless exceptionally high, for previous exercise or present excitement will often lift the pressure to a high point in susceptible people (to 160 or 200 mm Hg).

may also appear Uraemia is very occasional Oedema of renal type does not occur

Inflammation—These patients are unusually prone to develop inflammations, such as pneumonia, pleurisy and pericarditis, and infections of skin and subcutaneous tissues

A FINAL STATE

Breathlessness is an unusually conspicuous feature of essential hypertension in its later phases, and recognition of this fact is important In the failure of rheumatic heart disease, breathlessness is not very prominent in patients lying quietly in bed unless the veins are very deeply congested When the veins are moderately swollen, respiration is a little deepened and its rate is perhaps 25 or near this In the failure of hypertension venous congestion of this depth is usually accompanied by much greater breathlessness The breach of the usual relationship is important clinically because, if it is quickly observed, it will very often lead almost at once to a correct diagnosis Although orthopnoea is the rule in such patients there are cases in which the patient prefers a recumbent position and in these there may be urgent dyspnoea but no sign of venous congestion, this, however, is quite exceptional When there is slight or moderate congestion and unusual breathlessness depth being increased, and the rate passing above 30 to 35 or 40 per minute a simple diagnosis of congestive failure will not suffice If bronchitis is the cause, this will be evident, otherwise it will usually transpire that the disease is hypertensive failure The point is the more important diagnostically because at this stage other signs may fail Thus the quantity of

g
t

sometimes the fall comes earlier Thus blood pressure cannot always be relied upon to decide the diagnosis

Cheyne Stokes breathing is present in a large proportion of the patients at this stage, though it may consist of inconspicuous waxing and waning of the breathing In such it must be watched for closely to be discovered. The periodic breathing is most emphatic during sleep When apnoea comes, the patient becomes drowsy and is difficult to rouse, in the true hyperpnoeic phase he is restless and on occasion much distressed

Apathy to surroundings, a state ultimately bordering upon or reaching delirium, a pulse which if regular is rapid (100 to 120), a

temperature chart that has for weeks been subnormal in a man often wasted and sallow combine with the respiratory disturbances described to form a characteristic picture that is often seen

DIFFERENTIATION OF ESSENTIAL HYPERTENSION FROM CHRONIC NEPHRITIS

To distinguish between chronic glomerulo nephritis (secondarily contracted kidney) and the much commoner essential hypertension is not always possible clinically. A history of acute nephritis decisively favours the former. Essential hypertension is rare before and chronic nephritis rather rare after the fortieth year. The higher blood pressures (250 and over) are rare except in essential hypertension. In nephritis the pressures are more stable than in essential hypertension in which they tend to fall with rest and vary in other circumstances. Albumen occurs freely and red blood cells are found in the urine in nephritis. In nephritis retinitis is much commoner and takes severer forms and signs of renal insufficiency are more usual. Death is usually from uraemia.

COURSE AND PROGNOSIS

Most cases of essential hypertension present little change in health from year to year but continue to enjoy fair health or still complain of the minor symptoms which first brought them under observation. The duration of life from the onset of symptoms is variable but is now considered to average about ten years. The onset is usually so insidious however that the total duration of the malady is represented by a longer period probably double that already stated. Many patients are now known to have maintained high blood pressure for as long as twenty and some for as long as thirty years. The average duration of life should be considered in relation to the fact that the onset of symptoms is heaviest between the fiftieth and seventieth year and in relation to the natural expectation of life at these ages. Nevertheless the death rate amongst high tension cases is decidedly above normal. High tension determines the cause of death in at least half those who suffer from it. About one in three (or rather more) dies of heart failure, about one in four or five of cerebral haemorrhage, a few die in the anginal state, a very few develop grave renal insufficiency and become uraemic.

The prognosis for the individual is far less gloomy than it was at first thought to be by the profession, it is far brighter than it is still thought to be by the educated public. Distinct elements of un

certainly there are such as the cerebral accidents but it is not right that prognosis should be dominated by these though they naturally call for reserve in forecasting

Statistics show that the expectation of life shortens as the blood pressure is higher. Thus the death rate is very decidedly higher in cases in which the pressure ranges above than in cases in which it ranges below 200 mms Hg. Naturally little emphasis will be placed on pressures that approach to normal limits but in the usual higher ranges (200 mms or over) the actual height cannot be used to any great advantage the exception being extreme pressures such as 300 mms which are not consistent with many years of life and are not often displayed except by those obviously and very seriously ill.

The chief consideration in individual prognosis is the state of the heart. The outlook is brightest when high tension is found accidentally as in insurance examinations and the patient is leading an active life untroubled by symptoms and presenting few or no cardiac signs. It is kindly and legitimate in speaking of the future to name such periods as ten or twenty years in the hearing of anxious patients according to their ages. The naming of such periods will mean more to an older than to a younger man but should not be extended in the latter case. The outlook is a little less promising in those who become unduly short of breath for their years on active exercise. It is definitely less promising in those who are often breathless after moderate exercise or suffer occasional anginal pain on effort. It is dis-

of the feet should not be used by itself elderly and especially obese subjects often display it when the heart is sound. Its presence should be judged by the state of the veins and liver. Recovery from failure

ness to be the last

Varying degrees of enlargement of the heart are often paired with

appropriate degrees of failure. Great enlargement is unfavourable. Abnormality of the electrocardiogram such as curves of bundle branch block or other high grade anomalies and continuous pulsus alternans are other signs of grave cardiac involvement which even in the absence of congestion indicate a brief course.

Cerebral haemorrhage introduces the chief element of uncertainty into the prognosis the possibility of coronary thrombosis occurring also contributes a little. Though these factors naturally render dogmatic promises unwise they should not be introduced spontaneously into any prognostic statement. Transient cerebral attacks may be premonitory and may call for a warning to the friends but even these form no certain basis upon which to predict. In those who do not display premonitory symptoms cerebral haemorrhage cannot be foreseen. For these we can think truthfully that it is unlikely that if it is going to occur it will probably occur years ahead. A cerebral haemorrhage though often fatal may be survived for many years.

A general and important consideration in the prognosis is the manner of living. Those with high tension who can and will live quietly live longer than those whom necessity or desires compel to live strenuously.

TREATMENT

Introductory — An important present day aspect of the care of high tension is to ensure that the patient's attitude to his condition is reasonable. The public has been allowed to acquire incomplete and dangerous knowledge of high blood pressure believing it greatly to shorten life and to threaten apoplexy. Any indication of anxiety about the height of pressure should at once be allayed by reassuring statements. Patients in whom high pressure is found should rarely be informed of the fact it is a technical detail that should not be allowed to concern them. The habit of following blood pressure by frequent readings during treatment and of allowing patients access to such readings is strongly to be deprecated. The patient becomes obsessed by blood pressure and the manometer is regarded as the gauge of health and readings are watched closely and anxiously. Such use of the manometer is also a mark of ignorance there is no virtue in the frequency of readings and blood pressure is not in fact a chief and not often a very important guide to the progress of the malady. Readings may be taken at intervals of weeks or months and in a manner to ensure that the measurement is regarded as but incidental. It is often necessary that a patient should be made to understand that his activities must be limited, this can be done almost

always by a simple statement concerning advancing age and general health, with at the most a reference to the circulatory system.

It is better for patients in whom high blood pressure has been discovered, but who are without complaint or symptoms, to be reassured by a statement that the condition is not dangerous and that they should continue their usual life.

DOCTORS' OPINION

Mental Quiet —Anxieties of mind are well known to increase blood pressure and to maintain it when high. Many patients with high pressure are harassed people, full of business worries, family difficulties and cares. The relief of worry, anxiety, and mental distress is a very material part of effective treatment. In the presence of troubles and to feelings of ill health, the dread of illness or disastrous accident must not be added to the existing troubles. These patients must be reassured and their minds must be at ease.

Work and Exercise —Hours of work that are excessive should be shortened. Strenuous forms of exercise are being forbidden. Patients having few symptoms and no pain can continue in sedentary or lightly active work. Indoor workers are advised to take exercise in the open air. The amount of exercise should be governed by the reaction of the body. It should be increased gradually.

Exercise that produces distress, such as headache or fatigue that is unsuitable, should be discontinued. The effect of exercise upon subsequent blood pressure should be noted especially.

Sleep —The pressure falls during sleep, and the patient should be reassured that this is normal.

DIET AND NUTRITION

Diet —Protein free or protein poor diet may be advised in severe cases.

unpalatable. There is no objection to moderate amounts of alcohol in the form of spirits and light wines. Beer or excessive quantities of fluid of any kind should not be drunk.

Obesity should be treated by withdrawing first of all any excess of carbohydrates or fats from the ordinary diet and later by a general reduction in the amount of food (to about 1500 calories value). Rigid dieting is not often well tolerated; it causes undue weakness, sudden reductions of weight should not be attempted. Regular exercise helps to reduce weight; so does massage. Turkish baths are not advised. Thyroid may be given to stout women who develop high pressure at the menopause.

Lowering Blood Pressure — The fear of cerebral hæmorrhage and the frequency with which heart failure ultimately supervenes have suggested the use of remedies calculated to lower blood pressure. Known substances and proprietary preparations have been and continue to be introduced in large numbers. Nitrites, nitrates, acetyl choline, theobromine and other xanthine derivatives, benzyl benzoate, thyroid extract, liver extract are but a few of the more widely tried. For each of these strong claims have been put forward, but none has come to be regarded as reliable or even as offering much or any prospect of benefit. Perhaps the most striking reflection upon their value is the short period of popularity that each enjoys before it is displaced by a rival. Even the long used remedy, potassium iodide, beneficial in some high pressure syphilitic cases, has fallen much in favour. The amount by which pressure is lowered by these remedies is not usually great, and the fall is either transient or restricted to the period of the drug's administration. Many patients complain that they feel less fit under such treatment than without it, and it is doubtful if the policy of lowering blood pressure is sound. No doubt vasodilator substances will continue to be introduced and used. There is no harm in trying them on occasion, but their administration should not be allowed to disorganise a patient's life, or to form the incentive to repeated blood pressure readings, or to divert attention from more natural and more effective remedies. There is no known method of lowering pressure materially and over long periods of time; the most effective treatment is a suitably restful regime.

A Regime — The most successful method of treating hypertension cases is by a careful regime, reducing physical and mental strains by imposing actual rest periods, short and frequent, or prolonged by increasing hours of sleep, by shielding the patients from worry of any kind, by keeping the bowels lax with salines, and the diet at

the minimal quantity required to prevent loss of weight by forbidding tobacco altogether or in anything but most moderate quantities. Such a regime, associated with intervals of light open air

for an hour or so — 25 — 41

W

and often effective

Angina Pectoris — In angina of high pressure the usual remedies are employed

Cardiac Failure — The treatment of cardiac failure has been dealt with earlier. There are a few supplementary notes to add. It is important that treatment by rest in bed should be used early in the cardiac failure of hypertension, namely when breathlessness occurs during brisk walking exercise and before signs of congestion appear. Several months of complete rest will usually relieve for a much longer period; it should be followed by long rest periods each day and especially after meals for the rest of life. In treating the terminal phases of cardiac failure nocturnal dyspnoea is often prominent and calls for morphia. Oedema is not often massive but if it should be the very strictest precautions should be taken when draining the legs; sepsis is much more likely to appear than in the younger cases of rheumatic heart disease. The usual diuretics may be used.

CHAPTER XXV

ARTERIOSCLEROSIS SENILE HEART MYOCARDIUM

ARTERIOSCLEROSIS

AFTER the age of thirty the arteries of the body are rarely quite normal and the pathological change increases steadily with advancing years. The condition varies greatly in severity, arteriosclerosis that comes early and progresses quickly runs in families. The disease is well known to be associated with high blood pressure but often occurs apart from this it is stated to be more prevalent in those consuming much alcohol and in those engaged in heavy manual work. Its cause, however, is unknown.

PATHOLOGICAL ANATOMY

The earliest signs of arteriosclerosis are to be found usually in the aorta where small yellow flecks are visible in the intima. In more advanced cases the aorta is studded with small and larger nodular masses of the same fatty or gelatinous substance. The deposits occur early around the mouths of outgoing vessels. Ultimately the whole normal intima is concealed and rendered irregular by the deposits in which rough calcareous plaques are laid down, and by denudation of the surface. The visible changes in the intima are generally regarded as compensatory to atrophic changes in the medial coat, the adventitial coat becomes thickened and sclerotic. In very advanced and rare cases the aorta is converted into a rigid and brittle tube. Similar changes can occur in all the arteries of the body, they are usually affected in the order of their size. When affecting smaller arteries the disease may fall with especial heaviness upon a given group of vessels like those of brain, kidney, heart or limb. Damage to the tissues in the corresponding territory occurs when the flow of blood through the affected vessel declines. This happens in one of several ways. Thrombosis in a patent vessel may yield acute and massive necrosis, which may or may not become converted ult-

mately into a local scar. When the wall of a smaller vessel becomes thickened, the lumen often narrows materially so that little blood or, when the tone of its wall increases, no blood can pass, and with inadequate collateral channels, the nutrition of the tissues in its territory will suffer. Thrombosis completing a closure already far advanced may happen quietly. Consecutive changes may be brought about in the muscle of the heart in one or other of these ways, and consists of fatty degeneration or replacement fibrosis.

RECOGNITION OF ARTERIAL DISEASE

Advanced arteriosclerosis of the aortic wall leads to loss of elasticity and to reduced resistance of its walls: the aorta elongates, becomes a little sinuous, and dilates (Figs 39 and 40). These changes



FIG 39—Orthodiagram. Man aged 50. Advanced general arteriosclerosis, angina pectoris, gout. Systolic blood pressure 165 mm. No valve disease. Wassermann reaction negative. The aorta is very tortuous; its probable course is marked by dotted lines.




FIG 40—Orthodiagram. Man aged 63. weight 108 lb. Advanced general arteriosclerosis, systolic blood pressure 210 mm. No valve disease. The ascending aorta is elongated and unusually curved.

can occur independently of, but are exaggerated by, high blood pressure. Occasionally, with nicely adjusted exposures, calcareous deposits in the aorta are displayed radiographically, as they can be more frequently in the vessels of the limbs.

At the bedside, arterial disease is recognised by noting prominence of the vessels in their courses and by their tortuosity. In the normal arm of the young adult no pulse is visible, but in arterial disease pulsation appears over the course of the vessels, at first where the vessels normally lie most superficially, as at the wrist and before the elbow joint, ultimately throughout the whole length of the upper arm, and even on occasion throughout the forearm. The pulsa-

tion is naturally more conspicuous in cases in which the pulse is large, and it is also more conspicuous when the arm is flexed, for the slack in a vessel that has lost its elasticity is not taken up when its course is shortened, and so the artery is forced to become sinuous. When the disease is more advanced, the vessel is actually elongated and its course is then tortuous when the arm is extended. Sometimes tortuosity is so great that the brachial artery is bent until its course is reversed for a short distance, and in very rare cases may actually make a complete loop in the arm. Tortuosity of the superficial temporal vessel is of less value because these vessels naturally follow a wavy course and, when dilated, their very superficial position renders them unusually prominent.

At the wrist, thickening of the radial artery may often be felt. A length of vessel should be emptied of blood by occluding it above, wiping out the blood and occluding it also below. If the stretch of empty vessel between can then be felt and rolled under the finger, or if its surface is irregular, it is diseased. A normal artery when full and tense can often be rolled under the finger in this way.

Strictly speaking, disease is only recognised in the artery in which it is observed, but because arteriosclerosis is almost always widespread, conspicuous changes throughout the vessels of the arm may be used as sufficiently convincing evidence of widespread disease. It cannot be used to diagnose appreciable disease of particular vessels, coronary or cerebral though it does indicate that the case is of a type in which such disease may be expected to exist, and it helps to interpret corresponding groups of symptoms in cases of doubt. Irregularity of the retinal vessels provides a further hint.

CONSEQUENCES OF ARTERIAL DISEASE

Although essential hypertension and arteriosclerosis are separate pathological conditions, they are very often associated. That is so because they are both common maladies of the same age period and because, when there is a tendency towards arterial change, its development seems to be aided by high pressure, it is a mistake to think that diseased arteries cause high pressure.

Essential hypertension is so frequent and so important clinically, while at the same time sufficiently defined, that it is important to give it the emphasis of separate portrayal. By so doing, many of the phenomena associated with, and originally regarded as, simple manifestations of arterial disease, become included under essential

hypertension Some of these, for example cardiac hypertrophy, are properly regarded as pertaining to high pressure and not to arteriosclerosis, which, unless extreme, throws little burden upon the heart Others are to be regarded as pertaining to both, thus angina pectoris in part results from an increased burden, and in part from the disease of coronary vessel and cerebral haemorrhage, while occurring almost exclusively in high pressure cases, would not happen unless the unusually strained vessels were diseased Lastly, there are phenomena that pertain to arterial disease itself, for example thrombotic accidents Very many of the manifestations described as accompanying essential hypertension might, with equal reason, have found their place in this chapter To divide strictly and accurately is impossible and the attempt would have separated phenomena that are usually found together The present brief account is intended in part to correct any false impressions that might otherwise arise, and in part to supplement

Apart from rupture of the diseased vessel wall, an accident sufficiently noticed in the last chapter, the material consequences of arterial disease are nutritional The special symptomatology of arteriosclerosis is to be found in the abrupt failure or decline of blood supply to various organs, such as the brain the kidneys, the pancreas, the limbs and the heart The more important cerebral manifestations have been named in the last chapter The deeply scarred arteriosclerotic, contrasting with the granular arteriolosclerotic, kidney has no great clinical significance In the arteries of the pancreas disease gives rise to diabetes, in the vessels of the limb to distal gangrene or when arteries to the muscles of the calf are involved to intermittent claudication Acute closure of the coronary vessels leads to the syndrome fully considered in Chapter VI gradual closure is responsible for angina of effort, and by decreasing the musculature and robbing it of strength, leads up ultimately to failure with congestion, clinical states already considered at length

A frequent association of arteriosclerosis and the degenerative processes of advancing years is auricular fibrillation Its development may determine failure with congestion, but more often the change passes unnoticed, the rate of ventricular response being slow in the elderly Alternation of the heart beat is often seen

HEART DISEASE IN THE AGED

PATHOLOGICAL ANATOMY

Coronary arterial disease is the chief feature. The senile heart may be enlarged apart from high blood pressure not infrequently it is normal in size or may even be a little reduced. There is often a diffuse increase of fibrous tissue in it apart from such larger and denser areas that have arisen obviously from vascular thromboses. The muscle is often pale and special stains reveal diffuse fatty degeneration. It may be brown in colour and show pigmentary degeneration microscopically.

An ascending degeneration of the aortic cusps of which the mild forms are common in adults is distinct pathologically from arterio sclerosis which usually accompanies it. It begins in the substance of the cusp near its base and passes on to thickening that spreads up the cusp. Then large calcareous deposits occur at the bases of the cusps and project into the sinuses of Valsalva. The craggy masses cause obstruction. This degeneration is often responsible for aortic disease especially stenosis in elderly people. Similar degenerative change sometimes happens in the aortic leaf of the mitral valve and calcareous deposits are laid down at its base and in the adjoining part of the ventricular septum in the last situation occasionally breaking through the auriculo ventricular bundle.

CLINICAL FEATURES

In elderly or in old people the heart presents many different clinical pictures. There are instances of long standing rheumatic disease of the aortic and mitral valves. There are long surviving or late developing cases of syphilitic aortitis. But the degenerative lesions named earlier dominate in the cardiac disease of old age and produce the acute phenomena of coronary thrombosis or spasmodic angina and failure with congestion often supervenes upon one or other of these or occurs without these preliminaries as the amount of sound muscle in the heart dwindles. Here too must be mentioned again craggy aortic stenosis and the occasional case of heart block which may result from similar calcareous lesion or from fibrosis in the ventricular septum. The development of angina will be expedited as will failure with congestion in other cases by the increasing arterial pressures which are so frequent with advancing years. These various forms of disease occurring singly or in combination have been described already.

There remains for brief notice the heart that fails in old age without displaying any of the major antecedent troubles and in a less conspicuous way. As age advances a general lowering of vitality occurs and nutrition begins to fail generally and the body wastes. A systolic apical murmur appearing for the first time may have value as an early guide to a dilating heart. The bodily resources have usually begun to decline long since and amongst these is cardiac reserve, displaying itself by breathlessness on effort. In the decline there is often seemingly a slow race between cardiac capacity and the powers of the body generally. The heart is losing its reserve, but the man is becoming progressively enfeebled, owing to local arterial trouble or general weakness his limbs serve him less well, his will or inclination to carry out his old pursuits declines, he requires and takes less food and thus, as the heart weakens so simultaneously the body is brought more and more to rest and the burden falling on the heart lightens. The enforced decline of bodily activity constitutes a spontaneous remedy, which tends to prolong life, sometimes lengthening it remarkably though always precariously. The balance is unstable since the reserves are gone, and is quickly broken by any unusual event, such as an enforced period of activity, or an infectious disease such as bronchitis or pneumonia. In old men failure with congestion, when it comes, terminates life quickly and often with relatively few signs. The aged pass out unobtrusively, after brief illness, or without warning, while sitting in their chairs or sleeping in their beds.

THE MYOCARDIUM

In this and previous chapters much has been said of cardiac failure and of anginal pain, and very little directly about the muscle of the heart. This is as convenient a place as any to consider the relation of diagnosis to the myocardium. There are some physical signs as yet undescribed, which are relevant from this standpoint. These will be discussed briefly with comments upon their values.

SOME PHYSICAL SIGNS OF MYOCARDIAL INVOLVEMENT

Pulse Strength—The size of the pulse rarely has much value in gauging the state of the cardiac muscle, it is largely controlled by the local state of the artery and by the rate at which the heart is beating. Thus, the pulse is normally much smaller when the hands are cold than when they are warm, and it is usually very small in

paroxysmal tachycardia. Decreased pulse tension (low blood pressure) is not much more significant. It is true it may result from deficient output of the heart at each beat consequent upon weakness of the cardiac muscle but it may result also from deficient return of blood to the heart as in haemorrhage and as in profound loss of vasomotor tone. Central and peripheral factors are often combined and notably in the acute infectious fevers. The pulse weakens to imperceptibility in the last stages of many diseases, including cardiac failure but by that time the state of the patient is usually apparent at a glance. It is to be remarked that patients frequently present advanced signs of cardiac failure and high blood pressure simultaneously.

Heart Sound Intensity —The intensity of the heart sounds and especially of the 1st sound has long been thought to throw light upon the strength of the heart. It is true that decrease in the intensity of the 1st sound often occurs when the circulation is failing rapidly as in the acute infections a failure that is in part cardiac and in part vasomotor. It is also true that it occurs in terminal cardiac failure but at a time when there is a redundancy of signs. Ordinarily the intensity of the sound varies greatly according to the amount of covering of the heart and chest wall according to the way in which the heart is acting and with less easily determined factors. Emphysema and obesity both greatly diminish the intensity of the sounds. Simple but very rapid action as in paroxysmal tachycardia weakens the 1st sound relative to the 2nd and produces sounds having much the spacing and intensity of the foetal heart sounds. It is not questioned that weakening of the heart beat results in lowered intensity of the sounds it is merely pointed out that other factors are involved. Actual experience shows that in gauging the essential soundness of the heart in patients who are up and about the intensity of the sounds helps very little. Similarly, signs based upon change in the character of individual sounds though of some theoretical interest are insufficiently precise, and lack enough constancy in given circumstances to be of any great practical value.

Gallop Rhythm —This is constituted by three distinct heart sounds, usually of almost equal loudness. Usually the heart rate in these cases is a little rapid (90 or 100) the gallop is best heard at the impulse, though often over a wide area. The extra sound is found in graphic records (Fig. 41) to lie in presystole or in mid diastole, and to be related to the contraction of the auricle. This gallop rhythm is found especially in hypertension cases in which the

heart is failing, or in the heart failure of elderly subjects. There is as a rule no valve disease. The sign is often associated with prolongation of the auriculo-ventricular conduction interval, or with bundle branch block. The cardiac impulse is often double. Gallop rhythm may help to determine the cardiac origin of breathlessness or pain, by demonstrating the heart to be abnormal, for only patients with abnormal hearts present this sign, and they are usually serious cases. There is no constant underlying lesion, though hypertrophy of the auricle is common.

Wide reduplication of the 2nd sound, which also gives a sort of gallop rhythm, occurs especially in cases of mitral stenosis with slower heart action. The extra sound, in the early part of a long diastole, is unrelated to auricular systole, as it may occur when the auricles are fibrillating, it is probably due to an unusual intensification of the normal 3rd heart sound, and tends to be confined to the region of the impulse.

"T" wave inverted — In young and healthy adults *T* is always upright in an electrocardiogram taken from the right arm and left leg. An inverted *T* (Fig 23, page 113) is abnormal but occurs in a number of conditions. Temporarily it is seen under heavy dosage with digitals or after drinking quantities of ice cold water. As a persistent condition it is common in elderly heart patients, and most frequent in those in whom there are abundant other signs of cardiac weakness. It frequently appears a week or more after coronary thrombosis (Fig 6, page 46), and often persists. Although the death rate amongst those displaying it constantly is heavy, it has been known to persist for twenty years in a patient whose health remained unchanged during that period. It should not be used for prognostic purposes.

Bundle Branch Block — Large abnormal electrocardiograms in




FIG 41 — Simultaneous electrocardiogram (lead II) and sound record from the apical region. The electrocardiogram exhibits block in a branch of the auriculo-ventricular bundle. The sound record shows gallop rhythm the 1st sound of the heart being double and its first element falling in presystole. The black rectangle indicates the limits of ventricular systole. Time in $\frac{1}{4}$ sec.

which the initial phases are very wide and the chief and final deflections are in opposite directions (Fig 42), are caused by interference with one or other branch of the auriculo-ventricular bundle. The pathology of the condition and its clinical associations are similar to those already described under heart block, with which it is often combined though bundle branch block is most frequent in cases of aortic disease and coronary disease. As a temporary condition it indicates active mischief in the heart, and may be the only clear indication of this. As a persistent condition it is again significant of a myocardial affection. As in the case of heart block, it is usually to be regarded as indicating changes that are not confined to the special tract, but are more widely diffused. It is almost always accompanied by abundant cardiac symptoms and signs pointing to serious cardiac involvement such as those of cardiac failure, anginal pain, Cheyne Stokes breathing gallop rhythm, pulsus alternans, and attacks of nocturnal breathlessness and most of the patients who exhibit it are dead within two years. But some, and especially those in which there are not the ominous associations, survive for many years, and so the prognostic significance of the sign itself is indecise.




FIG 49.—I electrocardiogram from leads I, II, and III illustrating the common type of bundle branch block. The main deflections in leads I and III are in opposite directions. The initial and final deflections in each of these leads is opposite in direction. The initial deflections in each lead are conspicuous in amplitude and unduly prolonged.


Its chief value is that in cases of obscure breathlessness or pain it may indicate the heart as the site of mischief.

Pulsus Alternans—The pulse beats are rhythmic but alternate in height (Fig 43). Its cause is thought to be failure of a proportion of the fibres of the ventricle to

contract at alternate beats of the heart. Alternation often occurs when the heart is beating very rapidly, it is usual in auricular flutter, and frequent in paroxysmal tachycardia. It occurs also when the heart is beating slowly, and is then most frequently associated with hypertension, cardiac failure, angina, coronary thrombosis, or syphilitic aortitis. Its heaviest incidence is between the years fifty and seventy. Whenever it occurs there is reason to believe that the heart, or a substantial part of it, is struggling

to perform work of which it is scarcely capable. If the heart is disposed to alternate added strain such as high blood pressure or rapid beating will determine it. An extrasystole will often disclose alternation for a few cycles (Fig. 44).

Although of common occurrence it cannot be recognised by feeling the pulse or cardiac impulse but must usually be shown



sphygmographically. Not infrequently when persistent it can be detected by means of the blood pressure apparatus alternate beats forcing their way through the cuff at pressures 5 or 10 mm Hg higher than the remainder.

Occurring at high rates of beating it has relatively little significance. If persistent at slow rates of beating it is a grave sign. Few patients displaying it survive for one and very few for two




FIG. 44.—A radial pulse curve showing alternation in its second half. The alternation has been provoked by the extrasystole (Ex) which stands in the middle of the curve. Time 1 sec.

years. It is usually accompanied by abundant other evidence of grave heart affection: signs of failure, angina, attacks of nocturnal dyspnoea, Cheyne Stokes breathing, gallop rhythm, or bundle branch block. Though never the sole sign of disease, persistent alternation is occasionally found in the absence of the named phenomena and then still retains its ominous significance. There is no lesion of the muscle with which this curious and important disorder of beating is known to be correlated.

MYOCARDITIS FATTY AND FIBROID HEART

Three morbid conditions of the muscle alone call for brief consideration here. They are "myocarditis", "fatty heart", and "fibroid heart."

Myocarditis—The only form of inflammation of the heart muscle that is at all common is rheumatic in origin, and it takes an acute or more often subacute form. Occasionally inflammation occurs in other acute and fatal infections, diffusely in diphtheria locally in pyæmia. The term myocarditis has often been allowed most loosely to comprise all altered states of the muscle, real or supposed. It has been employed improperly as an equivalent of myocardial weakness. Actually the use of the term myocarditis is scarcely to be justified clinically except in relation to rheumatism, and here the term "carditis" is sounder. The description of myocarditis, acute or chronic, as a separate disease with its special symptoms and signs needlessly complicates a general description of cardiac disease which it is essential should remain simple.

Fatty Heart should not be diagnosed. If a patient has cardiac pain and is obese, it is easy to guess that around the heart an unusual amount of fat has accumulated. The guess, however, would be equally correct if there were no pain. A diagnosis of fatty heart in such circumstances has no more value than has that of "obesity", and thus it is objectionable because it pretends to be more than it is. If we mean by fatty heart a fatty degeneration of the muscle, then this cannot be diagnosed with any pretence to accuracy, unless perhaps in acute poisoning by infective disease or phosphorus, or in anaemia, and in these the desire to make the diagnosis is infrequent. In severe diphtheria the cardiac muscle may be so degenerate with fatty particles that it is throughout of a pure yellow colour, yet the heart continues beating. Because diffuse fatty degeneration occurs in the senile heart, or in coronary arterial disease, that is no reason for regarding fat in the heart as a cause of sudden death. Equal change is found in the heart of pernicious anaemia and here sudden death is most unusual. If a man dies suddenly in the street, a court is required to find the reason, and if nothing else is forthcoming, "fatty heart" by usage proves convenient and acceptable. The enquiry may not rest upon the answer "Death from natural causes" unless an anatomical basis for death is found. There is evidence that the ventricles of elderly people, and especially those presenting coronary disease, are prone to fibrillate. This fatal fibrillation is an

event in living muscle it leaves no message written upon the dead heart. The conception that correlates fatty heart and sudden death may be useful and therefore still justifiable in forensic practice but while fatty heart is usually unrecognisable in the living and its prognostic significance is uncertain the conception can find no place in clinical diagnosis.

Fibroid Heart—Fibrosis in the heart is found in one of two chief circumstances. Firstly it occurs very diffusely as an end product of rheumatic inflammation it is then mainly an interstitial though in part a replacement fibrosis. Secondly it occurs in the senile heart either as an after effect of vascular thrombosis and representing areas of healed infarction or more diffusely and more commonly as a consequence of diminished nutrition in narrowing of the coronary arteries and arterioles these two forms of fibrosis are often combined. That fibrous tissue is present in excess in the ventricle may be inferred from a history of rheumatic carditis or of coronary thrombosis but its extent cannot be ascertained. The patient's outlook depends not on the fibrosis but upon the state of the remaining muscle. For these reasons fibrosis of the heart should not be diagnosed the patient's condition being summed up in terms of its functional ability.

MYOCARDIAL WEAKNESS

It will be gathered from what has gone before that an attempt to correlate physical signs with specific states of the cardiac muscle offers infrequent prospects of success. Inflammation of the myocardium can be diagnosed safely in subacute rheumatism when the heart is clearly involved but the intensity of the morbid change the numbers of submiliary nodules present cannot be gauged. The weakness of the muscle correlates with more subtle processes. Fatty degeneration of the muscle in chronic heart disease is of unproved significance and cannot be diagnosed accurately. Fibrosis of the heart can only be diagnosed satisfactorily when the fibrous tissue has followed locally a coronary thrombosis. Myocarditis fatty and fibroid degeneration are the only gross changes that are at all common in the heart. Heart block or bundle branch block may indicate focal lesions in the conducting tissue and may allow us generally to infer that the changes are actually more diffuse in the muscle but the type of lesion cannot often be stated and its extent is quite problematic. Fibrillation of the auricles is usually but not constantly accompanied by morbid changes in the auricles. Thus

the basis on which to diagnose myocardial lesions remains a very slender one. Moreover, most patients dying of cardiac failure show little change of any of these kinds.

An outstanding fact of cardiac pathology is that we are unable to demonstrate, after death, the changes in cardiac muscle that have led to its failure. Minor changes are found in failure, but they are inconstant in kind, occurrence, and degree. We are rarely able to predict before death the specific changes that will ultimately be found in the ventricle after death, and never the extent of the change. It is sounder in practice not to make the attempt to diagnose in terms of anatomy, but in the more certain terms of function. Angina pectoris and cardiac failure are clinical states, which may be regarded as the chief products of defective nutrition or of weakness of the myocardium. It is in these terms that diagnosis must usually be made, because they convey with sufficient precision states that are recognisable and measurably significant. When we merely substitute for the functional diagnosis such terms as coronary disease, myocardial disease, or whatever is supposed to be the underlying basis of the disturbance, we introduce assumption and, while gaining nothing substantial, sometimes depart gravely from truth. Angina pectoris and cardiac failure never can find equivalents in the terms of structure, for each depends not only upon something in the heart but upon something that the heart is called upon to struggle against that lies outside itself.

The desired end point, the diagnosis of heart failure, in one or other of its degrees, or of angina pectoris, is often aided by the proper use of certain of the physical signs described earlier in this chapter. A patient has breathlessness or pain, but in neither instance is the origin of the symptom manifest, the diagnosis will often be brought to the point of justification, or actually be rendered certain, by the finding of a persistent phenomenon such as auricular fibrillation, heart block, bundle branch block, gallop rhythm, or, a little less emphatically, by inversion of *T*. For each of these phenomena points to definite abnormality of an organ from which the symptom under consideration often is derived. Alternation of the pulse is of decisive value, for it not only shows the heart to be abnormal, but actually presents clear evidence that the heart, or a substantial part of it, is labouring under difficulty.

CHAPTER XXVI

BRONCHITIS AND EMPHYSEMA

INTRODUCTORY

In this chapter the effects of chronic bronchitis and emphysema upon the heart will be discussed briefly. The reduction in the vascular bed of the lungs in emphysema, following destruction of alveolar walls and changes in the small vessels, increases the resistance to the output of blood by the right ventricle. The reduction of alveolar walls, the thickening of those that remain, and deficient power of expiration, reduce the capacity of the lungs to aerate the blood passing through them. Bronchitis, according to its severity, aggravates this defect and the blood passes to the systemic vessels in a state of imperfect oxygenation. Bronchitis also imposes upon the heart the strain of persistent coughing, and lastly, and perhaps most important of all, it introduces the factor of infection.

The clinical picture of bronchitis and emphysema is sufficiently familiar, it is enough here to note main features. The patient is usually a middle aged man, giving a history of recurrent cough of many years duration, the chest is fixed in the extreme inspiratory position, it is short, deep, and presents a wide intercostal angle. The chest wall is rigid and its movements small, expiration being particularly difficult. Evidence of expansion of the lung is found in the universally hyperresonant note, and in the extension of lung resonance over the heart and beyond its usual limits over the lower ribs. Breathlessness varies a good deal in its amount according to the accompanying bronchitis, which may be chronic, but which more usually recurs in acute or subacute attacks, or becomes exaggerated in the winter time.

PATHOLOGICAL ANATOMY OF HEART

The heart is generally enlarged, though usually not remarkably, occasionally very large hearts are found, sometimes they weigh no

more than normal. The enlargement affects both ventricles, but it is the rule for the right to be hypertrophied to an obviously greater extent than the left. This disproportion does not always exist, and is not in general so conspicuous as in mitral stenosis and decidedly less so than in pulmonary stenosis. General systemic arterial disease is commonly present, the coronary arteries are not infrequently involved and may show old standing thrombosis. The pulmonary arteries usually present atheromatous changes.

THE CAUSE OF THE CARDIAC FAILURE

Venous congestion in this malady is usually ascribed to failure of the right ventricle owing to the extra work it has to do. Doubtless this increased burden contributes to the end result, but it is not the sole factor. In cases of congenital pulmonary stenosis the burden during life if we may judge from the weight of the right ventricle after death is much greater than in emphysema, yet failure in pulmonary stenosis does not usually occur, and, occurring, is delayed for very many years. Failure in emphysema is not a simple affair of pulmonary arterial pressure. In not a few of the cases there is systemic hypertension, in others there is coronary disease, in still others auricular fibrillation points to an affection of the muscle and itself burdens the heart. The heart is subject to the strain of powerful coughing which raises blood pressure. The condition of the arterial blood is such that the heart muscle, like other tissues, is more or less asphyxiated. Lastly, there is the factor of infection. Emphysema is incurable, it can become worse but not better, yet congestion when it comes in these cases, though it may be progressive, is by no means always so frequently it comes with the attack of acute and subacute bronchitis and clears away again. Failure with congestion is not determined by emphysema in such cases but by bronchitis, it is determined, therefore, not by increased work of the right ventricle but by factors associated with the acute process, acting upon a heart already burdened.

CARDIAC SYMPTOMS AND SIGNS

It will suffice to note the special points that concern the heart, and its failure in this malady.

Recession of the Heart—The expansion of the lungs causes the heart to recede from the chest wall. The impulse is obscured, its superficial dulness disappears and the heart sounds are rendered

distant It is therefore impossible to determine the heart's size with accuracy, except by X ray examination

Cardiac Failure —When the heart fails, the signs and symptoms are similar to those described earlier in this book There are a few notable peculiarities Breathlessness, already present, becomes unusually severe The rigidity and urgency of breathing is notably greater in relation to the amount of venous congestion than in ordinary failure with congestion Cyanosis, which is of central type, is of unusual depth Thus, with the veins a little swollen in the neck, and the liver a little enlarged, cyanosis and breathlessness may be conspicuous, when the paroxysm of coughing comes, the veins in the neck swell greatly and the patient's skin and lips become deeply suffused and intensely cyanotic

PROGNOSIS AND TREATMENT DURING FAILURE

The immediate prognosis depends upon the course of the bronchitis If this has been chronic, the outlook is unpropitious If an acute or subacute attack is relieved the signs of cardiac failure clear away also actually recovery usually occurs in these attacks though it is gradual and may be spread over several or many weeks The patient may or may not be left measurably the worse after the attack is over Extensive infarction of the lungs sometimes kills these patients

When patients with emphysema begin to show definite signs of cardiac failure whether in attacks or in exacerbations of bronchitis or otherwise they are entering upon the last stages of their malady Life is not usually prolonged for more than one or two, exceptionally for a few years This prognosis conforms to what is usually found in failure with congestion when this develops in middle aged or elderly men

The treatment of the cardiac failure is carried out on lines similar to those already laid down Rest and efficient nursing are essential Auricular fibrillation when present is treated appropriately Vene section is useful when the veins are much engorged, depth of cyanosis is not the guide Efforts should be directed to relieve the strain of coughing Ammonium carbonate or chloride (5 grain doses, four times daily) are most valuable remedies when the sputum is scanty and viscid potassium iodide may be used in similar doses for the same purpose Ipecacuanha may be employed if it causes no nausea Relief may be secured by saturating the air which the patient breathes with moisture by the use of the steam kettle The patient

should endeavour to avoid strain in coughing, and may be helped to do so by the use of codeine ($\frac{1}{2}$ grain) or heroin ($\frac{1}{16}$ grain), but sedatives should not be given to the extent of making expectoration inefficient

Patients of this class live longer in warm dry climates, they should avoid dusty atmospheres, public buildings, and crowded rooms

CHAPTER XXVII

THYROTOXIC STATE

INTRODUCTORY

CHIEF manifestations of the thyrotoxic state, and the most diagnostic clinical signs of the disease, are to be found in the thyroid gland itself and in the eyes. The condition occurs chiefly in women between the ages of twenty five and forty five. It is enough here to name the firm enlargement, either uniform or nodular, and the vascularity of the gland, together with the protruding, staring eye balls. Associated with these most characteristic phenomena are others. There is change in the nervous system represented by an alertness, restlessness, quickness of movement, emotionalism, and tremor of the hands. Sweating — in excess, there is sometimes a diffuse pigmentation of the skin. Metabolism is increased, and with this may be linked wasting, weakness, and occasional glycosuria.

Lastly, in all but exceptional cases, there are disturbances of circulation that here claim particular attention, and not only because they may be crucial in leading to diagnosis. The circulatory system is found to be in a state of increased activity, which is often remarkable and always important. The development of circulatory failure sometimes jeopardises life, and it leads too often to the indefinite postponement of surgical treatment.

CARDIOVASCULAR MANIFESTATIONS

Pathological Anatomy — In patients dying of cardiac failure the heart is found to be dilated and not infrequently hypertrophied, though not greatly. No distinctive histological change has been found in the muscle, which is usually little altered.

Vasodilatation — Cutaneous vasodilatation is the rule, and evidence of it is often conspicuous. The two most valuable signs are increased temperature of the skin and capillary pulsation. It is increased temperature of exposed parts such as face and hands that

must be observed, the face feels hot and the hands keep warm or hot when outside the bed clothes in cool or cold rooms. The patient feels hot and prefers cool room temperatures to warm. The face is often flushed sometimes deeply, and this flush extends to the neck and covers the body, though less conspicuously. It is a mistake to regard this depth of colour as important, freshness of tint, and associated warmth and capillary pulsation, are of much more consequence. Not infrequently the depth of colour is natural or there may even be pallor. Capillary pulsation is always to be seen in the face when there is appreciable vasodilatation, though experience is required to detect it. As a spontaneous event it is plainest in the cheeks and forehead and should be looked for with keen attention from a little distance. On no account should the skin be rubbed, but a glass slide may be pressed on cheek, lobe of ear, and lip if the pulsation cannot otherwise be detected. It should also be sought in the pads of the fingers if, as is usual, these are warm.

Pulse —When regular the pulse is rapid, the rate being often as high as 100 to 120 while the subject rests, and rising higher with exercise or excitement. It is unusually high in sleep. The stroke is full and abrupt. Systolic blood pressure is in general raised, ranging around 140 or 150 mm Hg, diastolic readings show no rise.

Precordial Signs —The patients are usually thin and the heart is overacting. Therefore a diffuse impulse is customary, it almost slaps the examining hand and not infrequently imparts the feeling of a slight thrill. The maximal impulse is often displaced outward an inch or a little more in the 5th space. The character of the impulse is apt to convey the impression of a larger heart than is there. Abrupt and accentuated heart sounds are frequent, a systolic murmur is common over 2nd and 3rd left interspace and, when the impulse is displaced outward, over this also. A cardiorespiratory murmur in the left axilla is usual, when the heart is overacting.

Auricular Fibrillation is present either in paroxysmal form or as a continuous affair, in one fourth to one third of the cases as these come for treatment. The ultimate reason for this frequent and important accompaniment is unknown, though it is recognised to be the direct or indirect, though impermanent, result of the thyroid activity. The incidence of fibrillation rises strikingly with the age of the subjects, and is much more frequent in those presenting a previous history of rheumatism. The rate of ventricular response to the fibrillating auricle tends in these cases to be high. When

fibrillation comes it increases the degree of cardiac failure or introduces it. Quite occasionally flutter or paroxysmal tachycardia takes its place.

Cardiac Failure — Breathlessness on effort the symptom of failure is the rule in these patients. This failure is to be ascribed to overaction of the heart in the presence of thyroid intoxication. The energy expenditure of the heart is increased in part because its rate of beating is high and in part because its output of blood is raised and this output takes place against a systemic pressure that is often above the normal. General venous congestion supervenes in some being chiefly confined to those presenting auricular fibrillation. In identifying general venous congestion stress should not be laid upon engorgement of the cervical veins in patients who present signs of increased vascularity of the thyroid gland. For when the gland has a thrill over it many unusually large subcutaneous veins can often be seen returning blood from its region in unusual quantity and this tends to overfill the cervical veins as a whole. Enlargement of the liver is in this instance a more valuable indication of failure than swelling of the cervical veins.

Anginal Pain at first with effort and later at rest occurs in some of the more elderly patients whose auricles may or may not be fibrillating. It may be severe and the attacks repeated by day and night. It probably occurs only in those predisposed to it already.

DIAGNOSIS

The diagnosis of the thyrotoxic state presents no difficulty when exophthalmos, a characteristic thyroid tumour and overacting heart are found together. It is the latent case in which ocular signs are absent and the thyroid is but little or not recognisably enlarged that presents difficulty and in such the suggestion that the thyroid gland is at fault comes first from the circulation. All cases of paroxysmal auricular fibrillation should be considered from the standpoint of thyroidism so should cases of persistent but recent fibrillation. The question may also be raised advantageously in cases of longer continued fibrillation that are unexpectedly resistant to treatment. If distinct symptoms of cutaneous vasodilatation as previously detailed are present in such patients the diagnosis is probable. If similar symptoms of vasodilatation are associated with a long continued and unexplained simple tachycardia the thyroid is to be suspected. The occurrence of severe thyroid intoxication in middle aged women with little thyroid enlargement is to be

emphasised In these types, hints may be obtained from nervousness of the subject from slight diffuse pigmentation, from occasional but unexplained diarrhoea, from loss of weight, and from signs revealed by close scrutiny of the thyroid An estimate of basal metabolic rate will usually decide A rate raised more than 25 per cent above the normal value is considered, in the absence of fever or other recognised cause, to be strongly suggestive, values raised 40 or 50 per cent or more are not infrequent A normal basal rate excludes hyperthyroidism a matter of occasional importance in differentiating cases presenting severe effort syndrome (page 158)

TREATMENT AND PROGNOSIS

Whether it is looked at from the standpoint of the heart or not, the treatment is essentially surgical namely, subtotal thyroidectomy For several weeks the patient is kept as quiet as possible in bed, while becoming accustomed to surroundings During this initial period the pulse rate usually falls a good deal without medicinal treatment Iodine (5 minims well diluted of a 5 per cent solution thrice daily) is given, it lowers metabolism temporarily, the pulse falling to 90 80, or even 70 per minute and symptoms subsiding A maximal effect is to be obtained and the operation then performed If the reaction is unsatisfactory during the first course of ten to fifteen days a second course on slightly heavier dosage may be instituted after two weeks interval if the maximal point is passed in the first course, the course may be repeated after a similar interval

This preparatory treatment to thyroidectomy greatly decreases the risk of post operative thyroid intoxication and from the same point of view iodine is often continued for a week after operation Under this regime, and in the hands of a skilful and experienced surgeon, the mortality from the operation is stated now to be less than 1 per cent the beneficial effects are usually conspicuous and often lasting In some cases removal of a second portion of the gland becomes advisable owing to recurrence of symptoms

Patients who have already had paroxysms of auricular fibrillation often develop an attack during the first two days following operation Such patients may be given digitalis before operation, otherwise they should be brought under the influence of the drug rapidly (page 88) as soon as fibrillation appears, if the patients are sick, the digitalis may be given by rectum

In patients presenting persistent auricular fibrillation, digitalis

should be used at once to lower excessive ventricular rate, and should be continued until signs of failure have been reduced as far as possible. There need be no hesitation in giving full dosage, provided that it does not induce vomiting. While continuing under digitalis, a ten days' course of iodine is begun and the case is treated surgically when this is complete.

It is becoming clear that few cases, whether they present auricular fibrillation or not, and with or without congestion of the venous system, are unsuitable for treatment along these lines. Although the risk from the operation is raised by the presence of failure, this risk is more than compensated by the fact that the chances of really substantial recovery, or of saving the patient from terminal cardiac failure, are very greatly increased by surgical intervention. The operation should be undertaken even when after medical treatment there are residual signs of failure. Improvement of the circulation in these cases with failure is usually remarkable after operation.

It is not unusual for the auricular fibrillation previously present to stop spontaneously within a few weeks of successful operation. If it does not do so within three months, quinidine can be employed to arrest it if, on grounds previously considered (page 90), this is thought desirable.

CHAPTER XXVIII

CONGENITAL MALFORMATION

AMONGST the great variety of congenital malformations of the heart that occur, the physical signs of only a few are distinct. The chief pictures of clinical importance are described briefly in the present chapter. A number of the grosser defects are incompatible with extrauterine life or are survived only for a short period after birth. Different forms of congenital disease are particularly difficult to distinguish in very young children.

PERSISTENT DUCTUS ARTERIOSUS

This duct unites the arch of the aorta beyond the origin of the three main vessels to the left branch of the pulmonary artery. In the foetus it carries blood from pulmonary artery to aorta. The duct closes at birth, when the pulmonary falls below the aortic pressure, and it soon becomes impermeable and atrophies. In cases of its persistence the blood flows through it continuously from aorta into pulmonary artery. The characteristic sign is therefore a murmur waxing in intensity as the aortic pressure rises in systole and continuing in diastole until the next systole, this murmur is usually a harsh and loud one and accompanied by a thrill. Its maximal intensity is at the 2nd or 3rd left cartilage, but being intense it is heard over a wide area at the base of the heart, and usually in the interscapular region, but not in the neck. Sometimes the murmur is cut short in diastole. When this murmur is heard, the diagnosis is almost certain. Continuous murmurs of an identical kind, however, are heard over anastomoses between large arteries and veins, these are usually in the limbs or neck, but occur occasionally within the chest as a result of bullet wounds. A continuous murmur varying in intensity with respiration and heart beat is heard over the jugular veins in many cases of severe anaemia. Although this noise is sometimes audible a little way down into the chest, it is unlikely to

be confused with a ductus murmur. If in patency of the ductus the leak is sufficiently free, arterial signs similar to those in aortic regurgitation may appear, the pulse rising more steeply, the arteries throbbing more, and the diastolic pressure being lower than usual. These signs are rarely more than slightly developed.

The defect causes no appreciable enlargement of the heart but is often associated with dilatation of the pulmonary artery, which may give dulness in the 2nd and 3rd left interspaces near the sternum.

It may occur alone or in combination with other malformations such as pulmonary stenosis. Occurring alone and in its usual form, its chief prognostic significance is that it predisposes the subject to infective endocarditis, otherwise it has little known effect on the duration of life.

BICUSPID AORTIC VALVE

This is a very common defect, being found about once in every 100 or 200 autopsies. It is not diagnosable, and its only importance lies in the frequency with which it is responsible for subacute bacterial endocarditis (page 183), about one fifth of the cases succumb to this malady.

COARCTATION OF THE AORTA (ADULT TYPE)

When the ductus arteriosus shrinks, the aorta sometimes becomes constricted at the insertion of the ductus, the obstruction is partial or more usually complete, the aorta appearing as though a cord were tied around it. The constriction occurs just beyond the origin of the subclavian artery. In complete obstruction the blood is carried to the aorta beyond the obstruction by greatly dilated anastomotic vessels, branches of the subclavian artery, especially scapular and internal mammary arteries, conveying blood to the lower aorta through the intercostal and superficial epigastric arteries, the defect is commoner than has been thought, it usually escapes diagnosis. Sometimes the first sign found is the pulsation of one of these anastomotic vessels, enlarged and tortuous beneath the skin in the back or axillary region of the chest, or a local systolic murmur heard when by accident the stethoscope rests over such a vessel in examining the chest. More often the anastomotic vessels are found only after deliberate search has been made for them. High blood pressure (200 mms Hg) is the rule in the brachial vessels, and

ese and the carotid arteries often pulsate with unusual vigour. The carotid arteries at the base of the neck are often dilated. A systolic murmur at the aortic cartilage is frequent. The post manubrial dulness of aortic dilatation may be found and X ray examination may show dilatation of the aorta or its main branches. Whenever high blood pressure is discovered the femoral pulse should always be examined. In cases of coarctation this is absent or it is feeble and occurs after the pulse at the wrist. Signs of ventricular enlargement are found clinically or by orthodiagraph in about half the cases. The condition is quite compatible with a long life of arduous work. The average life is about thirty years. Most of the patients are without arterial symptoms the heart carrying on its functions well for many years. The patients succumb to a variety of maladies. Cardiac failure and sudden death usually from rupture of the aorta are almost usually the most frequent causes and account for about half the cases. Others die of cerebral haemorrhage, infective endocarditis, renal disease and of other and diverse causes.

OPEN FORAMEN OVALE

Persistence of the foramen ovale occurs in 20 to 30 per cent of all hearts. Patency of the foramen a term usually applied to a defective septum and wider opening is much rarer. So far as I am aware it does not give rise to any direct physical signs by which it may be recognised. It can hardly produce a systolic murmur because the auricle contracts before systole begins. A presystolic murmur has been described but must be of great rarity. The open foramen does not occur in cyanosis and is not known to shorten life. Clots liberated from the systemic veins are sometimes carried through it into the systemic arterial circulation thus missing the lungs (crossed embolism).

DEFECTIVE INTERVENTRICULAR SEPTUM

The septum of the ventricles may be open in its upper membranous part without other defect in the heart. When the aperture is a full one the right ventricle is increased in size and the X ray boulette tends to be circular in outline. A loud and long systolic murmur often accompanied by thrill and presumably caused by blood passing from left to right ventricle is maximal about the 4th rib near the sternum. The murmur is usually well heard over the precordium but is not conducted above the base of the heart or to the left axilla. Cyanosis is unusual and there are few symptoms.

The septal defect may be large and involve in greater or less degree the muscular partition, there may be no septum. Admixture of the right (venous) and left (arterial) blood happens according to the extent of the defect. The opening is usually associated with other defects, such as malposition or transposition of the great arteries, or with pulmonary stenosis, or with suppression of the pulmonary artery. In such the work of the right ventricle is greatly increased, and this chamber is of the size and thickness of the left. In many the admixture of venous and arterial blood is free, septal defects are in fact the only common cause of cyanosis in congenital heart disease, the mixed blood being expelled into the aorta. The resultant cyanosis is of the kind previously described as arising centrally (page 36), it may be transient and is then often slight, or it may be persistent and deep. In the last case clubbing of the finger tips, and polycythaemia up to eight or ten million red cells, or exceptionally more, are usual, breathlessness is conspicuous and bodily growth is defective. When clubbed fingers are fully developed, they are bulbous at their tips and the fingers are conspicuously cyanotic. In these cyanosed cases too, epistaxis, haemoptysis, or cerebral haemorrhage may occur, and sometimes alarming cerebral symptoms appear in the form of attacks of an epileptic nature or of coma.

In cases of septal defect, deep and persistent cyanosis, breathlessness and enlargement of the heart are the chief adverse signs, and yet some of the cyanotic cases live on into their forties. Infective endocarditis, pulmonary tuberculosis, and other infections often terminate life, some of the cases die of acute heart failure.

UNEEXPANDED INFUNDIBULUM (PULMONARY STENOSIS)

Amongst congenital malformations, lack of full expansion of the infundibulum, the half of the right ventricle that forms the outlet to the pulmonary artery, is one of the most frequent. When the defect is slight there is a little constriction between right ventricle proper and infundibulum, the moderator band forming one side of the constriction. When less expanded, the infundibulum forms a small separate chamber, united to the ventricle by a narrow orifice below and to the pulmonary artery above. Pulmonary stenosis, so called, in which the valve cusps are fused, is probably associated with a complete lack of infundibular expansion, in these cases a strong membranous shelf is found at the origin of the pulmonary artery, perforated centrally by a small circular aperture. The condition is comparatively rare. In general the less expanded the infundibulum, the

smaller is the stenosed aperture through which the blood has to pass out of the right ventricle, the nearer is the stenosed aperture to the pulmonary valve and the more apt is it to be combined with a patency of the interventricular septum. Other malformations are often associated. In high grades of stenosis, much blood may be transferred from the right to the left ventricle, and deep and persistent cyanosis appears with its accompaniments, finger clubbing, polycythæmia, and breathlessness. In high grade stenosis also, the right ventricle hypertrophies greatly and gives corresponding physical signs. Curiously enough the pulmonary artery is often dilated in these cases (Fig 45).



FIG 45.—Ortoid agram (1 lb) aged 1st weight 1 lb. Probably congenital pulmonary stenosis. Systolic thrill third left space attacks of cyanosis. The patient has been rotated bringing the left shoulder forward a little. The base of the greatly dilated pulmonary artery (P) is shown. The apex of the heart is probably formed by the right ventricle.

The stenosis displays itself through a harsh systolic murmur and thrill maximal to the left of the sternum, at the 2nd, 3rd, or 4th costal cartilage, but heard over a wide area of the upper chest, as high as the clavicles. It is transmitted to the interscapular region but not into the neck, very occasionally an early diastolic murmur is heard to the left of the sternum, and is caused presumably by pulmonary regurgitation. The prognosis is variable, some patients living beyond thirty or forty, but the death rate is much above normal, and especially in those much cyanosed and presenting much cardiac enlargement. Infections often terminate life. Despite the greatly increased work of the right ventricle, failure with congestion is infrequent.

DIAGNOSIS

The diagnosis of congenital malformation of the heart should not be made upon the basis of a systolic murmur in the pulmonary area. Such murmurs are amongst the commonest heard over the heart, they may be soft or harsh, and may be heard over the 2nd, 3rd, or 4th left cartilages. In the great majority of instances the murmur is without significance, but may be used as a reminder that the base of the heart should be felt for thrills. A diagnosis should not be made though malformation is often rightly suspected because a loud murmur of unusual distribution is found over the

heart It is clear that the only sign may be a systolic murmur but in such cases no great harm is done if the underlying condition remains undiagnosed The presence of a basal thrill especially if this is maximal to the left of the sternum is distinctly helpful Cyanosis or a history of cyanosis in cases of the kind occurring on and off since birth usually decides the diagnosis Deep cyanosis that is proved to be of central origin and that is known not to be caused by the presence of an abnormal blood pigment is almost diagnostic in itself provided there is no venous engorgement and no disease of the respiratory system Otherwise the recognition of congenital malformations depends upon the presence of one of the characteristic groups of signs that have been described earlier

PROGNOSIS AND TREATMENT

It should be clear from what has been said already that the prognosis in cases of malformation of the heart regarded generally is by no means so grave as it has often been thought There are certain conditions which make extrauterine life impossible or so precarious that it is not maintained for many weeks or months but for the rest the subjects of congenital heart disease although specially prone to pulmonary tuberculosis and infective endocarditis often live on well into adult life are active workers and bring up families In attempting to prophesy in cases in which an exact diagnosis fails to be made the main considerations are the presence or absence of persistent cyanosis and breathlessness and enlargement of the heart and its degree

The malformations are not open to curative treatment The treatment is for symptoms as these arise The patient's activities are to be guided by the simple but sound principle that he lives within the limits of his exercise tolerance

CHAPTER XXIX

CHILD BEARING ANAESTHETICS AND OPERATIONS

CHILD-BEARING

HEART disease amongst young women is almost always rheumatic in origin. Questions relating to pregnancy apply almost exclusively, therefore to this form of heart disease and especially to cases presenting mitral stenosis.

RISK OF CHILD BEARING

Advice is often sought by young women with heart disease as to the risk they will run by bearing a child. Often too advice should be given unasked to married women with heart disease. Child bearing entails an increased load upon the heart because the circulation has to serve both mother and child and the mother in moving must carry extra weight. During normal labour a relatively heavy load is thrown upon the heart intermittently for a period of many hours. In late pregnancy the heart is displaced upwards and its action is impeded in unknown but probably not high degree. The extra work involved by carrying is negligible in the first and almost so in the later months of pregnancy, the extra weight of the child is but a small fraction of the whole weight of the body. The work done in labour varies: it is greater in primipara, it is rarely great enough in normal subjects to produce more than transient breathlessness and pulse quickening: it is largely controllable by the management of labour. If the patient belongs to the group in which valvular disease is the chief finding, there being little enlargement, exercise tolerance being good or fair, work of this order is not harmful and the risk proves in fact to be slight, for most married women of this class pass through one or more pregnancies without trouble. There is even in these, however, an element of uncertainty. Judgment is being given not as to advisability of undertaking certain acts at the moment but of undertaking them in nine or more months'

time, and the patient's condition may not then be what it now is. Thus, the possibility of a renewed rheumatic infection needs to be considered on the basis of the past history, and a patient suspected to have suffered infection or reinfection recently should be advised to postpone decision for a year or more. Cases with signs of enlargement of the heart and poorer exercise tolerance should always be advised that it is unwise to become pregnant. The urgency of desire for a child will sometimes impel women to accept a risk that would not otherwise be taken, the risk being realised, it is for their decision. Women presenting auricular fibrillation run more risk during pregnancy than those without, the muscle of these hearts is in general less dependable, and work increases the rate of the beating unusually. Those with fibrillation, and those with large hearts and poor tolerance of exercise, should be told that the risk is much too grave, and every effort made to dissuade them from facing it.

Where a second pregnancy is in question, experience of the first is one of the most valuable guides to the probable course of the second.

FAILURE AND CHILD BEARING

Because most women with rheumatic heart disease fail to seek advice, or, receiving it, are unable or unwilling to follow it, a number of those who have serious cardiac conditions give birth to children. The adverse effects upon the heart, and the increased death rate consequent upon child bearing, tend perhaps to be exaggerated, and for two reasons. Firstly, heart failure will develop in many rheumatic heart cases during the usual child bearing age whether children are borne or not. Secondly, many women attribute ill health to child bearing when they have no good reason for doing so. Nevertheless, there is no doubt that the mortality amongst heart patients who marry is unusually heavy, and the effects of given pregnancies upon the subsequent condition are often obvious, or life is clearly brought to an end by a pregnancy. Patients who develop congestion in the early months would almost certainly have developed the same degree of failure without becoming pregnant. Failure develops chiefly in the later months. In cases previously presenting normal rhythm or auricular fibrillation the onset may be gradual, with increasing breathlessness. Or the onset may occur abruptly with the appearance of a paroxysm of tachycardia or of auricular fibrillation, these cases are often called instances of acute dilatation of the heart. When congestion comes early or late it tends continually to increase and, if it becomes severe, the patient often dies during

the last few weeks of pregnancy or in the puerperal period. Death may result from progressive failure pulmonary oedema embolism or infective endocarditis. If the patient survives labour congestion usually disappears quickly a clear indication that pregnancy was responsible. Such cases may return to a condition seemingly no worse than that preceding pregnancy but often they are more breathless than previously and amongst those who return to work congestion soon redevelops.

MANAGEMENT OF PREGNANCY

When a woman with heart disease takes to carrying a child she should cease to do some at least of her work. Thus in the later months all being well with her she should cease to do her house work to take any form of brisk exercise or to undertake serious physical efforts of any kind. She should spend hours resting each day increasing their number as her time approaches. The amount of exercise and the amount of rest taken will be controlled by symptoms she must live without any distress of breathing.

The appearance of venous congestion with or without auricular fibrillation in the first three months of pregnancy is an adequate reason for emptying the uterus as soon as failure has been treated.

Congestion is more apt to display itself in the later months though it is then much more difficult to recognise in its early stages. Oedema of the legs is of little value as a sign a congested liver is displaced upwards by the uterus and its edge and lower percussion border may remain indistinct. Diagnosis then depends chiefly upon breathlessness upon colour and upon the condition of the veins. The most important of these three is breathlessness at rest persisting during sleep. Cyanosis is also a valuable sign if general or if seen in skin that is warm so are persistent rales at the bases of the lungs. In examining the veins an allowance must be made for the increased pressure in the abdomen normally the veins of the neck are distended more and to a higher level in pregnant than in non pregnant women. Patients should usually come under strict nursing care before actual signs of venous engorgement appear. (Auricular fibrillation is to be treated with digitalis in the ordinary way without any restriction of dosage. The patients who react well may be allowed to go to full term seemingly natural labour is less harmful than induction. Every care should be used to make labour as effortless for the woman as possible. An alternative is Caesarian section now a good deal practised in these cases. The cases are first fully treated

for cardiac failure. In those reacting well, the section is undertaken as soon as there is a good prospect of the child surviving, and can be carried through in most cases with complete success. In those failing to react well, delivery should be as soon as possible. An advantage of Caesarian section is that it saves the mother from the last and most dangerous weeks of pregnancy. The operation is often combined with a procedure that sterilises, to rid the woman from further risk of pregnancy. The justification or otherwise of this procedure is more an ethical than a medical question.

✓ Patients presenting signs of venous congestion should not suckle their babies, but there is no reason to terminate lactation in simple valve cases.

ANAESTHETICS AND OPERATIONS

The dangers involved in anaesthetising and in operating upon patients suffering from cardiac disease have been much exaggerated in the past. Surgical intervention in cardiac cases is often inevitable to save lives that are threatened, when many such operations are performed successfully confidence grows and operations of less immediate urgency are undertaken. Experience has widened greatly, and the results of very many hundreds of operations on patients suffering from heart disease have been recorded. The actual risks are now known for all the chief forms of heart disease. If we include all fatalities that can be connected even remotely with the heart such as any case of pneumonia following operation then the mortality attributable to the cardiac condition is about 5 per cent, this statement applies to unselected cases of conspicuous heart affections such as mitral stenosis aortic disease auricular fibrillation and including all more serious maladies it does not include trivial or doubtful cardiac maladies it applies to groups of patients containing equal numbers of minor and major operations.

It will simplify if it is here said that there is no condition of the heart, other than one which is itself immediately jeopardising life, that prohibits operations being undertaken to meet a grave emergency, and that there is none that forbids trivial operations, including tooth extraction, when these are necessary, under local anaesthesia.

The advisability or otherwise of anaesthetising and operating upon cardiac patients is decided by a close consideration of the disease from which the patient is suffering to determine actual risk. This risk is then balanced against the benefits to be expected from

operative interference. The presence of cardiac disease increases the risk of operations but in some cases the increase of risk is very slight and in other cases it is great.

In the case of uncomplicated mitral stenosis or aortic regurgitation or in that of any other subject able to follow his ordinary daily routine the extra risk introduced by the cardiac state is practically negligible in contemplating surgical interference. In cases of the kind, therefore operations such as gastroenterostomy or the radical cure of hernia may be undertaken under general anaesthesia without hesitation from the standpoint of the heart.

Unselected cases of auricular fibrillation, most of which are complicated by mitral stenosis and by enlargement, present an unexpected mortality of only 2 per cent and here mortality can be reduced if operations of pure convenience are debarred in the more serious of the cases. One of the most frequent operations performed in this group is that of thyroidectomy; the mortality is small. Naturally all cases of fibrillation should undergo preliminary and thorough medical treatment with digitalis before operations other than urgent ones are performed. The case of fibrillation and of flutter are almost the only ones to which digitalis should be given before operation, atropine should not as a rule be given to either.

Paroxysmal tachycardia does not contraindicate operations under anaesthesia though the onset of a paroxysm would naturally postpone any that was not extremely urgent.

In hypertension and in breathless cases of arteriosclerosis the ordinary risks of anaesthesia and operation are raised about 5 per cent. The subjects are elderly and post operative broncho pneumonia which is the chief cause of death is a little commoner than in more normal subjects of the same ages. Prostatectomy is a frequent operation in this group and is well borne as a rule. Patients presenting definite evidences of renal insufficiency show a high mortality.

The presence of mild angina pectoris does not contraindicate general anaesthesia but should rule out operations of pure convenience and these only. We come to cases running greater risks.

In severe angina, syphilitic aortitis and aneurysm operations that are not urgent are rarely advisable, for amongst these patients the added mortality is 10 to 20 per cent. In severe angina chloroform has sometimes been given with impunity, to stop pain.

In no case of coronary thrombosis unless there has been good recovery and many months have elapsed since the event, should operations other than those necessary to save life be performed.

Cases of failure with definite venous congestion will sometimes require surgical treatment in emergency. Otherwise the cases should be treated thoroughly and as bed cases before operations are considered, and the treatment should be prolonged for weeks or months after signs of venous congestion have disappeared. Operations definitely calculated to improve general health may then be undertaken, but not operations of convenience. It is particularly to be noted that elderly patients with venous congestion are not good subjects.

Heart cases stand general anaesthetics well, ether is well tolerated by the young and middle aged, but should be avoided in older subjects that are bronchitic. Chloroform is well taken. There is no objection to carbon dioxide in amounts sufficient perceptibly to deepen breathing. The use of spinal anaesthesia whenever possible, because the case is a heart case, is of doubtful wisdom. A more important consideration than the kind of anaesthetic used is probably the competence of the anaesthetist. Quick anaesthesia that avoids all possibility of struggling is essential. Similarly speedy emergence from the anaesthetic is most desirable.

In heart cases of any kind during natural labour gas and oxygen or chloroform, the latter lightly, can be given with advantage. With gas and oxygen asphyxial signs must be avoided quite strictly. Full general anaesthesia is used for Caesarian section.

It is important to realise how the fatalities occur. Neither anaesthesia nor operation imposes so far as I am aware any burden or stress upon the heart. When cardiac patients submit to operation under general anaesthesia those that die unexpectedly do not usually succumb under anaesthesia but in the hours or days that follow. The reason for many of the deaths is that patients who are suffering from the graver forms of heart disease are relatively intolerant of septic poisoning and exhaustion. Thus post operative infections such as broncho pneumonia, the commonest cause of death in elderly heart cases, or a large wound area left infected after operation, are not well borne, neither is the intoxication that sometimes follows acutely upon thyroidectomy, these complications are ill tolerated because they raise the heart rate while rendering the muscle less efficient for its work. Coronary thrombosis and cerebral thrombosis are not infrequent after events in elderly subjects, they cause fatalities, which are associated perhaps with feebleness of circulation. In mitral stenosis with or without auricular fibrillation the fatal event may be embolic, and in these and high tension

cases pulmonary oedema may supervene. Patients who present failure with congestion or are threatened by it before operation may progress to fatal venous stasis in the few days that follow.

Sudden catastrophes during anaesthesia or operation occur but are unusual. Much of what is said and thought about fatalities in cardiac disease under anaesthesia is due to a very widespread misconception. It is customary to attribute a large proportion of deaths occurring under anaesthetics to acute failure of the heart. There is good reason for doing so. Thus it has been shown that under light chloroform anaesthesia fibrillation of the ventricles is the usual cause of death. It is quite erroneous to suppose that the fault lies in these cases in the heart; it lies either in the method of administering or in the unfortunate choice of anaesthetic. This accident is particularly frequent in the case of children; it is a phenomenon displayed by perfectly healthy hearts. There is no evidence to suggest that any reduction of such fatalities will result by excluding from anaesthesia patients in whom a systolic murmur, extrasystolic irregularity, or other isolated sign attributable to the heart has been detected. And the exclusion of cases presenting serious heart lesions could not reduce this mortality appreciably for it is clear that only a very small proportion of patients dying under anaesthetics are properly to be regarded as heart cases. A real understanding of the practical problems will not be reached by enquiring what proportion of deaths under anaesthetics and operations can conceivably be attributed to the heart; but by enquiring what is the ascertainable risk in cases presenting clear evidences of heart disease. That there is an increased risk from anaesthesia when the heart is recognisably unsound is clear enough but where anaesthetic deaths in general are concerned this becomes a trivial issue. It is an important issue when it concerns the individual heart case requiring anaesthesia for operation and it is from this standpoint that the problem has been discussed in this chapter.

CHAPTER XXX

DIAGNOSIS, PROGNOSIS, AND TREATMENT

DIAGNOSTIC TERMS AND SUMMARIES

IN medicine generally diagnostic terms are often indispensable. They have particular value when they name specific diseases running well defined courses and calling for definite practical measures. The name "diphtheria" at once suggests isolation and specific treatment. "Cancer of the stomach" is a name that not only conveys a well-defined clinical picture but a disease running to a fatal termination.

The satisfactory use of short diagnostic terms in cardiac disease becomes increasingly difficult. The system of names for cardiac disease in the last century was mainly anatomical, it was this system which gave such undesirable emphasis to defects of the valves, the harmful effect of which, though now lessening, is still with us. These names of valve affections are quite inadequate. To speak of "aortic disease" conveys neither the origin of the malady, nor the prospect of life, nor the need of treatment, the malady may have arisen as part of rheumatic carditis or of syphilitic aortitis, the man may be moribund or in almost perfect health and requiring no treatment. In this old nomenclature there were few or no names that had the striking and helpful significance of those previously mentioned. We now possess a number, for example, "subacute bacterial endocarditis", "rheumatic carditis", "syphilitic aortitis", "auricular fibrillation", "auricular flutter", "coronary thrombosis", "hypertensive failure". Each of these frequently fills a definite and most useful purpose. This list might be lengthened with advantage. The terms "aortic disease", "mitral stenosis", in speaking of cases, might profitably be retained and confined to those instances that are almost uncomplicated, cases up and about their business in life.

Cases that have general venous engorgement of cardiac origin should be called "failure with congestion" or, shortly, "congestion", irrespective of other cardiac findings. The term "cardiac asthma"

might be employed perhaps for cases of failure presenting nocturnal attacks prominently. Short terms of the kind we must have for every day convenience. A term to be serviceable must emphasise the main feature of the case and this may be a mechanical defect a bacterial invasion a degenerative process a disorder of function it is a diverse nomenclature. No system of diagnostic terms has yet been devised which really fills this need of briefly summarising cardiac cases this is so because the many structural defects and the very many disorders of function that occur combine in an endless variety of ways. For the same reason the systematic classification of heart diseases under names is scarcely possible. It would seem that the attempt to find an inclusive series of terms must be abandoned. Such brief terms as those that have been named will continue to be used owing to their convenience in everyday conversation. The further practical need is of a simple system whereby we can place on brief record all the essential features of a case that are required for prognosis or for treatment so that these may be gathered at a glance. Such a system I have now long used and found adequate to its purpose. The summary of the case is recorded in eight lines.

	a	b		d	e
1 Exercise tolerance	None	{ Poor } angina	Good	Poor	Poor
2 Congest on	+				
3 Enlargement	Mod	Def		Cons I	Mod
4 Rhythm	A F	P A			
5 Valves	U S		Syst M	A R free	A R sl
6 Infect on					Subac to b
7 Vessels		{ B P } 100		Aneur	
8 Et ology	Rh			S	

The table here given provides a synopsis of five patients

- a Breathless at rest congestion moderate enlargement of heart auricular fibrillation mitral stenosis rheumatic origin
- b Exercise tolerance poor through angina slight definite enlargement pulsus alternans hypertension
- c Good exercise tolerance systolic murmur in mitral area
- d Poor tolerance considerable enlargement free aortic regurgitation aneurysm syphilitic
- e Poor tolerance moderate enlargement slight aortic regurgitation subacute bacterial endocarditis

The short dotted lines indicate that in the corresponding spaces

there is nothing significant to record The degrees of enlargement here referred to are defined on page 116

PROGNOSIS

A NORMAL HEART

If after carefully examining a patient and finding nothing definitely wrong we are to fear that vital things are still undiscovered, we shall remain at the mercy of every rumour having patient or patient's heart as its source, and be incapable of responsible decisions It is our patient who will suffer from the resulting diffidence and overcautiousness, these attributes in a medical man turn many healthy people into invalids But confidence, to be justified, must rest on knowledge and experience it is not to be substituted by mere temerity To be able to proclaim that a heart is normal upon a sufficiently sure basis is a very important accomplishment

The most valuable indication we possess of a heart's essential soundness is its power to engage fully in its work without complaint No patient having normal capacity for bodily exercise has grave heart disease, for bodily exercise calls upon the heart for extra work, it challenges the heart's reserves and, if it can soon exhaust these, it will induce unusual breathlessness, or, in the event of the heart's own blood supply proving inadequate, cause anginal pain Thus the presence of good exercise tolerance can be relied upon to exclude all the more serious maladies, though it should be understood that full capacity for work cannot be gauged always and entirely from the patient's own statements, valuable as these usually are Actual experience will show that only exceptional patients, having such full capacity, will present clear signs of disease

Because there are these exceptions, full tolerance of exercise is an insufficient basis upon which to declare the heart to be sound It is also necessary to exclude definite enlargement of the heart, definite evidence of valve disease in the form of mitral stenosis and aortic disease, congenital malformations, auricular fibrillation and high blood pressure, though the last two are so very rarely accompanied by normal exercise tolerance that it is hardly necessary to name them A short physical examination suffices to exclude the states of disease named, particular heed being given to the valves, for it is this form of disease which is most likely to consort with normal tolerance The heart must be listened to while the patient stands, lies, and lies after exercise on the left side Much enlargement of the

heart and progressive mischief syphilitic or rheumatic are incompatible with good exercise tolerance though the physical examination is an extra safeguard There must be a brief enquiry for paroxysmal disorders of heart action and these also excluded the heart may then be proclaimed sound/A systolic murmur over any part of the heart should lead to a sharper search for more definite evidences of disease and in young people to a close enquiry for recent rheumatism Neither the systolic murmur nor a modification of a heart sound nor a sound interpreted as exocardial nor an unusual form of electrocardiogram should deter us from pronouncing the heart essentially sound in these circumstances In this connection it should be noted that an isolated sign rarely has much significance Indications of disease are grouped together almost always The foregoing are the criteria that were used on a very large scale in judging men fit for active service during the Great War they proved themselves to be reliable

A more difficult case is where exercise tolerance is reduced in the absence of physical signs of disease in heart lungs or other organ for very many such patients have normal hearts Here for the time being we must proceed by rule of convenience declaring the heart sound in the young but holding it under suspicion or to blame in older subjects (see page 163)

CHRONIC HEART DISEASE AND PROGNOSTIC GROUPINGS

In considering on what general lines an opinion of the expectation of life is to be formed in a case of chronic heart disease we may profitably ask whether we are led by subdividing patients systematically into groups All cardiac patients may be divided up into important groups according to the manner in which the circulation is carried on They could be again subdivided according to the size of the heart unenlarged or enlarged in one of several degrees They could be subdivided again and again according to the nature of the valve defect and its degree according to etiology to blood pressure and to heart action Subdivision along these lines will be found to provide several hundred different classes of cardiac case The attempt has but one result namely to convince that prognosis can never be practised in this way the finer the subdivision the more cumbersome and impractical it becomes It is imperative to proceed upon a broader basis and to emphasise those features of cardiac disease that are of chief prognostic significance

Stated in the order of their importance, and omitting active in

fection for purposes of simplicity, the following factors mainly govern prognosis

(a) The state of the circulation at rest, or under exercise as judged by breathlessness, and the presence or absence of anginal pain in similar circumstances. These considerations are essential and intimate parts of any sound prognostic system, they are so important that prognosis of chronic disease carried through purely on this basis would not prove very unsatisfactory

(b) The size of the heart

(c) The presence or absence of syphilitic disease of the aorta

(d) The presence or absence of gross disturbances of heart rhythm (for example, auricular fibrillation), the presence or absence of high tension

(e) The state of the valves

Before proceeding to deal with the practical problem of prognosis from this standpoint, we must first understand quite clearly what is required. No enquiry, however minute, would carry our prophetic powers to such a pitch that we could state definitely how long a patient has to live. Fortunately it is not asked. We are asked to give an opinion along the following simple lines

1 Does the expectation of life seem normal?

2 Has the patient many years to live?

3 Has he a number of years, say ten, a little more, or less?

4 Has he but a few years?

5 Is life to be measured in months, a precarious life of perhaps one, or perhaps two years?

6 Is the patient in imminent danger or actually moribund?

These questions can usually be answered accurately. The first has been answered in enquiring how a normal heart is recognised. The remaining questions may be answered in tabular form. Sufficient types are here included to illustrate fully, the attempt is not made to cover all possibilities. It is assumed that the cases are all under full and proper treatment and that none are suffering from active rheumatism, or from infective endocarditis.

1 *Many years' expectation*

Valve disease in the young adults, with little or no sign of cardiac enlargement, no auricular fibrillation, and good or fair exercise tolerance

- 2 *A number of years' expectation (about ten years)*
 Angina of effort early crises
 High tension little or no sign of enlargement, fair tolerance
 Auricular fibrillation with or without valve disease, but little enlargement and fair tolerance
 Enlargement moderate with or without valve disease, but with fair tolerance
- 3 *A few years' expectation (about three to six years)*
 Angina of effort on walking a short way (with or without previous coronary thrombosis)
 Syphilitic aortitis
 High tension with moderate enlargement or angina
 Enlargement great or moderate with poor tolerance
 Auricular fibrillation and moderate enlargement
 Failure with congestion (relieved by treatment)
- 4 *Living precariously up to one or occasionally two years*
 Enlargement great congested
 High tension and cardiac asthma
 Congestion unrelieved or in syphilitic aortitis, in high tension following angina or after coronary thrombosis
 Angina unrelieved by rest
- 5 *In imminent danger*

Oedema of lungs	}	Often recovering
Coronary thrombosis (recent)		
Adams Stokes syndrome (epileptic state)		
Leaking aneurysm	}	Rarely or never recovering
Anginal state		
Gross failure with congestion unrelieved		

It is not intended that this table shall form a precise guide to prognosis in individual cases each of which must be considered upon its own merits as discussed in earlier chapters. It is only intended that it shall broadly and clearly illustrate the relative importance of the chief prognostic factors previously named, and how these come to be used in answering the relatively simple questions that are asked. It will be observed that apart from the first group of cases, valve disease can be made to take little part. That is so because as we proceed to more and more serious categories other factors become more and more decisive, and valve disease becomes relatively unimportant.

CONVERSING WITH THE PATIENT AND HIS FRIENDS

To a sick man his doctor's visit is the chief event of the day. Cardiac patients react quickly to the atmosphere surrounding them, they must be helped to keep or gather courage. Cardiac cases are naturally sanguine, depression, when it comes, usually derives from others. It is the doctor's plain duty to enter the sick room with cheerfulness, to give counsel thoughtfully but confidently, and to leave behind as he goes an appropriate word of encouragement, to comfort the patient in his waiting, to help his suffering, or to allay his fears. A countenance of gloom is as out of place in a sick room as is a coffin.

In speaking with a patient or his friend about the nature of the malady, truth of statement is a first essential. It is not always expedient to volunteer the whole truth, but answers must be given to direct questions, evasions usually defeat their purpose. When a man in his right mind demands to know if he is dying, as seldomly he does, he should have his answer, and if it is definitely adverse, it can be conveyed compassionately. Spontaneously to acquaint a patient that his life is in grave danger is rarely necessary or advisable. It sometimes becomes desirable to convey to him, directly or through his family or legal adviser, that it would be a simple act of wisdom if he set his house in order.

To patient or to patient's friend simple and unmistakable language should be used, veiled statements are too apt to be misinterpreted. The words must be chosen thoughtfully. The word "disease" should never be used, disease of the heart conveys at once to most patients the idea of something incurable and threatening. The language should be as untechnical as it can be made. A patient has a right to be told what ails him, if he so desires, in terms that convey an idea of the magnitude and significance of the trouble, he has no right to technical diagnostic terms, and it is very rarely wise to parade these before him. Accurate information cannot be conveyed by means of strange words, which to unaccustomed ears bring unintended meanings, and for many the word once caught up becomes a matter requiring search in medical books, or a topic of debate with other patients. Thus the word "angina" should never come first from a doctor, if it is understood at all it will convey in almost all instances an ominous meaning, yet the prognosis varies up to fifteen years. The word "dropsy" is to be avoided, and the adjective "malignant" eliminated completely from the medical man's vocabulary. The

valves should not be named, nor murmurs mentioned. These names and details should be of no concern to patients. Methods of examination likewise require little or no explanation, a clear example of the unhappy effects of unnecessarily disclosing technical detail is the well known instance of high blood pressure readings.

It is rarely desirable to speak of catastrophes that may arise out of over exertion, though a warning that certain acts are foolish and

that he or she may drop dead at any moment", a not very unusual phrase is guilty both of cruelty and folly. Apart from its total lack of consideration sudden death in heart disease is never so inevitable as to warrant such a statement. In grave angina, and in thoracic aneurysm, sudden death is frequent, but in most other common cardiac maladies it is comparatively rare. In the named instances, and in these only if the question is directly asked by the patient, it must be answered that there is the chance, such information should not be volunteered to him, though it is wise to tell a near and discreet relative that the life is a precarious one. It is important that the motive should be correct, that the statement should be made chiefly out of solicitude for the patient's safety, or withheld out of consideration for the family's mental comfort. A medical man who fails to warn relatives of a possible sudden ending may be a little blamed if and when the event occurs, but a doctor who sets out to safeguard himself against all risks of this kind, deliberately prefers his own to others' interests. Death without warning disturbs a patient's family enough, its long, and often unnecessary, anticipation inflicts more suffering.

A few well chosen words are far better than a diffuse statement out of which any detail may be seized upon and emphasised, while the real substance is forgotten. It is most necessary to be brief and to keep to essentials.

Every man will have his own way of saying things, it is not the precise form of expression that matters, it is the outlook. A doctor's watchword should be optimism and not pessimism, he should be hopeful, and not fearful of the unknown, and experience will show that this profits both his patients and himself. A statement of prognosis will be formed on the lines laid down elsewhere, and if the

prospect is uncertain, then it is the brighter prospect that should, within reason, be presented to the patient

TREATMENT

The treatment of heart disease consists chiefly in managing cardiac failure and angina pectoris of different grades of severity. The treatment of a patient convalescing from failure with congestion has been discussed at length, so has that of angina pectoris and of hypertension. Special points of treatment have been considered in relation to the several conditions to which they apply. Thus most of the ground has been covered. There is still something that may be said usefully to ensure that the management of heart cases that are up and about should be understood. There are supplementary statements, which, included here, save their reiteration in several places in this book.

Healthy Habits—Experience has led us to recognise healthy habits of living. Very prudent people live quietly and moderately, they have their simple daily routine of work and pastime, enlivened by occasional excursions and entertainments, social gatherings and visits to and from their friends. They are temperate in their eating, taking no more than will maintain them in robust health, arranging their diet to consist chiefly of plain good food, relieved occasionally of its monotony by a more elaborate, but not heavy meal. They are regular in their habits of work and exercise, and in their meal times. They are strictly temperate in their drinking. They control their emotions and their passions. They avoid all forms of excess. They use tobacco little if at all. They welcome the freshness of abundant air and open spaces, delighting in the feelings of invigoration that accompany active exercise; they love the warmth of sunlight playing on their skins, and the sleepiness of healthy fatigue. These are habits that few people in industrial countries now adopt, that fewer still maintain. The cares and distractions of an increasingly complex life, indoor or sedentary occupation, advancing years and decreasing energy, interfere less or more, and the prudent rules are neglected, neglect is the easier because the penalties, owing to the body's great power of resistance to disease, are uncertain in their incidence and often long deferred. *Healthy habits* would become more prevalent if the consequences of their neglect were more immediate and invariable. They are largely the habits of the young, to which many older people rapidly return of their own accord in

convalescing from some acute but passing illness. While immunity from consequences is often long enjoyed by the originally healthy it is not enjoyed to the same degree by defectives. Thus a regular life which may be said to mean unusual prudence in the former is a necessary safeguard to the latter and amongst these we include all patients suffering from well defined disease of the heart.

Avoidance of Infection —For cardiac patients infection is a special danger. Reasonable precautions will not eliminate infection but will reduce the frequency of its recurrence. Crowded rooms and gatherings in public buildings should be avoided especially in the winter season and at such times as infection is known to be prevalent. Those who are infective in the same house should keep their distance and visitors in an infective state should be discouraged by the household. A further measure of safeguard is a scrupulous hygiene of the mouth and throat.

Tobacco —It were best that cardiac patients did not smoke at all but if they will it must be done in very strict moderation. We may not know definitely that long continued smoking is injurious to the blood vessels as some would have us think. It is clear enough however that smoking is responsible in a very large number of its habits for much strenuous coughing that it leads in many people by irritation of the respiratory passages to infections of the throat and of the trachea. Tracheal and bronchial irritation and infection are particularly prevalent amongst but are not confined to those who inhale the smoke and in these a winter cold is often followed by a period of weeks or even months of morning cough. Such coughs and infections are most injurious to many heart cases.

Exercise and Manual Work —In all cardiac cases that are fit to walk about outdoor exercise is beneficial for it encourages health of mind and body. The principle to be laid down for these patients is that they live strictly within the limits set by their symptoms. Patients who suffer from breathlessness may take exercise that makes them breathe a little rapidly from time to time so that they begin to be conscious of the breathlessness but there must be no pushing on to a point of distress or to a point where the sense of breathlessness is not lost almost as soon as rest is taken. So too with anginal pain the man must live strictly within the limit of his pain stopping at once when it is felt and avoiding repetition of the act. In neither one case nor the other must so much exercise be taken as subsequently to bring about unpleasant fatigue. It is useless to attempt to lay down beforehand precisely what a man may or may not be

permitted to do. It is for each patient a matter of cautious trial, and, if a trial is carried too far, then a rest period of hours, or in some cases days, should follow as compensation. Exactly the same principle applies to work, the work should suit the case and be gauged by trial. Cardiac patients can rarely engage in heavy manual labour, such as farm labouring, heavy hammering, or heavy portage, some are fit for moderate manual work such as bricklaying and carpentering, most will find lighter work, such as painting or light fitting or purely sedentary work, most suitable. It is often better to curtail the hours than to attempt to change the work. A patient should not be told to leave his job, unless there is quite clear reason for his doing so, and unless it is thought he will comply. Occupation of mind and body banishes anxiety and cultivates contentment. Children should continue schooling although bodily activities may need strict limitation, and in the young must be controlled closely.

Patients who possess an indifferent tolerance of exercise should report frequently, declining tolerance is a matter for concern. It is unsound to allow congestion gradually to develop without interference, a fall in respiratory reserve or suspicion of venous overfilling should at once suggest a short period of rest in bed. Early treatment by rest is economical, for it often postpones attacks of serious congestion and the long periods of rest which these involve.

Strenuous Acts such as running, walking fast uphill, walking against strong winds, lifting weight in excess of 50 lb, struggling of any kind should be forbidden for all time to patients that have once suffered from cardiac failure, to anginal cases, syphilitic aortic disease and to cases recovered from coronary thrombosis. And the same patients must be instructed to avoid situations in which they may be tempted to perform such acts in emergency. Thus no such patient should walk on broken or precipitous paths or drive an automobile into the country unattended, they must look to others to be the first in helping in accidents or emergencies that are liable to strain the helper.

Diet—The diets here given are intended only to guide and not to be used rigidly, quantity and form of food may be varied within reasonable limits to suit individual cases.

DIET I

GRAVE CONGESTION, ABOUT 800 CALORIES

Breakfast (8.9 A.M.)

- (a) Bread or toast (with or without syrup or honey) $1\frac{1}{2}$ oz. and 1 cup of tea (with milk and 1 or 2 lumps of sugar) 5 oz.
or (b) 1 raw egg in milk 6 oz.

Dinner (1.30 P.M.)

- Minced meat or chicken 2 oz. or fish 3 oz.
Bread $1\frac{1}{2}$ oz. or spinach 2 oz.

Supper (7 P.M.)

- (a) Custard 5 oz.
or (b) Milk pudding 4 oz.
or (c) Bread and milk (milk 4 oz., bread 1 oz., with sugar) 8 oz.

Drinks in small quantities between meals. Total fluids, including tea and milk in above dietary rarely to exceed 20 oz. (570 c.c.s.)

DIET II

CONVALESCENCE, WITHOUT EXERCISE, ABOUT 1500 CALORIES

Breakfast (8.9 A.M.)

- (a) Bread 2 oz.
Butter $\frac{1}{2}$ oz.
1 egg poached or boiled or white fish (not fried) 3 oz.
Tea or coffee 4 oz.
Milk 1 oz.
Sugar 1 2 lumps.
Honey 1 oz.
or (b) Grapefruit or orange
Cereal, dry cornflakes 1 oz. or cooked oatmeal 3 oz.
Milk 5 oz.
Sugar $\frac{1}{2}$ oz.

Dinner (1.30 P.M.)

- (a) Fish, white (not fried) 4 oz. (or meat, chicken rabbit, mutton, sweetbread or game 2 oz.)
Potato 3 oz. or bread 1 oz.
Spinach, tomato, green of cabbage, peas or beans or salad 2 3 oz.
(b) Milk pudding 5 oz.
with cooked fruit (prunes, plums, apricots, baked apple) with sugar, or raw orange, peach or grapes.

Tea (4.5 P.M.)

- Tea 4 oz.
Milk 1 oz.
Sugar 1 2 lumps.

Supper (7 P M), only one course

- (a) Fish 4 oz
Potato 3 oz
- (b) Milk pudding 8 oz
- (c) Chicken 3 oz or fish 4 oz
Salad

Any of these (a) to (c), with bread 1 oz, butter $\frac{1}{2}$ oz

All meals to be taken without fluids. Fluids to be drunk an hour before chief meals, water or, for those accustomed to alcohol, whisky or brandy ($\frac{1}{2}$ oz) and water not more than twice daily, or once with 1 glass sherry, claret, hock with midday meal. Total fluids up to 40 or 50 oz

Diet I is suited to grave failure with congestion, to acute coronary thrombosis, to grave angina pectoris at rest. Such a diet can be maintained up to a week without causing serious loss of strength. As a case of failure with congestion progresses to convalescence, the diet is increased until it corresponds to Diet II.

Diet II is equally suited to other cardiac cases such as angina pectoris, or auricular fibrillation under active treatment with digitalis, that are confined to bed, or to cases of hypotension or senile heart sitting in their chairs. Where convalescence is more advanced and bodily work is undertaken more food will be needed.

Decrease in the starch content of the diet may be required if flatulence is present. Flatulence often results in undue breathlessness and disturbed rhythm, and not infrequently determines anginal pain. It is necessary to insist that no fluid is taken at meal times.

For cases of chronic heart disease that are free from major symptoms there are no special restrictions. In general the meals should be regular, well spaced, and light, highly seasoned and indigestible foods being avoided. It is especially important to cardiac subjects that, while well nourished, they should not become obese, obesity already present should be treated.

INDEX

Paroxysmal tachycardia and 72 153
 Pericardium guarding 117 153
 Pregnancy and 153 269
 Rheumatic fever and 197
 Strain and 70 152
 Adams Stokes syndrome 101
 Adherent pericardium *See* Pericardial
 adherency
 Adrenaline (effects of) 60
 After care of cardiac failure 32
 Age and forms of heart disease 202
 Aged (heart disease in the) 244
 Amyl nitrite 54 57
 Anacrotic pulse 127 220
 Anaemia—
 Anginal pain in 60
 Cardiac failure and 23
 Cyanosis and 39
 Fatty heart in 250
 Anaesthetics in heart cases 271
 Aneurysm 212, 213
 Arterial signs of 220
 Descending aorta 220
 Diagnosis of 224
 Mycotic 188
 Pressure manifestations of 217
 Prognosis of 224
 Symptoms and signs of 213
 Treatment of 225
 Aneurysmal dilatation of left auricle 112
 Angina pectoris—
 Aged people and 244
 Aids to diagnosis of 245 252
 Anaemia and 60
 Anaesthetics in 272
 Aortic regurgitation and 53 126
 Clinical associations of 50
 Coronary arteries and 49 48 50

Angina pectoris—(contd.)
 Diagnosis of 52 54
 Effort and 51
 Hypertension and 50 229 239
 Hyperthyroidism and 60 209
 Nitrites in 54 57
 Oedema of lung and 168
 Paroxysmal tachycardia and 60 73
 Prognosis of 54
 Rheumatic heart disease and 203
 Special cases 59
 Special syndrome 53
 Symptoms of 50
 Syphilis (cardioaortic) and 222
 Thyrotoxic state and 60 250
 Treatment of 56
 Vasomotor angina (Nothnagel) 60
 Anginal pain *See also* Angina pectoris
 Cause of 42
 Coronary thrombosis and 44
 Factors provoking 51 52
 Neurasthenia and 59
 Nocturnal 54
 Pulse and blood pressure and 54
 Tobacco and 59
 Aortic cusp ruptured 125 152 222
 Aortic diastolic murmur 123 124
 Aortic dilatation 213 202
 Coarcted aorta and 264
 Symptoms and signs of 213
 Aortic regurgitation 120
 Angina pectoris and 53 126
 Auricular fibrillation and 200
 Cardiac failure and 15
 Consecutive phenomena in, 125
 Degrees of 123
 Differential diagnosis of 124
 Digitalis in 143
 Enlargement and 106 126 203
 Fallor and 126 189
 Rheumatism and 120 202
 Signs and symptoms of 121
 Syphilis and 120 125 222
 Vasodilatation in, 125

- Aortic stenosis, 127
 Ascending degeneration and 120 244
 Recurrent and 120 203
 Symptoms signs and diagnosis of, 127
- Aortic systolic murmur 123 128 264
- Aortic valve *See also* Aortic stenosis and Aortic regurgitation
 Ascending degeneration of 244
 Bicuspid 263
 Pathological anatomy of 120
- Aortitis *See* Syphilis
- Apoplexy 167 234 232
- Arterial pulse *See* Pulse
 Arteriole spasm 231
- Arteriosclerosis 240
- Arterio-erythrocytosis 124 126
- Atrial degeneration of aortic valve 120, 244
- Atrial fibrillation 179
- Asphyxia causing cardiac failure 23
- Athlete's heart 168
- Atropine in cardiac 70
- Augmentation 70 704
- Auricular fibrillation 81
 Anaesthetics in 272
 Aortic disease and 704
 Cardiac failure and 23 31 63
 Cardiac reactions to 82
 Clinical associations of 81
 Digitalis in 31 86
 Embolism in 81 170
 Hypertension and 732
 Mitral stenosis and 203
 Murmurs in 124
 Paroxysmal 82
 Prognosis of 85
 Quinidine in 88 90 208
 Recognition of 83
 Rheumatic heart disease and 203 208
 Strophilanthus in 88
 Strophilanthus in 80
 Sudden death in 87, 104
 Symptomatology of 87
 Tryptoxenic state and 82 204 259
 Treatment of 86
- Auricular flutter 73
 Diagnosis of 75
 Prognosis of 78
 Rheumatic heart disease and 706
 Treatment of 79
- Auriculo-ventricular bundle 93
- Auriculo-ventricular valves 100 *See also* Mitral valve and Tricuspid valve
- Back pressure theory 141

- Bacterial endocarditis *See* Subacute bacterial endocarditis and Acute bacterial endocarditis
- Ball's ramblin, 206
- Bedstead (cardiac), 26
- Bicuspid aortic valve 183 263
- Blood culture in endocarditis, 190 193
- Blood pressure—
 Age and, 231
 Angina and rise of 54
 Aortic regurgitation and, 121, 122
 Diastolic, 120, 230
 Effort syndrome and 160
 Frequent readings unwise, 236
 High *See* Hypertension
 Lowering 238
 Mean 230
 Measuring 230
 Syncope and 97
 Systolic 121 230
- Blood flow in aorta 33 *See also* Vaso-dilatation
- Boat racing and heart disease, 154
- Brachial plexus (aneurysm pressing on) 220
- Bradycardia 92
 Differentiation of types, 94
 Simple form of 97 99
 Syncope and 99
- Brassy cough 218
- Breathlessness 1
 Bronchitis and 259
 Cardiac failure and 2 7
 Cause of in failure 2 165
 Cyanosis and 36
 Differentiation of forms 163
 Grades of 2
 Hypertension and 233
 In health 2 164
 Nervous 174
 Nocturnal 167
 On effort 2, 3 155
 Posture and 26
 Syphilitic aortitis and 213
 Venous congestion and 16, 154
- Broken compensation 115
- Bronchitis and emphysema 2 1
- Bronchovascular aneurysm 2 1
- Brown induration of liver 46
- Bumbe branch block 247
- Caffeine 30
- Capillary pulsation 40 100
- Cardiac asthma 168
- Cardiac bedstead 26
- Cardiac disease *See* Dilation
- Cardiac enlargement *See* Dilation 117

- Cardiac failure 1 7 22
 After care of, 32
 Aids to diagnosis of 245 252
 Anaemia and 23
 Anaesthetics and 273
 Angina pectoris and 55
 Aortic regurgitation and 155
 Asphyxia and 23
 Auricular fibrillation and 23 31 63
 Bowels in 26
 Bronchitis and emphysema and, 254 255
 Cause of 22
 Coarctation of aorta and 155
 Coronary thrombosis and 23 45
 Cyanosis and 38
 Definition of 1
 Delirium in 15
 Diet in 27 33
 Digitalis and 29 31 86
 Exercise in 33
 Exhaustion in 7
 General statement 1
 Hypertension and 22 233 239
 Hyperthyroidism and 23 259
 Hypnotics and 27
 Infection and 23
 Jaundice in 14
 Muscle change in 23 253
 Oedema in 15 18
 Oxygen in 29
 Pain and 7
 Paroxysmal tachycardia and 23 72
 Pericardial adhesions and 22 181
 Posture and 26
 Pregnancy and 153 269
 Prognosis in 24
 Prolonged work and 154
 Pulmonary disease and 22 254
 Refractory cases of 34
 Reserves and 166
 Rest in 26 32
 Rheumatic carditis and 23 197
 Rheumatic heart disease and 204 209
 Signs of 8
 Sleep in 27
 Splenic enlargement and 189
 Stages of 3
 Subacute bacterial endocarditis and 23 187
 Subtle change in muscle and 204
 Supervision of 33
 Symptoms of 7
 Syphilis (cardioaortic) and 222
 Thyrotoxic state and 23 259
 Tonics and stimulants in 29
- Cardiac failure—(contd)
 Treatment of 25
 Urine in 15
 Valve disease and 22
 Venesection in 28
 Venous congestion and 3 7 15
 Work and, 22 151 154
 Cardiac impulse 107 *See also* Pulsation
 of chest wall
 Diffuse 109 158 160 173 259
 Displaced 108
 Epigastric 110
 Episternal 214
 Heaving 112
 Immobile 178
 In children and lads 108
 Maximal 107
 Pericardial effusion and 173
 Recession of 254
 Ribs and sternum involved by 109
 Cardiac ischaemia 41
 Cardiac margins, 111
 Cardiac oedema *See* Oedema
 Cardiac reserve—
 Age and 245
 Failure and 3 22 156
 Valve defects and 143 156
 Cardiac syncope ■
 Cardiorespiratory murmur 136 160
 Carotid pulsation 14 122
 Carotid sinus reflex—
 Heart block and 94
 In flutter 76
 In paroxysmal tachycardia 79
 Central problem in heart work 1
 Cerebral faints 101 236
 Cervical veins 10
 Cheyne Stokes breathing 166 167 233 248
 Child bearing and the heart 153 268
 Chloroform and the heart 273 274
 Chorea 198
 Clubbed fingers 186 189 192 265 266
 Coarctation of aorta 155 263
 Collapsing pulse 122
 Compensation 114
 Complete heart block 93 94
 Congenital heart disease 262
 Bacterial endocarditis and 183 263 265
 Cyanosis in 36 265 267
 Diagnosis prognosis and treatment of 266 267
 Constrictive pericarditis 177 204
 Continuous murmur 263
 Conveying with patients 281

- Coronary arteries 41
 Angina an 42 50
 Disease of 41
 Fatty heart and 250
 Fibroid heart and 251
 In the aged 244
 Sudden death and 103
 Thrombosis in 43
- Coronary embolism 189
- Coronary thrombosis 43
 Cardiac failure and 45
 Diagnosis of 47
 Hypertension and 43 232
 Oedema of lungs and 45 169
 Prognosis and treatment of 48 40
 Sudden death in 44 48 103
 Symptoms and signs of 44
- Cough—
 Brassy or metallic 218
 Cardiac failure and 7 254
 Treatment of 30 255
- Coupled beating 65 87
- Cyanosis 36
 Bronchitis and emphysema and 255
 Cardiac failure and 15 38
 Congenital malformation an 36
 265 267
- Decompensation 115
- Degenerative changes and rheumatic heart disease 210
- Delirium (cardiac failure an) 15
- Diagnosis (*see various conditions also*)—
 Terms of and use of 275
- Diastole (heart rate and length of) 24
- Diastolic blood pressure 120 230
- Diastolic murmur *See* Murmur
- Diastolic shock 216
- Diet, 285
 In cardiac failure 27 33
- Diffuse impulse 109
- Effort syndrome and 160
- Exercise and 159
- Pericardial effusion and 173
- Thyrotoxic state and 258
- Digital nodes 188
- Digitalis—
 Aortic regurgitation and 149
 Auricular fibrillation and 31 86
 Cumulation of 87
 Flutter and 80
 Maintaining doses of 90
 Massive doses of 88
 Nausea and vomiting from 87
 Normal rhythm and 23
- Digitalis coupling 87
- Digoxin 88
- Dilatation of aorta *See* Aortic dilatation
- Dilatation of heart *See also* Acute dilatation
 Aneurysm of left auricle 112
 Cause of 105
 Efficiency increased by, 152
 Measure of 114
 Prognosis of 117
 Systolic apical murmur and 112 139
 197 245
 Treatment of 118
- Diuretics 30
- Dropy *See* Oedema
- Ductus arteriosus (persistent) 262
- Dulness (cardiac) 110
 Aneurysm an dilatation of aorta and 215
 Emphysema and 254
 Fixed absolute 180
 Pericardial effusion and 174
- Effort syndrome 158
- Embolism—
 Acute bacterial endocarditis and 192
 Auricular fibrillation and 91 170
 Coronary thrombosis causing 46
 Effects of 188
 Pulmonary 170
 Quinidine and 91, 170
 Rheumatic heart disease and 206
 Subacute bacterial endocarditis an, 188
- Emphysema 253
- Endocarditis (bacterial) *See* Acute bacterial endocarditis and Subacute bacterial endocarditis
- Endocarditis (rheumatic) 129 195
- Energy expenditure *See also* Work
 Tachycardia and 23 72
- Enlargement 105 *See also* Dilatation and Hypertrophy
 Aortic regurgitation and, 106 126
 203
 Degrees of 116
 Diagnosis of 116
 Exercise tolerance and 119 204
 Hypertension and 227 232
 In children and lads 109
 Measuring 107
 Pericardial adhesions and 181
 Prognosis of 117
 Rheumatic heart disease and 203
 209
 Symptoms of 106
 Syphilis (aortic) an 222
 Treatment of 118

- Epigastric pulsation 110
 Epistaxis 228
 Erosion by aneurysm 212 213 220
 Errors of measurement 110 132
 Essential hypertension *See* Hypertension
 Exercise—
 Blood pressure and 160
 Effects of 158
 Heart rate in 5 160
 In cardiac failure 33 284
 In effort syndrome 162
 Provoking paroxysmal tachycardia 153
 Exercise tolerance—
 Cardiac reserves and 156
 Decreased 3
 Degrees of 11
 Enlargement and 204
 Prognosis and 277
 Tests of 4
 Exhaustion in cardiac failure 7
 Exocardial murmurs 137
 Expansile pulsation 214
 Extrasystole 63
 Fainting *See* Syncope
 Faintness *See* Oddiness
 Fatigue in cardiac failure 7
 Fatty heart 250
 Fear of death in angina 53
 Fever—
 In bacterial endocarditis 185 192
 In heart disease 190
 In rheumatism 196
 Tachycardia in 69
 Fibrillation of the ventricles *See* Ventricular fibrillation
 Fibroid heart 251
 Flatulence 287
 Flint's murmur 136
 Foramen ovale (open) 264
 Friction (pericardial) 47 174 177
 Friction (pleuropericardial) 175 180
 Fullness of head in cardiac failure 7
 Gallop rhythm 23* 246
 Giddiness—
 Aortic regurgitation and 121
 Effort and 37
 Heart block and 100
 Posture and 97 102 121 159
 Group beating 65
 Gummata of heart 212
 Haemoptysis 170 171 213 217 218
 Headache 228 239
 Healthy habits 283
 Heart block 92
 Intermittence from 65
 Murmurs of mitral stenosis and 136
 Syncope and 100
 Heart rate *See also* Bradycardia and Tachycardia
 Blood pressure and 125
 In exercise 11 160
 In various states 69 92
 Normal 92
 Heart sounds—
 Accentuated *See* Accentuated sounds
 First sound replaced by murmur 139
 Gallop rhythm 246
 In aneurysm 216
 In complete block 94
 Intensified decreased 246
 Reduplicated *See* Reduplicated sounds
 Third sound 199 247
 Heart strain 75 151 161
 Hypertension (essential) 227
 Accompanying manifestations of 231
 Anaesthetics and 272
 Angina and 50 229 230
 Arterial disease and 242
 Cardiac failure and 22 233 239
 Course and prognosis of 234
 Final picture 233
 Hypertrophy in 106
 Oedema and 220 233
 Recognition of 229
 Symptoms of 228
 Syphilis (cardioaortic) and 222
 Treatment of 236
 Hypertension (paroxysmal) 60
 Hyperthyroidism *See* Thyrotoxic state
 Hypertrophy—
 Bronchitis and emphysema and 251
 Cause of 106
 Development of 112
 Electrocardiograms of 112
 Hypertension and 106 227 232
 Measure of 114
 Physiological 154
 Right and left 106 112
 Hypertrophy and dilatation 10,
 Balance between, 115
 Distinction between 113
 Hypnotics in cardiac failure, 27

Infarction of lung 170 255

Infectious—

Avoidance of 284

Cardiac failure and 23

Effort syndrome and 159

Essential hypertension and 233

Paroxysmal fibrillation and 82

Paroxysmal tachycardia and 71

Old age and 245

Inspiratory decline or failure of pulse 179

Inspiratory swelling of veins 179

Intermittent pulse 63

Irregular and disordered heart action 62 69 81 92

Isochaemia (pain of muscle in) 42

Jaundice in cardiac failure 14

Joint pains and effusions 187 194 197

Kidney *See* Renal

Leucocytosis 186 192 197

Liver (cirrhotic and congested) 17

Liver (congested) 14

Lung *See* Pulmonary

Lung (compressed) 175 217

Malar flush 39

Malnutrition in cardiac failure 15

Manual work (regulation of) 284

Marathon runners (heart in) 154

Maximal impulse 107 *See* Cardiac impulse

Mean blood pressure 230

Mediastino-pericarditis *See* Pericardial adhesion

Methaemoglobinæmia 36

Mitral regurgitation 136

Diagnosis of 139

Grades of 139

Prognosis of 146

Mitral stenosis 130

Auricular fibrillation and 205

Early diagnosis of 132 199

Infarction and 170

Murmurs in 131 134

Recognition of 130

Rheumatic fever and 199

Surgical treatment of 148

Third heart sound in 199

Mitral valve *See also* Mitral stenosis and Mitral regurgitation

Disease recognised in 139

Pathological anatomy of 129

Murmur—

Auricular fibrillation changing 134

Cardiorespiratory 136

Constant and inconstant 137

Continuous basal 262

Crescendo in mitral stenosis 131 134

Diastolic of mitral stenosis 131 134

Diastolic of tricuspid stenosis 141

Early diastolic aortic 123 124

Early diastolic mitral 132 134

Early diastolic pulmonary 124

Exocardial 137

Flint's 136

Mid diastolic mitral 132 136 199

Mitral systolic 137

Musical 125 139

Posture and 123 132

Presystolic 132

Steel's 124

Systolic aortic 123 128

Systolic apical 136

Systolic apical (dilatation and) 112 139 197 245

Systolic basal in coarctation aorta, 264

Systolic (exercise inducing) 158

Systolic in effort syndrome 160

Systolic of defective septum 264

Systolic pulmonary 266

Systolic tricuspid 141

Timing of 123 131

To and fro aortic 123

Muscular ischaemia and angina 42

Musical murmur 125 139

Mycotic aneurysm 188

Myocardial weakness 251

Myocarditis 250

Myocardium (signs of involvement) 245

Myomalacia cordis 44

Nausea—

From congestion 88

From digitalis 87

Syncope and 93

Neosalvarsan 226

Neptitis *See* Renal disease

Neurasthenia (anginal pain and) 59

Nitrites in angina 54 57

Nitroglycerine 58

Nodes (digital) 188

Nodules (subcutaneous) 194 196 199

Nodules (subcutaneous cardiac) 196

Normal heart (recognition of) 277

Nose bleeding 228

Nothnagel's syndrome 60

Novasurol 30

Obesity (fatty heart and) 250

- Obesity (hypertension and) 227 238
- Oedema 18
- Cardiac 18
- Differential diagnosis of 20
- Distribution of 19
- Drainage of 31
- Factors in causation of 18
- In elderly people 20 229
- In hypertension 223 233
- Of lung *See* Pulmonary oedema
- Posture and 19
- Recognition of 19
- Serous exudates 21 III 179
- Treatment of 30
- Oesophageal obstruction 213 217
- Operations in heart cases 271
- Orthodiagraph 107
- Overacting heart 70 258
- Oxygen therapy in cardiac failure 29
- Pain *See also* Anginal pain
- Aneurysm and 213 220
- Cardiac failure and 7 29
- Coronary thrombosis and 44
- Effort syndrome and 160
- Hepatic 14 29
- Pericarditis and 173 176
- Precordial 20 160
- Syphilitic aortitis and 213
- Pallor 35
- Acute bacterial endocarditis and 192
- Aortic regurgitation and 126 189
- Cause of 35
- Rheumatic fever and 196
- Subacute bacterial endocarditis and 186 189
- Syncope and 99
- Palpitation—
- Aortic regurgitation and 121
- Auricular fibrillation and 82
- Effort syndrome and 159
- Extrasystolic 66 68
- Paroxysmal tachycardia and 72 73
- Simple form 70 78
- Paraplegia (aneurysm and) 290
- Paroxysmal fibrillation 82 258 259
- Paroxysmal hypertension 60 168
- Paroxysmal tachycardia 71
- Abrupt onset and offset 71 74
- Anginal pain in 60 73
- Dilatation of heart in 72 153
- Exercise provoking 153
- Prognosis of 70
- Residual damage from attacks 77
- Rheumatic heart disease and 206
- Treatment of 78
- Patent foramen ovale 264
- Pericardial adhesion and mediastino-pericarditis 176
- Cardiac failure and 22
- Diagnosis of 181
- Hypertrophy and 106
- Rheumatic fever and 195 203
- Significance of 181
- Symptoms and signs of 177
- Treatment of 182
- Pericardial effusion 172
- Pericardial friction 47 174 177
- Pericardial sac (congenital absence of) 177
- Pericarditis 172
- Constrictive 177 204
- Course of 175
- Friction in 47 174 177
- Rheumatic 195 197
- Symptoms and signs of 173
- Treatment of 175
- With effusion 172
- Pericardium (dilatation guarded by) 117 153 172
- Perihepatitis (ascites and) 21 170
- Persistent ductus arteriosus 124 262
- Petoehiae 186 192
- Pistol shot sound 122
- Pleuropericardial friction 175 180
- Polycythaemia 39 263
- Posture—
- Giddiness and syncope and 97 102 121
- In cardiac failure 26
- Murmurs and 123 132
- Venous pulse and 13
- Pregnancy and the heart 153 268
- Presystolic murmur 132
- Prognosis (*see various diseases also*)—
- Basis of 277
- Enlargement and 117
- Experience the sound basis of 67 144
- Grouping for 207 278
- Principle of 67 144
- Pulmonary artery dilated 166 263 266
- Pulmonary congestion 165
- Acute 166
- Cyanosis in 38
- Passive 165
- Signs and symptoms of 15 163
- Pulmonary disease (cardiac failure and) 22
- Pulmonary infarction 170 255
- Pulmonary oedema 163
- Acute 166 169
- Coronary thrombosis and 45
- Cyanosis in 38
- Early 165
- Prognosis and treatment of 169

I *limbary oedema*—(contd.)
 Rheumatic disease and 207
 Symptoms and signs of 15 165
 Syphilis and 225
Pulmonary regurgitation 124 206
Pulmonary stenosis 265
Pulsating tumour 214
Illusion of chest wall 213 *See also*
Lurid impulse
Pulse (arterial)—
 Injurious decline of 179
 Intermittent 63
 Irregular *See* Irregular and disordered
 heart action
 ~ ~ 101 231

block
 Slowly and syncope 99
 Strength and myocardium 245
 Tension of 229
 Unequal (right and left) 220
 Venous pulse distinguished 14
Pulse alternans 232 248
Pulsus paradoxus 180
Purpura—
 Acute bacterial endocarditis and 192
 Hypertension and 232
 Rheumatic fever and 198
 Subacute bacterial endocarditis and
 187

Quinine—
 Auricular fibrillation and 88 90 208
 Flitter and 80
 Intravenous 79
 Paroxysmal tachycardia and 79

Rapid pulse *See* Tachycardia
Recession of the heart 254
Recurrent laryngeal nerve (aneurysm
 and) 218

Red granular kidney 228

Reduplicated sounds—
 First apical 130 134
 In complete block 95
 Second apical 247
 Second pulmonary 130 134

Renal asthma 166

Renal disease—
 Arteriosclerotic (and arterio-sclerotic)
 kidney 243
 Bacterial endocarditis and 187
 Chronic glomerulo-nephritis, 234
 Congested kidney 15
 Re-granular kidney 228 243
Renal efficiency in hypertension 232

Renal embolism 189
Retinal arteries 231
Retinal embolism 188
Retracted lung 109 178 213
Rheumatic earitis 194
Rheumatic fever 191
Rheumatic heart disease (chronic) 202
Rheumatic nodules 104 196 194
Rheumatic valvulitis 120 129, 195 202
Rib movement and prominence 109
Rowing and the heart 154
Ruptured aortic cusp 125 152 222
Ruptured ventricle 46 48 103

Senile heart 244
Serous exudates 21 III 179
Simple bradycardia 92
Simple tachycardia 69 76 78
Sinus arrhythmia, 62
Skin colour 35
Skin temperature and cyanosis 30
 sleeplessness 27
Slow pulse *See* Bradycardia
Slow rising pulse 127
Sold or stertor 158
Southey's tubes, 31
Sphygmomanometry 230
Splenic embolism 188
Splenic enlargement 186 189
Status anginosus 47
Steell's murmur 124
Sternum moving 109
Strenuous acts 285
Stridor in aneurysm 217
Strophantus 89
Subacute bacterial endocarditis 183
Sudden death 103
 Angina pectoris and 55 56
 Auricular fibrillation and 87 104
 Ball thrombus and 206
 Coronary thrombosis and 44 48 103
 Doctor's references to 282
 Fatty heart and 250
 Paroxysmal tachycardia and 73
 Vagal inhibition and 104
 Ventricular fibrillation and 103 274
 Superior vena cava obstructed 16 179
 219
Sweating (syncope and) 99 101
Sympathetic (aneurysm pressing on) 218
Syncope 96
Symphylis (cardio-aortic) 211 221 *See*
also Aortic regurgitation
Systolic blood pressure 121 230 *See*
also Hypertension
 Aortic valve disease and 121 127
 Higher in leg than arm 122

- Systolic murmur *See* Murmurs
- Systolic retractions with pericardial adhesions 177
- Tachycardia 69 *See also* Simple tachycardia and Paroxysmal Tachycardia and Auricular flutter
- Anginal pain and 60
- Aortic regurgitation and 125
- Diastole shortened in 24
- Differentiation of 73
- Effect on sound and impulse 70
- Energy expenditure and 72
- Irregular *See* Auricular fibrillation
- Syncope and 76 101
- Thyrotoxic state and 259
- Tapping serous effusions 31 176
- Test exercises 4
- Theobromine 30 58
- Theocin 30
- Third heart sound 199 247
- Thoracotomy for pericardial adhesions 182
- Thrill—
- Aortic stenosis and 128
- Congenital affections and 262 264 266
- Mitral stenosis and 130 131
- Thrombotic accidents in rheumatic heart disease 208
- Thyrotoxic state 257
- Timing murmurs 123 131
- Tobacco 59 284
- Tortuosity of vessels 241
- Trachea (aneurysm pressing on) 17
- Tracheal tug 215
- Treatment (*see various conditions and symptoms also*) 283
- Tricuspid regurgitation 141
- Tricuspid stenosis 130 141
- Tricuspid valve 130 141
- T wave inverted 247
- Unexpanded infundibulum 265
- Uræ—
- Cardiac failure and 15
- Flow in cardiac failure 30
- Hypertension and 239
- Vagal inhibition (sudden death and) 104
- Vagal standstill 100 104
- Valve disease (*see various valve defects also*)—
- Anæsthetics and 271
- Burden of 106
- Cardiac failure and 29
- Exercise and 149
- Management of 148
- Prognosis of 143 209
- Rheumatic heart disease and 209 207
- Significance of 143
- Vasoconstriction (symptoms of) 60
- Vasodilatation 40
- Aortic regurgitation and 125
- Thyrotoxic state and 257
- Vasovagal syncope 98 102 159
- Venæsection 28
- Venous congestion 7 8
- Breathlessness and 10 33 33
- Cyanosis and 39
- Diagnosis of 15
- Liver in 14
- Pericardial disease and 173 179
- Superior caval obstruction and 16 219
- Venous pressure measurement 8 11 12
- Venous pulse—
- Arterial pulse distinguished 14
- Congestion and 13
- Normal 19
- Posture and 13
- Venous pressure and 12
- Venous swelling and collapse 9
- Ventricular arrest 99 103
- Ventricular fibrillation 103
- Coronary thrombosis and 44
- Sudden death from 103 274
- Syncope from 101
- Vomiting (from congestion or digitalis) 8 87 89
- Water hammer pulse 121 174
- Work—
- Burden of 151
- Cardiac failure and 29 151
- Prolonged and failure 154
- X ray—
- Aneurysm and dilatation and 216
- Cardiac enlargement and 107 175
- Orthodagraph 107
- Pericardial adhesions and 151
- Pericardial effusion and 175

THE END

